


## Gold(I)-Catalyzed Inter- and Intramolecular Additions of Carbonyl Compounds to Allenenes

Tania Jiménez,<sup>†</sup> Javier Carreras,<sup>†</sup> Julien Ceccon,<sup>†</sup> and Antonio M. Echavarren<sup>\*,†,‡</sup><sup>†</sup>Institute of Chemical Research of Catalonia (ICIQ), Barcelona Institute of Science and Technology, Av Paisos Catalans 16, 43007 Tarragona, Spain<sup>‡</sup>Departament de Química Analítica i Química Orgànica, Universitat Rovira i Virgili, C/Marcel·li Domingo s/n, 43007 Tarragona, Spain


## Supporting Information

**ABSTRACT:** The gold(I)-catalyzed intramolecular reaction of allenenes with oxoalkenes leads to bicyclo[6.3.0]undecane ring systems, although in the case of terminally disubstituted allenenes, seven-membered rings are formed. The related intermolecular addition of aldehydes to allenenes also gives seven-membered rings.




Gold(I)-catalyzed enyne cycloisomerization reactions are powerful tools for the stereoselective construction of complex carbon skeletons.<sup>1</sup> These transformations have been used as the key steps in the total synthesis of diverse natural products.<sup>2,3</sup> We have developed a particularly useful transformation based on the gold(I)-catalyzed [2 + 2 + 2] intramolecular cycloaddition between oxo-1,6-enynes in substrates such as **1**, which leads to oxatricyclic derivatives **2a,b** by a cascade process in which two C–C and one C–O bonds are formed.<sup>4</sup> This methodology was applied in the total syntheses of (+)-orientalol **3a** and (–)-englerin **A**<sup>3b,5</sup> (Scheme 1). A mechanistically similar transformation was also developed with oxo-1,5-enynes.<sup>6</sup>

## Scheme 1. Gold(I)-Catalyzed [2 + 2 + 2] Intramolecular Cycloaddition between Alkynes and Oxoalkenes



The intermolecular reaction of terminal alkynes with 5-oxoalkenes also gives [3.2.1]oxabicycles.<sup>7</sup> A somewhat analogous intermolecular gold(I)-catalyzed reaction of 5-, 6-, and 7-oxoalkenes with allenamides was developed by the group of Mascareñas to form 7–9-membered rings depending on the length of the tether of the oxoalkenes.<sup>8a</sup> In addition, by use of chiral ligands, this transformation was further developed as an enantioselective process. The same group recently reported the

gold(I)-catalyzed [2 + 2 + 2] cycloaddition of allenamides, alkenes, and aldehydes for the synthesis of functionalized tetrahydropyrans.<sup>8b,9</sup> However, similar transformations involving simple oxo-1,7-allenenes have not been described. Herein, as part of a program on the development of new gold(I)-catalyzed cascade reactions for the synthesis of complex sesquiterpenes,<sup>3,10</sup> we report a ready access to the bicyclo[6.3.0]undecane ring system,<sup>11</sup> a motif conspicuously present among sesquiterpenes and higher terpene natural products (Figure 1).<sup>12</sup>




**Figure 1.** Representative natural terpenoids containing eight-membered rings.

We envisioned that in oxo-1,7-allenenes or 1,7-allenenes reacting intermolecularly with carbonyl compounds the attack of the alkene to a gold(I)-activated allene **I** would generate tertiary carbocation **II**, which would be rapidly trapped by the carbonyl group to form oxonium cation **III** (Scheme 2). Prins cyclization would then form intermediates **IV** or **V** by *endo*- or *exo*-pathways, respectively. Gold(I) carbene **IV** could give rise

**Received:** February 2, 2016

**Published:** February 26, 2016

Scheme 2. Proposed Mechanism for the Gold-Catalyzed Reaction of 11-Oxo-1,7-allenenes



to eight-membered ring compounds VI or VII when  $R^2 = H$ , whereas V would lead to seven-membered ring VIII. In addition to the regiochemical uncertainties, control of the final relative configuration was a second matter of concern. It is important to remember that unlike the reaction of the analogous 1,6-enynes **1**, where the cyclization proceeds stereospecifically through cyclopropyl gold(I) carbenes as intermediates,<sup>4,13</sup> reactions of allenes with alkenes are fundamentally different since the configuration of the final products is not mechanistically determined. Thus, the initial cyclization could give *cis*- or *trans*-five-membered rings II, and furthermore, new stereocenters are generated in the formation of intermediate III from tertiary carbocation II as well as in the Prins cyclization leading to IV.


Initially, we investigated the intramolecular reactivity of 11-oxo-1,7-allenenes using allene aldehyde **3a** as the substrate in the presence of different catalysts (5 mol %) at room temperature in  $\text{CH}_2\text{Cl}_2$  (Table 1). In all the cases, we obtained tricyclic derivative **4a** as a result of a cascade reaction proceeding selectively through gold(I) carbene IV. The best results were obtained with gold complexes **A**<sub>1</sub>, **A**<sub>3</sub>, and **D** bearing NHC and phosphite ligands (Table 1, entries 1, 3, and 7). Lower yields were obtained with gold(I) complexes **B** and **C** with bulky biphenyl phosphines<sup>14</sup> (Table 1, entries 4–6). Best results were obtained with *i*Pr–gold(I) cationic complex **A**<sub>1</sub> with  $\text{SbF}_6^-$  as the counteranion, leading to the formation of oxatricyclic derivative **4a** in 71% isolated yield as a single diastereomer (Table 1, entry 1). The reaction could also be carried out with  $\text{PtCl}_2$ , although the yield was poor (Table 1, entry 8). The relative configuration was confirmed by determining the X-ray crystal structure of diol **4a'**,<sup>15</sup> prepared by reduction of **4a** with  $\text{LiAlH}_4$ .

Having the optimal conditions in hands, we sought to evaluate the generality of the reaction, considering the substitution pattern in the allene, the configuration of the alkene, and also the intermolecular reaction between allenes and aldehydes (Scheme 3). By performing the reaction with the monosubstituted allene **3b**, we observed the same type of reactivity, leading in this case to a mixture of diastereomers.

Table 1. Catalyst Optimization in the Intramolecular Cycloaddition of 11-Oxo-1,7-allenene **3a**<sup>a</sup>


entry	catalyst	time (h)	yield <sup>b</sup> (%)
1	<b>A</b> <sub>1</sub>	0.25	78 (71) <sup>c</sup>
2	<b>A</b> <sub>2</sub>	0.25	45
3	<b>A</b> <sub>3</sub> + $\text{AgSbF}_6$	0.25	63
4	<b>B</b>	6	37
5	<b>C</b> <sub>1</sub>	24	25
6	<b>C</b> <sub>2</sub> + $\text{AgSbF}_6$	24	55
7	<b>D</b>	2 min	67
8	$\text{PtCl}_2$	16	10

<sup>a</sup>Conditions: allene **3a** and catalyst (5 mol %) in  $\text{CH}_2\text{Cl}_2$  (0.1 M).  
<sup>b</sup>Yield of **4a** determined by <sup>1</sup>H NMR; see the Supporting Information for details.  
<sup>c</sup>Isolated yield.



Catalysts **A**<sub>1</sub> or **A**<sub>2</sub> led to a 3:1 mixture of diastereomers **4b** in moderate yields (42–46%) after 72 h, whereas more electrophilic catalyst **D** gave **4b** as a 1:1 mixture of diastereomers after just 10 min.<sup>16</sup> Surprisingly, reaction of trisubstituted allene **3c** with catalysts **A**<sub>1</sub> gave seven-membered ring **5** in 78% yield instead of an eight-membered cyclic derivative with a *trans* configuration at the fusion between the 5- and the 7-membered rings, which was confirmed by X-ray crystallographic analysis.<sup>15</sup> A similar reaction was observed with phosphite gold complex **D** (88%, 2 min); however, a separable 3:1 mixture of diastereomers was obtained with this catalyst. The lower stereoselectivity observed using catalyst **D** is probably a consequence of the very high electrophilicity of this gold(I) complex, which favors formation of both intermediates *cis*- and *trans*-II through early transition states of closely similar energy. Reaction of substrate **3d** with a *Z* configuration at the alkene led to eight-membered ring **4d** in 40–46% yield using catalysts **A**<sub>1</sub> or **D**. Compound **4d**, which was also observed as a minor product in the cyclization of **3a**,<sup>16</sup> could arise by a proton elimination from an intermediate IV with a configuration different from that involved in the formation of **4a**.<sup>17</sup> We also investigated the intermolecular version of this reaction using 1,7-allenene **6** and excess aliphatic or aromatic aldehydes (Scheme 3). In the presence of phosphite gold(I) catalyst **D**, fast reactions were observed (5–10 min), yielding hexahydro-1*H*-cyclopenta[*c*]oxepines **7a–i** in moderate to good yields as mixture of diastereomers as a result of a cyclization/*endo*-Prins process. The use of complex **A**<sub>1</sub> moderately increased the

Scheme 3. Gold(I)-Catalyzed Intra- and Intermolecular Cycloaddition of 11-Oxo-1,7-allenenes



diastereoselectivity, although the yields were lower and longer reaction times were required.<sup>16</sup>

The cyclization of 11-oxo-1,7-allenenes was extended to ketones instead of aldehydes (Table 2). Under the optimized conditions used for aldehyde 3a, we were pleased to observe the formation of the tricyclic compounds 4e–k in 60–87% yields using gold complex A<sub>1</sub>. X-ray crystallographic analysis of 4g unambiguously confirmed its relative configuration.<sup>15</sup> Alkyl

Table 2. Gold(I)-Catalyzed Intramolecular Cycloaddition of 11-Oxo-1,7-allenenes 3e–k<sup>a</sup>

entry	R	time (h)	product (yield, %) <sup>b</sup>
1	Me (3e)	6	4e (63)
2	<i>n</i> -Hex (3f)	0.25	4f (64)
3	Cy (3g)	0.15	4g (72)
4	<i>t</i> -Bu (3h)	3	4h (79)
5	Ph (3i)	6	4i (72)
6	3,5-(CF <sub>3</sub> ) <sub>2</sub> C <sub>6</sub> H <sub>3</sub> (3j)	2	4j (87)
7	CH <sub>2</sub> Cl (3k)	1	4k (60)


<sup>a</sup>Conditions: allenene 3e–k and A<sub>1</sub> (5 mol %) in CH<sub>2</sub>Cl<sub>2</sub> (0.1 M).

<sup>b</sup>Isolated yields.

ketones, including those with bulky groups, and aryl ketones react similarly. Interestingly, oxoallenenes with electron-withdrawing groups at the ketone, such as 3j and 3k (Table 2, entries 6 and 7), reacted smoothly and gave yields similar to those obtained from other ketones, which supports the hypothesis that formation of carbocation II is rate-determining and attack of the carbonyl group is, comparatively, a faster process.

Finally, we applied this intramolecular cycloaddition for the enantioselective formation of an oxatricyclic system. The synthesis of the oxoallenene began by alkylation of methyl isobutyrate with geranyl bromide to give ester 8,<sup>18</sup> which was converted into known aldehyde 9<sup>19</sup> in two steps by reduction with LiAlH<sub>4</sub> and subsequent oxidation with Dess–Martin periodinone (Scheme 4). Enantioselective allenylation of

Scheme 4. Enantioselective Synthesis of Oxatricyclic Compound 12



aldehyde 9 was performed by applying Corey's method<sup>20</sup> using a chiral bromoborane and propadienyltri-*n*-butylstannane followed by benzylation under standard conditions to yield allene 10. Treatment of 10 with Admix- $\alpha$  and methylsulfonamide followed by oxidative cleavage using NaIO<sub>4</sub> on silica<sup>21</sup> led to aldehyde 11. At this point, the excellent enantioselectivity of the allenylation reaction was confirmed by chiral GC (98:2 er). Exposing 11 to cationic gold(I) complex A<sub>1</sub> for 16 h at 25 °C gave 12 in 75% yield as the only isolated tricyclic compound. The intermolecular reaction of 11 occurred with complete retention of the configuration.

In summary, we have found that the intramolecular reaction of 11-oxo-1,7-allenenes gives rise to bicyclo[6.3.0]undecane ring systems, usually in good yields and selectivities, which is remarkable for a reaction that most probably proceeds through an open carbocation. Only in the case of 1,1-dialkyl-substituted alkenes are seven-membered ring systems formed as a consequence of an *exo*-selective Prins-type cyclization. This

reaction can also be carried out intermolecularly between 1,7-allenenes with aliphatic or aromatic aldehydes.

## ■ ASSOCIATED CONTENT

### ■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.6b00342](https://doi.org/10.1021/acs.orglett.6b00342).

Experimental procedures and characterization data for compounds as well as the X-ray crystal structures of **4a'**, **5**, and **4g** (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [aecharvarren@iciq.es](mailto:aecharvarren@iciq.es).

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank MINECO (Severo Ochoa Excellence Accreditation 2014-2018 (SEV-2013-0319), Project CTQ2013-42106-P), the European Research Council (Advanced Grant No. 321066), the AGAUR (2014 SGR 818 and Beatriu de Pinós Postdoctoral Fellowship to J.C.), the ICIQ X-ray Diffraction and Chromatography units, and the ICIQ Foundation.

## ■ REFERENCES

- (1) (a) Zhang, L.; Sun, J.; Kozmin, S. A. *Adv. Synth. Catal.* **2006**, *348*, 2271–2296. (b) Fürstner, A.; Davies, P. W. *Angew. Chem., Int. Ed.* **2007**, *46*, 3410–3449. (c) Hashmi, A. S. K. *Chem. Rev.* **2007**, *107*, 3180–3211. (d) Li, Z.; Brouwer, C.; He, C. *Chem. Rev.* **2008**, *108*, 3239–3265. (e) Arcadi, A. *Chem. Rev.* **2008**, *108*, 3266–3325. (f) Jiménez-Núñez, E.; Echavarren, A. M. *Chem. Rev.* **2008**, *108*, 3326–3350. (g) Gorin, D. J.; Sherry, B. D.; Toste, F. D. *Chem. Rev.* **2008**, *108*, 3351–3378. (h) Michelet, V.; Toullec, P. Y.; Genêt, J.-P. *Angew. Chem., Int. Ed.* **2008**, *47*, 4268–4315. (i) Aubert, C.; Fensterbank, L.; Garcia, P.; Malacria, M.; Simonneau, A. *Chem. Rev.* **2011**, *111*, 1954–1993. (j) Krause, N.; Winter, C. *Chem. Rev.* **2011**, *111*, 1994–2009. (k) Obradors, C.; Echavarren, A. M. *Acc. Chem. Res.* **2014**, *47*, 902–912. (l) Fensterbank, L.; Malacria, M. *Acc. Chem. Res.* **2014**, *47*, 953–965. (m) Dorel, R.; Echavarren, A. M. *Chem. Rev.* **2015**, *115*, 9028–9072.
- (2) (a) Hashmi, A. S. K.; Rudolph, M. *Chem. Soc. Rev.* **2008**, *37*, 1766–1775. (b) Rudolph, M.; Hashmi, A. S. K. *Chem. Commun.* **2011**, *47*, 6536–6544. (c) Rudolph, M.; Hashmi, A. S. K. *Chem. Soc. Rev.* **2012**, *41*, 2448–2462. (d) Fürstner, A. *Acc. Chem. Res.* **2014**, *47*, 925–938. (e) Zhang, Y.; Luo, T.; Yang, Z. *Nat. Prod. Rep.* **2014**, *31*, 489–503.
- (3) Work from our group on the total synthesis of terpenoids by using gold(I) catalysis: (a) Jiménez-Núñez, E.; Molawi, K.; Echavarren, A. M. *Chem. Commun.* **2009**, 7327–7329. (b) Molawi, K.; Delpont, N.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2010**, *49*, 3517–3519. (c) Gaydou, M.; Miller, R. E.; Delpont, N.; Ceccon, J.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2013**, *52*, 6396–6399. (d) Carreras, J.; Livendahl, M.; McGonigal, P. R.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2014**, *53*, 4896–4899.
- (4) Jiménez-Núñez, E.; Claverie, C. K.; Nieto-Oberhuber, C.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2006**, *45*, 5452–5455.
- (5) Zhou, Q.; Chen, X.; Ma, D. *Angew. Chem., Int. Ed.* **2010**, *49*, 3513–3516.
- (6) Huguet, N.; Echavarren, A. M. *Synlett* **2012**, *2012*, 49–53.
- (7) Obradors, C.; Echavarren, A. M. *Chem. - Eur. J.* **2013**, *19*, 3547–3551.
- (8) (a) Faustino, H.; Alonso, I.; Mascareñas, J. L. *Angew. Chem., Int. Ed.* **2013**, *52*, 6526–6530. (b) Faustino, H.; Varela, I.; Mascareñas, J. L.; López, F. *Chem. Sci.* **2015**, *6*, 2903–2908.
- (9) Review on catalytic [4 + 2] and [4 + 3] cycloaddition of allenes: López, F.; Mascareñas, J. L. *Chem. Soc. Rev.* **2014**, *43*, 2904–2912.
- (10) Ardckhean, R.; Caputo, D. F. J.; Morrow, S. M.; Shi, H.; Xiong, Y.; Anderson, E. A. *Chem. Soc. Rev.* **2016**, DOI: [10.1039/C5CS00105F](https://doi.org/10.1039/C5CS00105F).
- (11) Leading references on the formation of eight-membered rings by cycloaddition: (a) Yu, Z.-X.; Wang, Y.; Wang, Y. *Chem. - Asian J.* **2010**, *5*, 1072–1088. (b) Wang, Y.; Yu, Z.-X. *Acc. Chem. Res.* **2015**, *48*, 2288–2296. (c) Wender, P. A.; Ihle, N. C. *J. Am. Chem. Soc.* **1986**, *108*, 4678–4679.
- (12) For examples, see the following. (a) Dumortienol: Toyota, M.; Bardon, A.; Kamiya, N.; Takaoka, S.; Asakawa, Y. *Chem. Pharm. Bull.* **1997**, *45*, 2119–2121. (b) Toxicodenane C: He, J.-B.; Luo, J.; Zhang, L.; Yan, Y.-M. *Org. Lett.* **2013**, *15*, 3602–3605. (c) Astericanolide: Limanto, J.; Snapper, M. L. *J. Am. Chem. Soc.* **2000**, *122*, 8071–8072. (d) Ophiobolin F: Nozoe, S.; Morisaki, M.; Fukushima, K.; Okuda, S. *Tetrahedron Lett.* **1968**, *9*, 4457–4458. (e) Nitidasine: Hog, D. T.; Huber, F. M. E.; Mayer, P.; Trauner, D. *Angew. Chem., Int. Ed.* **2014**, *53*, 8513–8517.
- (13) Dorel, R.; Echavarren, A. M. *J. Org. Chem.* **2015**, *80*, 7321–7332.
- (14) Ranieri, B.; Escofet, I.; Echavarren, A. M. *Org. Biomol. Chem.* **2015**, *13*, 7103–7118.
- (15) CCDC 1451024 (**4a'**), CCDC 1451022 (**5**), and CCDC 1451023 (**4g**) contain the supplementary crystallographic data. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- (16) See the Supporting Information for more details.
- (17) An isomerization of **3a** into **3d** by a gold(I)- or proton-catalyzed process seems unlikely since **3a** was calculated to be 3 kcal·mol<sup>-1</sup> more stable than **3d** (DFT calculations, B3LYP/6-31G(d), solvent = CH<sub>2</sub>Cl<sub>2</sub>).
- (18) Kodama, M.; Shiobara, Y.; Sumitomo, H.; Mitani, K.; Ueno, K. *Chem. Pharm. Bull.* **1987**, *35*, 4039–4042.
- (19) Negishi, E.; John, R. A. *J. Org. Chem.* **1983**, *48*, 4098–4012.
- (20) (a) Corey, E. J.; Imwinkelried, R.; Pikul, S.; Xiang, Y. B. *J. Am. Chem. Soc.* **1989**, *111*, 5493–5495. (b) Corey, E. J.; Yu, C.-M.; Lee, D.-H. *J. Am. Chem. Soc.* **1990**, *112*, 878–879.
- (21) Zhong, Y. L.; Shing, T. K. M. *J. Org. Chem.* **1997**, *62*, 2622–2624.