

Welcome Universitaetsbibliothek Johann Christian Senckenberg

(Log on / register)

BioMed Central home | Journals A-Z | Feedback | Support | My details

home | issues | articles | reports | supplements | weblinks | conferences | search | information | subscribe | my details

This article is part of the supplement: [Sepsis 2009](#) .

Poster presentation

## T-cell-specific peroxisome proliferator-activated receptor gamma depletion inhibits T-cell apoptosis and improves survival of septic mice via an IL-2-dependent mechanism

MV Schmidt<sup>1</sup>, P Paulus<sup>2</sup>, A-M Kuhn<sup>1</sup>, V Meilladec-Jullig<sup>1</sup>, K Zacharowski<sup>2</sup>, B Bruene<sup>1</sup> and A von Knethen<sup>1</sup>

1 Institute of Biochemistry I, Faculty of Medicine, Goethe-University Frankfurt, Germany

2 Department of Anaesthesia, Intensive Care Medicine & Pain Therapy, University Hospital Frankfurt, Germany

■ corresponding author email

from Sepsis 2009

Amsterdam, The Netherlands. 11-14 November 2009

Critical Care 2009, **13**(Suppl 4):P37 doi:10.1186/cc8093

The electronic version of this abstract is the complete one and can be found online at:

<http://ccforum.com/content/13/S4/P37>

Published: 11 November 2009

© 2009 BioMed Central Ltd.

### Introduction

Immune paralysis with massive T-cell apoptosis is a central pathogenic event during sepsis and correlates with septic patient mortality. Previous observations implied a crucial role of peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) during T-cell apoptosis.

### Methods

To elucidate mechanisms of PPAR $\gamma$ -induced T-cell depletion, we used an endotoxin model as well as the caecal ligation and puncture sepsis model to imitate septic conditions in wild-type versus conditional PPAR $\gamma$  knockout (KO) mice.

### Results

PPAR $\gamma$  KO mice showed a marked survival advantage compared with control mice. Their T cells were substantially protected against sepsis-induced death and showed a significantly higher expression of the pro-survival factor IL-2. Since PPAR $\gamma$  is described to repress nuclear factor of activated T cells (NFAT) transactivation and concomitant IL-2 expression, we propose inhibition of NFAT as the underlying mechanism allowing T-cell apoptosis. Corroborating our hypothesis, we observed up-regulation of the pro-apoptotic protein

#### Critical Care

Volume 13

Suppl 4

#### Viewing options:

Full text

PDF (1.9MB)

#### Related literature:

Other articles by authors

[on Google Scholar](#)

[on PubMed](#)

Related articles/pages

[on Google](#)

[on Google Scholar](#)

#### Tools:

[Download citation\(s\)](#)

[Download XML](#)

[Email to a friend](#)

[Order reprints](#)

[Post a comment](#)

#### ■ Nominate for award

#### Post to:

[Citeulike](#)

[Connotea](#)

[Del.icio.us](#)

[Facebook](#)

[Twitter](#)

BIM and downregulation of the anti-apoptotic protein Bcl-2 in control mice, which are downstream effector proteins of IL-2 receptor signaling. Application of a neutralizing anti-IL-2 antibody reversed the pro-survival effect of PPAR $\gamma$ -deficient T cells and confirmed IL-2-dependent apoptosis during sepsis.

## Conclusion

---

Apparently antagonizing PPAR $\gamma$  in T cells might improve their survival during sepsis, which concomitantly enhances defence mechanisms and possibly provokes an increased survival of septic patients.

[Have something to say? Post a comment on this article!](#)

[Terms and Conditions](#) [Privacy statement](#) [Information for advertisers](#) [Jobs at BMC](#) [Contact us](#)

© 1999-2009 BioMed Central Ltd unless otherwise stated. Part of Springer Science+Business Media.