

University of Groningen

In situ impact of solar ultraviolet radiation on photosynthesis and DNA in temperate marine phytoplankton

Helbling, E.W.; Buma, A.G.J.; de Boer, M.K.; Villafane, V

Published in:
Marine Ecology Progress Series

DOI:
[10.3354/meps211043](https://doi.org/10.3354/meps211043)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2001

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Helbling, E. W., Buma, A. G. J., de Boer, M. K., & Villafane, V. (2001). In situ impact of solar ultraviolet radiation on photosynthesis and DNA in temperate marine phytoplankton. *Marine Ecology Progress Series*, 211, 43-49. <https://doi.org/10.3354/meps211043>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

In situ impact of solar ultraviolet radiation on photosynthesis and DNA in temperate marine phytoplankton

E. Walter Helbling^{1,*}, Anita G. J. Buma², M. Karin de Boer², Virginia E. Villafañe¹

¹Estación de Fotobiología Playa Unión, Casilla de Correos No. 153 – (9100) Trelew, Chubut, Argentina and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)

²Department of Marine Biology, University of Groningen, PO Box 14, 9750 AA Haren, The Netherlands

ABSTRACT: *In situ* experiments were conducted at various depths in the water column to determine the impact of solar UV radiation (280 to 400 nm) upon photosynthesis and DNA of natural phytoplankton assemblages from mid-latitudes of Patagonia (Bahía Bustamante, Chubut, Argentina; 45° S, 66.5° W). The effects of UV radiation were significant at the surface; however, the impact decreased rapidly with depth: at 3 m there was no measurable DNA damage accumulation, whereas at 6 m photosynthetic inhibition was almost zero. UV-A radiation (315 to 400 nm) was mostly responsible for photosynthetic inhibition, while UV-B radiation (280 to 315 nm) had a lesser effect on this process. However, UV-B radiation was very effective in damaging the DNA through the formation of cyclobutane pyrimidine dimers (CPDs) in surface waters. The high initial CPD level found in the natural phytoplankton assemblage decreased when samples were incubated at 3 or 6 m, indicating that at these depths repair, dilution or disappearance of damage occurred. Phytoplankton assemblages were dominated by cells less than 2 µm in effective diameter; this cell size category seems to be more resistant to photosynthetic inhibition, but vulnerable to CPD accumulation, as compared with larger eukaryotic phytoplankters (i.e., *Phaeodactylum* sp.).

KEY WORDS: UV · Phytoplankton · DNA · CPD · Thymine dimers · Photosynthesis inhibition · Mid-latitudes

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Phytoplanktonic organisms are affected in different ways by ambient levels of solar radiation (Holm-Hansen et al. 1993a, Häder 1997). In particular, the effects of solar ultraviolet radiation (UVR, 280 to 400 nm) in autotrophic organisms have been addressed due to the discovery of the stratospheric ozone depletion (i.e., the ozone 'hole'), as this phenomenon results in an increase of short wavelengths of UV-B radiation (280 to 315 nm) reaching the earth's surface (Madronich 1993). One of the most studied effects of solar UVR upon

phytoplanktonic organisms is the inhibition of photosynthetic rates, which has been observed in many regions, such as polar areas (Helbling et al. 1992, Smith et al. 1992, Neale et al. 1998), and temperate (Behrenfeld et al. 1993, Helbling et al. 1993) and tropical environments (Helbling et al. 1992, Behrenfeld et al. 1993, Villafañe et al. 1999). Another effect of UVR that has been particularly addressed in phytoplanktonic organisms is the damage of genetic material (i.e., DNA) as a consequence of the formation of cyclobutane pyrimidine dimers (CPDs), mainly thymine dimers – T<>T (Karentz et al. 1991, Buma et al. 1997, Boelen et al. 1999, unpubl.). Most of the UVR effect studies have been carried out in polar areas (especially Antarctica), as it is thought that organisms of these high latitudes would be

*E-mail: efpu@efpu.com.ar

especially affected by enhanced levels of solar UV-B radiation during springtime ozone depletion events (Smith et al. 1992, Holm-Hansen et al. 1993b, Arrigo 1994). However, as responses to UVR depend on the particular biological, physical and optical characteristics of the ecosystem under study, it is not possible to safely extrapolate the results obtained in polar areas to other aquatic systems of the planet.

There are not many studies that address the impact of solar UVR upon phytoplanktonic species of temperate systems (Behrenfeld et al. 1993, Helbling et al. 1993). In particular, we are not aware of any studies of the effects of solar UVR on phytoplanktonic species of the temperate region of Patagonia in South America. This is rather surprising, considering the high productivity of the coastal areas of Patagonia (Charpy & Charpy-Roubaud 1980) and the very interesting characteristics that would warrant this type of photobiological study. The mid-latitude aquatic environments of Patagonia, especially the area including the Atlantic coast, receive high daily doses of solar radiation during the austral spring and summer due to a combination of several factors: (1) relatively high irradiances and daylight periods of up to 18 h (Orce & Helbling 1997); (2) low cloudiness (Lubin & Jensen 1995), and (3) clear skies with very low amounts of particles. In addition to these factors, the region is under the influence of periodic events of low column ozone concentrations due to its proximity to the Antarctic polar vortex and the ozone 'hole' (Orce & Helbling 1997), thus receiving sporadically higher levels of solar UV-B radiation.

This study evaluates the effects of solar UVR upon photosynthesis and DNA in phytoplanktonic species of the Argentinean Sea–Atlantic Ocean, in the area of Bahía Bustamante, Chubut (Patagonia), which is a nutrient rich-high primary productivity region, and

commercially important for the collection of the alga *Gracilaria verrucosa* (Hudson) Papenf. (Rhodophyta). To our knowledge this is the first study that describes *in situ* UVR effects on photosynthetic performance and, simultaneously, DNA damage induction in marine plankton organisms.

MATERIALS AND METHODS

This study was conducted at Bahía Bustamante, Chubut (45° S, 66.5° W), Argentina (Fig. 1), during January 1999. Surface water samples were collected early in the morning using acid-cleaned (1 N HCl) polycarbonate bottles and incubated *in situ* (0, 3, 6, and 9 m depth) for 6 h centered around local noon (i.e., from 09:00 to 15:00 h)

In order to determine the effects of solar UVR on photosynthetic rates, duplicate samples were placed in 50 ml quartz tubes and inoculated with 5 μ Ci (0.185 MBq) of labeled sodium bicarbonate (Steeman Nielsen 1952). Three different radiation treatments were implemented at each depth: (1) uncovered quartz tubes (samples receiving both photosynthetic available radiation [PAR, 400 to 700 nm] and UV), (2) tubes covered with a Mylar-D film, transmitting UV-A (315 to 400 nm) + PAR, and (3) tubes covered with Plexiglas filter UF-3, so that samples received only PAR. The tubes were then placed in aluminum anodized frames that were connected to a buoy, down to the depths mentioned above. After the incubation period, the samples were filtered onto Whatman GF/F filters (25 mm in diameter), exposed to HCl fumes overnight, dried and counted using a liquid scintillation counter (Holm-Hansen & Helbling 1995).

To evaluate the formation of CPDs (i.e., DNA damage) and the potential of repairing DNA, water sam-




Fig. 1. Map of the Chubut Province, Argentina, indicating the sampling and experimentation site at Bahía Bustamante. Inset: relative location of Chubut in South America

ples containing natural phytoplankton assemblages were placed in 10 l UV-transparent polypropylene bags hanging next to the aluminum frames. These bags have a very high transparency for all solar wavelengths (Visser et al. 1999). At the start of the experiment ($t = 0$) and after the incubation period, the samples were filtered and fractionated onto 10, 2, and 0.2 μm polycarbonate membrane filters (Poretics, 47 mm) and immediately frozen in liquid nitrogen (-180°C) until analyses, which were carried out at the University of Groningen, The Netherlands. DNA was extracted using a modified method from Doyle & Doyle (1991). Filters were incubated at 60°C for 30 min with 750 μl preheated CTAB isolation buffer (2% [w/v] CTAB Sigma, 1.4 M NaCl, 0.2% [v/v] β -mercaptoethanol, 20 mM EDTA, 100 mM Tris-HCl pH = 8.0). An aliquot of 750 μl CIA (chloroform/isoamylalcohol 24:1) was added to extract the DNA from cell debris and proteins. After centrifugation (14 000 rpm [$20\,000 \times g$], 10 min), 0.5 ml of cold isopropanol was added to the upper (water) phase to precipitate the DNA (1 h at 4°C). After centrifugation (14 000 rpm [$20\,000 \times g$], 30 min, at 4°C) the supernatant was removed and the pellet was washed with 1 ml of 80% ice-cold ethanol (15 min, at -20°C , followed by centrifugation, 30 min at 4°C). Finally the DNA pellet was dried under vacuum and resuspended in TE buffer (1 mM Tris-HCl pH = 8.0, 0.1 mM EDTA). To remove the RNA, the DNA was incubated for 1 h with 75 $\mu\text{g ml}^{-1}$ RNase (Boehringer Mannheim) at room temperature. The amount of DNA was determined fluorometrically using Picogreen dsDNA quantitation reagent (dilution 1:400, Molecular Probes) on a 1420 Victor multilabel counter (EG&G Wallac, excitation 485 nm, emission 535 nm).

The amount of CPDs was determined using the method of Boelen et al. (1999, unpubl.), employing a primary antibody directed mainly to thymine dimers. In short, 100 ng of heat-denatured DNA samples were blotted onto nitrocellulose membranes (Schleicher and Schuell, Protran 0.1 μm). The membranes were baked at 80°C to immobilize the DNA. After a 30 min blocking step, followed by 3 washing steps, the membranes were incubated with the primary antibody H3 overnight at 4°C . After repeated washing, incubation with the secondary antibody (HRP rabbit-antimouse, Dako P0260) was done for 2 h at room temperature. CPDs were detected using ECL detection reagents (RPN2106 Amersham) in combination with photosensitive films (Kodak-X-AR-5). Finally, the films were scanned and the dimers were quantified using Image Quant software (version 4.2, Molecular Dynamics). Each blot contained 2 dilution series of standard DNA with known amounts of CPDs (Boelen et al. 1999, unpubl.). In order to determine DNA effective radia-

tion and attenuation, a DNA biosimulator was used as described in Boelen et al. (1999, unpubl.). Duplicate acid-cleaned quartz tubes, containing bare DNA, were placed close to the polypropylene bags and incubated for the duration of the incubation experiment.

Samples were also taken for analysis of chlorophyll *a* (chl *a*), floristic composition and absorption characteristics of phytoplankton. The analyses of these samples were done as follows: (1) chl *a* analysis: 100 ml of sample was filtered onto a Whatman GF/F filter (25 mm in diameter) and the photosynthetic pigments extracted in absolute methanol for 1 h (Holm-Hansen & Riemann 1978). Chl *a* concentration was then calculated from the fluorescence of the extract (Holm-Hansen et al. 1965), using a Turner Designs fluorometer (model TD 700). (2) Samples for identification and enumeration of phytoplankton were placed in 125 ml brown bottles and fixed with buffered formalin (final concentration of 0.4% in the sample); after settling 25 to 50 ml of sample, they were analyzed with an inverted microscope (Leica DM IL) following the technique described in Villafañe & Reid (1995).

Incident solar radiation was measured continuously using a broad band ELDONET radiometer (Real Time Computers Inc.), which measures UV-B (280 to 315 nm), UV-A (315 to 400 nm) and PAR (400 to 700 nm) with a frequency of 1 reading min^{-1} . Optical characteristics of the water column were determined using a profiling broad band ELDONET radiometer (Real Time Computers Inc.), which measures UV-B, UV-A, PAR, water temperature and water depth; a total of 6 profiles were done with this sensor. This instrument was deployed by hand from a Zodiak at a site next to our array for *in situ* incubations

RESULTS

UV-B radiation decreased with depth in the water column, and the attenuation coefficient ($K_{\text{UV-B}}$) was 0.80 m^{-1} (Fig. 2A), with the 1% radiation level found at 5.8 m. The accumulation of CPDs in bare DNA from the biosimulator was high at 0 m: 800 CPDs per 10^6 nucleotides. However, DNA effective radiation was so rapidly attenuated that virtually no CPDs could be detected at 3 m. UV-A radiation and PAR penetrated deeper into the water column, with the 1% radiation level for UV-A at 8.7 m, while the irradiance level for PAR at the bottom (i.e., 15 m with high tide) was 10% of the surface irradiance (calculated from K_{PAR}). The attenuation coefficients for UV-A ($K_{\text{UV-A}}$) and PAR (K_{PAR}) were 0.53 and 0.16 m^{-1} , respectively (Fig. 2B,C). Phytoplankton distribution in the water column was rather homogeneous and no stratification was noticeable from the temperature profile (Fig. 3). Although




Fig. 2. Representative profiles showing the underwater radiation field next to our *in situ* experiments; radiation in $W m^{-2}$. (A) UV-B radiation; $K_{UV-B} = 0.80 m^{-1}$; (B) UV-A radiation; $K_{UV-A} = 0.53 m^{-1}$; and (C) photosynthetic active radiation (PAR); $K_{PAR} = 0.16 m^{-1}$. Profiles done on January 13, 1999 (14:00 h local time)




Fig. 3. Water temperature profile (in $^{\circ}C$) as a function of depth in waters close to our *in situ* experiments. Data obtained with the temperature sensor on the ELDONET radiometer

we did not measure salinity, we estimated that there were no significant changes in the density of the upper part of the water column because there was no input of freshwater into this system. Chl *a* concentration in the surface sample was $3.1 \mu g l^{-1}$. Mean surface incident irradiance values during the time of our experimentation were 2.2, 60 and $450 W m^{-2}$ for UV-B, UV-A and PAR, respectively; the doses during the 6 h incubation period were 47, 1291 and $9661 KJ m^{-2}$ for UV-B, UV-A and PAR, respectively.

Fig. 4 represents the photosynthetic characteristics of natural phytoplankton assemblages when exposed *in situ* to the 3 radiation treatments mentioned above. The amount of carbon fixed in surface waters during the 6 h incubation period (Fig. 4A) was about $17.5 \mu g C l^{-1}$ in the treatment that received only PAR. The carbon fixation for PAR was rather constant with depth, hav-

ing a very slight increase at 3 to 6 m, suggesting little photoinhibition at the surface due to high PAR levels. The amount of carbon fixed by phytoplankton receiving UV-A + PAR and UV + PAR was 12.6 and $11 \mu g C l^{-1}$, respectively, and no significant differences ($p < 0.05$) were observed among treatments below 6 m depth (Fig. 4A). Assimilation numbers (Fig. 4B) were rather constant with depth (about $0.9 mg C [mg chl a]^{-1} h^{-1}$) for the treatment that received only PAR, with maximum numbers found at 3 m. Photosynthetic inhibition (Fig. 4C) at the surface was 29 and 7% for UV-A and UV-B, respectively, as compared with the treatment that received only PAR. This inhibition decreased with depth so that no significant differences ($p < 0.05$) among treatments were found at 6 m. UV-A was responsible for most of the observed inhibition in the upper water column (i.e., upper 3 m).

No DNA could be extracted from the 2 and 10 μm fractions, indicating that cells bigger than 2 μm represented a negligible proportion of the plankton biomass in terms of DNA. Microscopic observation of phytoplankton samples also revealed that the phytoplankton crop was mainly composed of picoplanktonic cells (less than 2 μm in diameter), with very few representatives of larger cells such as diatoms (e.g., *Pseudonitzschia* spp., *Skeletonema costatum*, *Licmophora* sp.). The natural phytoplankton assemblage already had a high level of DNA damage at the time the incubation started, with a mean of 375 T<>T per 10^6 nucleotides ($t = 0$, Fig. 5B). The high levels of CPDs in the phytoplankton were consistently observed in all morning samples collected for other experiments (data not presented). At the end of the incubation period, the formation of CPDs increased significantly in surface waters (mean value of 650 T<>T per 10^6 nucleotides). In contrast, CPD levels at 3 and 6 m depth diminished to about 250 CPDs per 10^6 nucleotides (Fig. 5B).

DISCUSSION

Two of the most important physiological forms of natural UVR stress in phytoplanktonic organisms are photosynthetic inhibition and DNA damage. Many studies have evaluated the role of UVR in inhibiting photosynthesis (Helbling et al. 1992, Smith et al. 1992, Neale et al. 1998) and damaging DNA (Karentz et al. 1991, Buma et al. 1997, Boelen et al. unpubl.) in various regions. In this study we considered both targets (i.e., photosynthesis and DNA) at the same time, thus pro-

viding a powerful tool to evaluate the overall impact of solar radiation in the very productive waters of the Patagonian coast. Our results show that UVR can simultaneously affect photosynthesis and DNA as observed in surface waters (Figs. 4 & 5).

Phytoplankton photosynthesis in our study was more inhibited by UV-A than by UV-B (Fig. 4), with PAR causing relatively minor photoinhibition. The greater inhibition by UV-A, as compared to UV-B, has also been observed in other freshwater and marine environments (Bühlmann et al. 1987, Kim & Watanabe 1993, Villafaña et al. 1999). The inhibition of phytoplankton photosynthesis due to solar UVR decreased with depth but was mostly limited to the upper 3 m of the water column (Fig. 4); below this depth, solar UVR decreased just enough (Fig. 2) to cause a mild effect. In particular, UV-B radiation in our study area (Fig. 2A) had a relatively high attenuation coefficient (i.e., 0.80 m^{-1}), as compared to Antarctic areas with similar particle concentration (Helbling et al. 1994). The high $K_{\text{UV-B}}$ in our study most probably reflects the amount of terrigenous material at the coastal station where we conducted the experimentation. However, the attenuation coefficients at Bahía Bustamante were lower than in other coastal areas (Booth & Morrow 1997). When compared with other environments of Patagonia, such as the sub-Antarctic waters of the Beagle Channel, Tierra del Fuego (Helbling et al. 1996), the total amount of photosynthetic inhibition at the surface was similar at both sites (about 40%); however, its depth distribution was different (Villafaña et al. 2001). When comparing the overall impact of UVR at these sites, we considered the photosynthetic inhibition as a function of the optical depth (the optical depth of 4.6 is equal to $K_{\text{PAR}} \cdot Z_{\text{eu}}$). In the Beagle Channel, photosynthetic inhibition in the water column reached down to 3 optical depths (Helbling et al. 1996, Villafaña et al. 2001), while at Bahía Bustamante it reached down to 1.5 optical depths (Fig. 4C), decreasing almost exponentially. Thus, the depth integrated loss of carbon fixation (upper 10 m of the water column) due to UV-B (determined when compared to the PAR treatment that was taken as 'no inhibition') would be higher for the Beagle Channel assemblages (calculated value of 24%) than for Bahía Bustamante phytoplankton (calculated value of 3%). The integrated inhibition at Bahía Bustamante is closer to the 4.9% value reported for UV-B inhibition in the Antarctic (Holm-Hansen et al. 1993b).




Fig. 4. Depth distribution of (A) carbon fixation, (B) assimilation numbers and (C) percentage of photosynthetic inhibition for natural phytoplankton populations from Bahía Bustamante exposed to 3 radiation treatments; photosynthetic inhibition in the PAR treatment has been set to zero. +: 1 standard deviation




Fig. 5. Depth profiles of CPD formation ($\text{T}<>\text{T}$ per 10^6 nucleotides). (A) Bare DNA samples, and (B) natural phytoplankton populations. The symbol at $t = 0$ indicates the amount of CPDs at the start of the experiment. +: 1 standard deviation

Initial CPD levels were very high in this study (Fig. 5B) and higher than those found in other areas, for instance in marine tropical picoplankton (Boelen et al. unpubl.), in the Antarctic (Buma et al. 2001) or in the plankton from Lake Titicaca, Bolivia (Helbling et al. 2001). This would seem to indicate a prolonged history of previous UV-B exposures in the water column, combined with a low repair capacity of cells. On the other hand, CPD levels decreased at 3 and 6 m depth, indicating that either repair was taking place, or damage was diluted by *de novo* DNA synthesis in viable cells. Also, CPD formation may result in an increasing proportion of non-viable cells in the plankton assemblage, eventually leading to a loss of cells from the

water column through lysis (Boelen et al. unpubl.). It has to be stressed that our CPD results reflect not only picophytoplankton cells but also heterotrophic bacteria, all retained on the 0.2 μm filter. However, as has been demonstrated recently by Boelen et al. (1999b), no significant difference in CPD accumulation could be observed in these 2 functional picoplankton groups in tropical marine assemblages. Some studies had determined that small cells (i.e., nanoplanktonic <20 μm in diameter) were generally more resistant than large cells when looking at photosynthetic inhibition (Helbling et al. 1992, Laurion & Vincent 1998). Simulated *in situ* experimentation data (not shown), conducted parallel to the *in situ* studies presented in this paper, were used to calculate the biological weighting function (BWF) for inhibition of phytoplankton photosynthesis. Eight independent experiments were conducted using sharp-cut of filter (Schott); a mean BWF was calculated using a BWF-PI model (Cullen et al. 1992, Neale & Kieber 2000), and the spectral dependence of the BWF in the broadband intervals was extracted using the method of Rundel (1983). A comparison of Bahía Bustamante BWF with the one for *Phaeodactylum* sp. (Cullen et al. 1992) showed that phytoplankton from Bahía Bustamante were significantly less sensitive to UVR for wavelengths higher than 300 nm. However, in contrast with this, small cells (less than 2 μm) seem to be more sensitive to UVR when looking at CPD formation (Karentz et al. 1991, Boelen et al. unpubl.). Also, natural Antarctic picoplankton assemblages displayed significantly higher CPD levels compared with larger, diatom-containing size fractions, as a result of exposure to solar UV-B radiation (Buma et al. 2001).

Clearly, CPDs were accumulated in the surface, but no accumulation was observed at 3 and 6 m depth, in accordance with the strong attenuation of DNA effective UVR (Fig. 5A). It should be considered that cells move in the upper mixed layer (see Fig. 3 for temperature profile) but in an *in situ* experimentation they are kept at a fixed depth. This means that the surface samples would receive higher irradiance than they would in the upper mixed layer, so an increase in CPDs would be observed. However, the cells incubated at 3 and 6 m receive less irradiance, thus a decrease in CPDs was noticed (Fig. 5). We have recently shown for plankton organisms from Lake Titicaca (Helbling et al. 2001) that both UV-A and UV-B inhibit photosynthesis, with UV-A having a stronger effect than UV-B. At the same time CPD accumulation was only related with UV-B exposure, as found in this study in Patagonia. Moreover, daily patterns of both photosynthetic inhibition and DNA damage accumulation were far from similar. Evidently, DNA damage accumulation and photosynthetic inhibition patterns reveal effects on various, rel-

atively independent cell targets, i.e., nuclear DNA and the photosynthetic apparatus, although interactions between the 2 processes are imaginable (Helbling et al. 2001). Assimilation numbers in our coast reach 5 mg C [mg chl a]⁻¹ h⁻¹ during the spring bloom, but were lower in the mid-summer post-bloom assemblages (Villafañe, unpub. data). The relatively low assimilation values (about 1 mg C [mg chl a]⁻¹ h⁻¹), compared to the spring bloom, found in this study (Fig. 4), together with the high CPDs values found in the water column, might in fact reveal an interactive process between DNA damage and photosynthetic apparatus.

In conclusion, this study has demonstrated that UV stress may be brought about by multiple target effects, at least at the surface. The extent to which these targets are affected will be determined by irradiance conditions or species-specific differences in vulnerability, for instance related to cell size.

Acknowledgements. We thank M. José, C. Soriano, R. Cariaga and M. Hernández for their help during field work; L. Sala carried out ¹⁴C analysis and L. Taibo helped with computer drawings. We also thank the Soriano SA and Fundación Playa Unión for logistic support. We thank the comments of 4 anonymous reviewers. This work was supported by Agencia Nacional de Promoción Científica y Tecnológica, ANPCyT/BID (Project PICT97 07-00000-02206); The Third World Academy of Sciences, TWAS (Project 98-036 RG/BIO/LA) and Fundación Antorchas (Projects A-13622/1-100 and A13669/1-3). This is Contribution No. 13 of Estación de Fotobiología Playa Unión.

LITERATURE CITED

- Arrigo KR (1994) Impact of ozone depletion on phytoplankton growth in the Southern Ocean: large-scale spatial and temporal variability. *Mar Ecol Prog Ser* 114:1–12
- Behrenfeld M, Hardy J, Gucinski H, Hanneman A, Lee H II, Wones A (1993) Effects of ultraviolet-B radiation on primary production along latitudinal transects in the South Pacific Ocean. *Mar Environ Res* 35:349–363
- Boelen P, Obernosterer I, Vink AA, Buma AGJ (1999) Attenuation of biologically effective UV radiation in tropical Atlantic waters measured with a biochemical DNA dosimeter. *Photochem Photobiol* 69:34–40
- Booth CR, Morrow JH (1997) The penetration of UV into natural water. *Photochem Photobiol* 65:254–257
- Bühlmann B, Bossard P, Uehlinger U (1987) The influence of longwave ultraviolet radiation (u.v.-A) on the photosynthetic activity (¹⁴C-assimilation) of phytoplankton. *J Plankton Res* 9:935–943
- Buma AGJ, Engelen AH, Gieskes WWC (1997) Wavelength-dependent induction of thymine dimers and growth rate reduction in the marine diatom *Cyclotella* sp. exposed to ultraviolet radiation. *Mar Ecol Prog Ser* 153:91–97
- Buma AGJ, de Boer MK, Boelen P (2001) Depth distributions of DNA damage in Antarctic marine phyto- and bacterioplankton exposed to summertime ultraviolet radiation. *J Phycol* (in press)
- Charpy LJ, Charpy-Roubaud CJ (1980) La production primaire des eaux du golfe San Jose (Peninsule Valdes,

- Argentina). III. Estimation de la production phytoplantonique annuelle. *Hydrobiologia* 75:225–229
- Cullen JJ, Neale PJ, Lesser MP (1992) Biological weighting function for the inhibition of phytoplankton photosynthesis by ultraviolet radiation. *Science* 258:645–650
- Doyle JJ, Doyle JL (1991) Isolation of plant DNA from fresh tissue. *Focus* 12:13–15
- Häder DP (1997) Effects of UV radiation on phytoplankton. In Jones JG (ed) *Advances in microbial ecology*. Plenum Press, New York, p 1–26
- Helbling EW, Villafañe VE, Ferrario M, Holm-Hansen O (1992) Impact of natural ultraviolet radiation on rates of photosynthesis and on specific marine phytoplankton species. *Mar Ecol Prog Ser* 80:89–100
- Helbling EW, Avaria S, Letelier J, Montecino V, Ramírez B, Ramos M, Rojas W, Villafañe VE (1993) Respuesta del fitoplancton marino a la radiación ultravioleta en latitudes medias (33° S). *Rev Biol Mar* 28:219–237
- Helbling EW, Villafañe VE, Holm-Hansen O (1994) Effects of ultraviolet radiation on antarctic marine phytoplankton photosynthesis with particular attention to the influence of mixing. In: Weiler S, Penhale P (eds) *Ultraviolet radiation in Antarctica: measurements and biological effects*. Am Geophys Union Antarct Res Ser 62:207–227
- Helbling EW, Hernando MP, Holm-Hansen O, Villafañe VE (1996) Efectos de la radiación ultravioleta en el fitoplancton del Canal Beagle, Tierra del Fuego. XVI Jornadas de Ciencias del Mar, Concepción
- Helbling EW, Villafañe VE, Buma AGJ, Andrade M, Zaratti F (2001) DNA damage and photosynthetic inhibition induced by solar UV in tropical phytoplankton (Lake Titicaca, Bolivia). *Eur J Phycol* (in press)
- Holm-Hansen O, Helbling EW (1995) Técnicas para la medición de la productividad primaria en el fitoplancton. In: Alveal K, Ferrario ME, Oliveira EC, Sar E (eds) *Manual de métodos ficológicos*. Universidad de Concepción, Concepción, p 329–350
- Holm-Hansen O, Riemann B (1978) Chlorophyll a determination: improvements in methodology. *Oikos* 30:438–447
- Holm-Hansen O, Lorenzen CJ, Holmes RW, Strickland JDH (1965) Fluorometric determination of chlorophyll. *J Cons Int Explor Mer* 30:3–15
- Holm-Hansen O, Lubin D, Helbling EW (1993a) Ultraviolet radiation and its effects on organisms in aquatic environments. In: Young AR, Björn LO, Moan J, Nultsch W (eds) *Environmental UV photobiology*. Plenum Press, New York, p 379–425
- Holm-Hansen O, Helbling EW, Lubin D (1993b) Ultraviolet radiation in Antarctica: inhibition of primary production. *Photochem Photobiol* 58:567–570
- Karentz D, Cleaver JE, Mitchell DL (1991) Cell survival characteristics and molecular responses of Antarctic phytoplankton to ultraviolet-B radiation. *J Phycol* 27:326–341
- Kim DS, Watanabe Y (1993) The effect of long-wave ultraviolet radiation (UV-A) on the photosynthetic activity of natural population of freshwater phytoplankton. *Ecol Res* 8: 225–234
- Laurion I, Vincent WF (1998) Cell size versus taxonomic composition as determinants of UV-sensitivity in natural phytoplankton communities. *Limnol Oceanogr* 43: 1774–1779
- Lubin D, Jensen EH (1995) Effects of clouds and stratospheric ozone depletion on ultraviolet radiation trends. *Nature* 377:710–713
- Madronich S (1993) The atmosphere and UV-B radiation at ground level. In: Young AR, Björn LO, Moan J, Nultsch W (eds) *Environmental UV photobiology*. Plenum Press, New York, p 1–39
- Neale PJ, Kieber DJ (2000) Assessing biological and chemical effects of UV in the marine environment: spectral weighting functions. In: Hester RE, Harrison RM (eds) *Causes and environmental implications of increased U.V.-B. radiation*. Royal Society of Chemistry, Cambridge
- Neale PJ, Davis RF, Cullen JJ (1998) Interactive effects of ozone depletion and vertical mixing on photosynthesis of Antarctic phytoplankton. *Nature* 392:585–589
- Orce VL, Helbling EW (1997) Latitudinal UVR-PAR measurements in Argentina: extent of the 'Ozone hole'. *Global Planet Change* 15:113–121
- Rundel RD (1983) Action spectra and estimation of biologically effective UV radiation. *Physiol Plant* 58:360–366
- Smith RC, Prézelin BB, Baker KS, Bidigare RR, Boucher NP, Coley TL, Karentz D, MacIntyre S, Matlick HA, Menzies D, Ondrusek M, Wan Z, Waters KJ (1992) Ozone depletion: ultraviolet radiation and phytoplankton biology in Antarctic waters. *Science* 255:952–959
- Steeman Nielsen E (1952) The use of radiocarbon (^{14}C) for measuring organic production in the sea. *J Cons Int Explor Mer* 18:117–140
- Villafañe VE, Reid FMH (1995) Métodos de microscopía para la cuantificación del fitoplancton. In: Alveal K, Ferrario ME, Oliveira EC, Sar E (eds) *Manual de métodos ficológicos*. Universidad de Concepción, Concepción, p 169–185
- Villafañe VE, Andrade M, Lairana V, Zaratti F, Helbling EW (1999) Inhibition of phytoplankton photosynthesis by solar ultraviolet radiation: studies in Lake Titicaca, Bolivia. *Freshw Biol* 42:215–224
- Villafañe VE, Helbling EW, Zagarese HE (2001) Solar ultraviolet radiation and its impact on aquatic systems of Patagonia, South America. *Ambio* (in press)
- Visser PM, Snelder E, Kop AJ, Boelen P, Buma AGJ, van Duyl FC (1999) Effects of UV radiation on DNA photodamage and production in bacterioplankton in the coastal Caribbean Sea. *Aquat Microb Ecol* 20:49–58

Editorial responsibility: Otto Kinne (Editor), Oldendorf/Luhe, Germany

*Submitted: March 22, 2000; Accepted: August 8, 2000
Proofs received from author(s): January 18, 2001*