

University of Groningen

## Functioning of *Plantago major* and *Urtica dioica* exposed to elevated CO<sub>2</sub>

den Hertog, Jeroen

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1997

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

den Hertog, J. (1997). Functioning of *Plantago major* and *Urtica dioica* exposed to elevated CO<sub>2</sub>: Analysis of growth pattern in relation to C and N allocation s.n.

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Summary

The atmospheric CO<sub>2</sub> concentration has risen from 280  $\mu\text{l l}^{-1}$  at the start of the industrial revolution, around 1880, to over 350  $\mu\text{l l}^{-1}$  at present. This rise is expected to increase far into the next century, even if the release of CO<sub>2</sub> to the atmosphere would stabilise at the present level. The often predicted increase in the mean global temperature, the so-called 'greenhouse effect', is the most conspicuous consequence of this rise in CO<sub>2</sub> concentration. Although the eight o'clock news sometimes refers to CO<sub>2</sub> as an air-polluting gas, its presence is a prerequisite for plant growth and agricultural production. CO<sub>2</sub> is a substrate for photosynthesis, and is often in short supply and thereby it limits plant growth. An increase in the concentration of CO<sub>2</sub> may cause a rise in the rate of photosynthesis and an increase in the production of individual plants, crops and natural vegetation. Indeed, since the early sixties, in the dutch greenhouse vegetable production 'CO<sub>2</sub>-fertilisation' is a common practice to increase yield.

In most plants from temperate climate zones grown under favourable conditions, in the short term, a rise in CO<sub>2</sub> concentration causes an increase in the rate of photosynthesis through an increased substrate supply as well as a suppression of photorespiration. The rate of growth closely follows this short-term effect on photosynthesis. Within a day the relative growth rate (RGR) is increased in plants grown at elevated CO<sub>2</sub> as compared to plants grown at ambient CO<sub>2</sub> levels. A plant's long term response to elevated CO<sub>2</sub> is more uncertain. In various plant species a continued stimulation of the rate of photosynthesis as well as a return of this rate to the level observed in control plants has been found.

For dutch agriculture an elevated CO<sub>2</sub> concentration may contribute to a rise in yield, since growth conditions approach optimal water and nutrient supply in many years. However, in regions with more extreme climates or poor soils, conditions of water and nutrient supply are often less favourable for plant growth and a rising atmospheric CO<sub>2</sub> concentration may not lead to the stimulation of agricultural production. The same applies to natural vegetation, which in many cases grows in marginal areas with respect to climatic and soil conditions. Under these circumstances, an elevation of the CO<sub>2</sub> concentration will interact with e.g. plant nutrition and water supply in determining the rate of photosynthesis and the growth pattern of the plant.

Over the last decades many reports have dealt with effects of elevated CO<sub>2</sub> on agricultural production. Since the early eighties wild plant species, natural vegetations and ecosystems begun to attract scientific attention. If plant species differ in their response to variations in the atmospheric CO<sub>2</sub> level, this would imply possible shifts in the interaction between species in their natural environment. In

addition, other growth conditions interfere with the plant's reaction to elevated CO<sub>2</sub> and even more changes in natural ecosystems could occur as a result of an increase in atmospheric CO<sub>2</sub> concentration.

This study focusses on the functioning of whole plants under conditions of elevated (700  $\mu\text{l l}^{-1}$ ) as compared to the ambient level of CO<sub>2</sub> (350  $\mu\text{l l}^{-1}$ ). The species used are *Plantago major* ssp. *pleiosperma* (great plantain) and *Urtica dioica* (stinging nettle), two species that grow in many (semi-)natural vegetations in Europe. All experiments were carried out in growth rooms, in nutrient solution, in order to have optimal control over environmental conditions.

The aim of the study was (i) to describe the growth response of plants, in the vegetative growth period, to this doubling of the atmospheric CO<sub>2</sub> concentration and (ii) to identify physiological and morphological traits that are sensitive to changes in the CO<sub>2</sub> level. The second aim was to relate the observed growth response to the nitrogen (N) metabolism of the plant, since nitrogen is quantitatively an important nutrient, that often limits plant growth under natural conditions. These two aims were a prerequisite for the following step, which was to analyse the response of *P. major* to elevated CO<sub>2</sub> under conditions of limiting N supply. Finally, the results of this study were used to evaluate if generalisations could be made of functional types of plant species, that respond either significantly or marginally to an increased atmospheric CO<sub>2</sub> concentration.

Both in *P. major* and in *U. dioica* the RGR was transiently stimulated by an elevation of the atmospheric CO<sub>2</sub> concentration (chapters 2 and 3). The higher RGR was accounted for by an increase of the rate of photosynthesis at elevated CO<sub>2</sub>, but the stimulating effect on photosynthesis was counteracted by decreases in the Specific Leaf Area (SLA, chapters 2 and 3). Accumulation of nonstructural carbohydrates could partly explain this effect on the SLA. In addition, differences in plant size, due to the CO<sub>2</sub> treatment and a coupling between leaf morphology and plant size, accounted for the observed decrease of the SLA at elevated CO<sub>2</sub> (chapter 3). In some experiments the increase in the rate of photosynthesis was accompanied by increases in the rate of shoot and root respiration as well (chapters 2 and 6), but the latter responses were rather variable and the stimulation of the rate of respiration seemed to be restricted to the phase of growth stimulation by elevated CO<sub>2</sub> (chapter 6).

*U. dioica* showed a longer-lasting stimulation of the RGR at elevated CO<sub>2</sub> than *P. major*. After a doubling of the CO<sub>2</sub> concentration, the former species, accumulated nonstructural carbohydrates, starch and soluble sugars in the leaves to a lesser extent than the latter (chapter 3), which led to relatively smaller effects of elevated CO<sub>2</sub> on SLA and the % dry matter in the leaves of *U. dioica* as well. The observed decreases in the concentration of total nitrogen and those of different nitrogen pools in plants grown in elevated CO<sub>2</sub> were largely due to this accumulation of

carbohydrates. The  
as a consequence of  
and the growth form  
of small secondary  
so-called more inde  
*U. dioica* to allocat  
branches, in compa

The interaction of a  
was studied in *P. m*  
various exponential  
exponential nutri  
experiment, wherea  
initial phase of non  
exhaustion of the n  
regime, and, consec  
supply. Furthermore  
nutrient solution is  
severe nitrogen lim  
supply the level of f  
for both CO<sub>2</sub> treatm

Growth of *P. major*  
various rates of nitro  
RGR upon elevation  
differences in respon  
treatments was restr  
the leaves at elevate  
doubling of the CO<sub>2</sub>  
extra carbon gained  
contribute to the syn  
and was fixed at a h  
neither lead to grow  
roots or storage orga  
suggested as a way  
nutrient-limited con  
when grown at a co

The data from chapt  
nutrient-limited gro  
relatively larger roo  
those experiments. I  
growth of *P. major* a  
regimes were model

carbohydrates. The difference in response between the two species was explained as a consequence of the species-dependent RGR, which was higher in *U. dioica*, and the growth form. *P. major* forms rosettes of leaves and has a limited number of small secondary shoots, whereas *U. dioica* forms a widely branched shoot, in a so-called more indeterminate growth pattern. Those characteristics seem to enable *U. dioica* to allocate more of the extra carbon gained at elevated  $\text{CO}_2$  to new branches, in comparison with that in *P. major*.

The interaction of an elevated  $\text{CO}_2$  concentration and a limited supply of nitrogen was studied in *P. major* (chapter 4). Nitrate in the nutrient solution was supplied at various exponential rates, instead of different concentrations. An advantage of the exponential nutrient supply is, that growth is limited right from the start of the experiment, whereas in an experiment at different nutrient concentrations, an initial phase of nonlimiting nitrate supply is immediately followed by a complete exhaustion of the nutrient solution. The latter procedure creates a variable nutrient regime, and, consequently the plant is exposed to a continuously changing nitrate supply. Furthermore, if the RGR is stimulated, as is the case at elevated  $\text{CO}_2$ , the nutrient solution is exhausted more rapidly and the plant thus experiences a more severe nitrogen limitation than under control conditions. With exponential nitrogen supply the level of nutrient stress can be maintained at a constant and equal level for both  $\text{CO}_2$  treatments.

Growth of *P. major* plants was compared between the two  $\text{CO}_2$  treatments and at various rates of nitrogen supply, in the period after the transient stimulation of the RGR upon elevation of the  $\text{CO}_2$  concentration. At this relatively long term, the differences in response to a growth-limiting N supply between the two  $\text{CO}_2$  treatments was restricted to a large accumulation of nonstructural carbohydrates in the leaves at elevated  $\text{CO}_2$  at all levels of nitrogen supply. No other shifts upon a doubling of the  $\text{CO}_2$  concentration in the allocation of carbon were observed. The extra carbon gained per leaf area was invested in carbohydrates, which did not contribute to the synthesis of new leaf area. Although  $\text{CO}_2$  kept entering the leaves and was fixed at a higher rate in plants grown at elevated  $\text{CO}_2$ , the extra  $\text{CO}_2$  did neither lead to growth of new photosynthetic tissue nor to the formation of extra roots or storage organs. An investment of the extra carbon in roots has often been suggested as a way in which plants could benefit from elevated  $\text{CO}_2$  even under nutrient-limited conditions. Apparently, however, this did not happen in *P. major* when grown at a constant exponential nitrate supply.

The data from chapter 4 were used to simulate, if the outcome of experiments with nutrient-limited growth at a constant nutrient concentration, which often includes a relatively larger root weight at elevated  $\text{CO}_2$ , could be explained by the scenario of those experiments. In chapter 5 a model was constructed, which mimicked the growth of *P. major* as described in chapter 4. Subsequently different nutrient regimes were modelled, which represented growth of plants at different concen-

trations of nitrate in a nutrient solution, with regular replenishment of the solution, such as occurs in the classical nutrient-limitation studies. The simulation indicated, that, indeed, the ratio of the root to total plant weight (root weight ratio, RWR) was increased in the period between free access to nitrate and severe nitrate limitation, when the RGR drops to almost zero. This study did not answer the question, which scenario for the availability of nitrate under natural field conditions is most likely. However, it did indicate that an increased supply of carbon in plants grown at elevated  $\text{CO}_2$  did not lead to a relative increase in root size at a given level of nutrient stress.

In future studies concerned with elevated  $\text{CO}_2$ , which are designed to gain insight in the functioning of natural ecosystems, what responses can be expected in various plant species and what characteristics determine the responsiveness of a species to changes in the  $\text{CO}_2$  concentration? The observed responses of *P. major* and *U. dioica* indicate, that differences in growth form and specific RGR are traits that may be indicative for the magnitude and duration of the growth response to be expected. Additionally, species with large non-photosynthetic organs, that only consume carbon, seem to possess a flexible so-called 'sink' that can use the extra carbon provided by growth at elevated  $\text{CO}_2$ . The degree to which a plant can respond to elevated  $\text{CO}_2$  with an increase in RGR seems to be indicated by the balance between the production of structural and nonstructural carbohydrates; if the extra carbon can be invested in new structural dry matter, the accumulation of nonstructural dry matter is limited, and, alternatively, when the plant is not able to invest in more structural growth, the extra carbon ends up as nonstructural carbohydrates, mainly starch, in the leaves.

In ecological studies species are divided into functional groups, that are described in terms of a coherent set of traits, a life-history strategy. Plants with a high potential RGR often occur in highly productive, ruderal and disturbed habitats. Growth of those species is likely to be responsive to elevated  $\text{CO}_2$ . If species within such a community have a large sink, i.e. a tuber or stolon, or when they have an indeterminate growth pattern, they are expected to respond to an elevation of the atmospheric  $\text{CO}_2$  concentration with a significant and prolonged increase in growth rate.

## Samenvatting

Sinds het begin van de industriële revolutie is de  $\text{CO}_2$  concentratie gestegen. Naar verwachting zal deze toename in de komende eeuwen erin zouden slaan. Het meest opvallende effect is de stijging van de globale temperatuur. Vanwege de klimaatverandering is het soms een belangrijke maatregel om de noodzaak van CO<sub>2</sub> te reduceren en dus ook voor de landbouw. In veel gevallen is de thesesnelheid, en de CO<sub>2</sub> concentratie in de atmosfeer, waardoor vervolgens de natuurlijke vegetatie verandert.

Wanneer planten worden opgekweekt in termen van CO<sub>2</sub>, leiden tot een groter substratum. groeisnelheid volgt. Binnen een dag naar een verhoogde CO<sub>2</sub> bij normale CO<sub>2</sub> n. verhoging van de voortdurende verhoging naar het

In de Nederlandse vochtcondities gekweekte CO<sub>2</sub> concentratie waar de klimaatsonderzoek voedingstoffen is, gunstig effect hebben op de vegetaties, die vaak een aanbod van voedingstoffen. interacties optreden enerzijds en de fotosynthese en de