

University of Groningen

## The role of Ynt1 in nitrate and nitrite transport in the yeast *Hansenula polymorpha*

Machin, F.; Medina, B.; Navarro, F.J.; Perez, M.D.; Veenhuis, M; Tejera, P.; Lorenzo, H.; Lancha, A.; Siverio, J.M.

*Published in:*  
Yeast

*DOI:*  
[10.1002/yea.1075](https://doi.org/10.1002/yea.1075)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2004

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Machin, F., Medina, B., Navarro, F. J., Perez, M. D., Veenhuis, M., Tejera, P., ... Siverio, J. M. (2004). The role of Ynt1 in nitrate and nitrite transport in the yeast *Hansenula polymorpha*. *Yeast*, 21(3), 265 - 276. DOI: 10.1002/yea.1075

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Yeast

Yeast 2004; 21: 265–276.

Published online in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/yea.1075

## Research Article

# The role of Ynt1 in nitrate and nitrite transport in the yeast *Hansenula polymorpha*

Felix Machín<sup>#</sup>, Braulio Medina<sup>†</sup>, Francisco J. Navarro<sup>†</sup>, M. Dolores Pérez<sup>##</sup>, Marten Veenhuis<sup>2</sup>, Paula Tejera<sup>###</sup>, Helena Lorenzo<sup>####</sup>, Ana Lancha<sup>3</sup> and José M. Siverio<sup>1\*</sup>

<sup>1</sup>Departamento de Bioquímica y Biología Molecular, Grupo del Metabolismo del Nitrógeno, Universidad de La Laguna, Tenerife, Canarias, Spain

<sup>2</sup>Eukaryotic Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute (GBB), University of Groningen, Kercklaan 30, 9751 NN Haren, The Netherlands

<sup>3</sup>Departamento de Microbiología y Biología Celular, Grupo del Metabolismo del Nitrógeno, Universidad de La Laguna, Tenerife, Canarias, Spain

\*Correspondence to:

José M. Siverio, Departamento de Bioquímica y Biología Molecular, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Canarias, Spain.  
E-mail: jsiverio@ull.es

<sup>†</sup>These two authors contributed equally to this work.

<sup>#</sup>Present address: Cell Cycle Group, MRC Clinical Science Center, Imperial College London, Hammersmith Hospital Campus, Du Cane Road, London W12 0 NH, UK.

<sup>##</sup>Present address: Instituto de Enseñanza Secundaria, Lanzarote, Canarias, Spain.

<sup>###</sup>Present address: Unidad de Investigación, Hospital Universitario Nuestra Señora de Candelaria, Tenerife, Canarias, Spain.

<sup>####</sup>Present address: Instituto de Enseñanza Secundaria, Gran Canaria, Canarias, Spain.

## Abstract

**Ynt1 is the only high-affinity nitrate uptake system in *Hansenula polymorpha*. Nitrate uptake was directly correlated with the Ynt1 levels and shown to be independent of nitrate reductase (NR) activity levels. Ynt1 failed to transport chlorate and, as a result, strains lacking *YNT1* were sensitive to chlorate, as is the wild-type. Nitrite uptake in a wild-type strain was partially inhibited by nitrate to levels shown by a *YNT1*-disrupted strain in which, in turn, nitrite transport was not inhibited by nitrate. It is concluded that nitrite uptake takes place by two different transport systems: Ynt1 and a nitrite-specific transporter(s). The nitrite-specific transport system was induced by nitrate; consistently, no induction was observed in strains lacking the transcription factor *YNA1*, which is involved in nitrate and nitrite induction of the nitrate assimilatory structural genes. Ynt1 presents its optimal rate for nitrite uptake at pH 6, while pH 4 was optimal for the specific nitrite uptake system(s). At pH 5.5, the contribution of Ynt1 to high-affinity nitrate and nitrite uptake was around 95% and 60%, respectively. The apparent  $K_m$  of Ynt1 for nitrate and nitrite is in the  $\mu\text{M}$  range, as is the specific nitrite uptake system for nitrite. The analysis of the effect of the reduced nitrogen sources on nitrate assimilation revealed that glutamine inactivates nitrate and nitrite transport, dependent on Ynt1, but not the nitrite-specific system. Copyright © 2004 John Wiley & Sons, Ltd.**

**Keywords:** nitrate; nitrite; transport; yeast; *Hansenula polymorpha*

Received: 11 December 2002  
Accepted: 9 November 2003

## Introduction

In the yeast *Hansenula polymorpha*, genes encoding nitrate transporter (*YNT1*), nitrate reductase

(*YNR1*) and nitrite reductase (*YNI1*) are clustered. The same cluster also contains two  $\text{Zn(II)}_2\text{Cys}_6$  transcriptional factors encoded by *YNA1* and *YNA2*, both essential for nitrate induction of *YNT1*,

*YNR1* and *YNI1*. *YNA1* or *YNA2* disruptants are unable to grow in nitrate or nitrite (Avila *et al.*, 1995, 1998, 2002; Brito *et al.*, 1996; Pérez *et al.*, 1997).

The nitrate transporter Ynt1 has 12 putative transmembrane domains (Pérez *et al.*, 1997). It has been included in the NNP (nitrate–nitrite porter) family, involved in high-affinity nitrate and/or nitrite transport (Forde, 2000), a member of the major facilitator superfamily (Pao *et al.*, 1998). Ynt1 shows similarity to NrtA (formerly designated CRNA) and NrtB from *Aspergillus nidulans* (Unkles *et al.*, 1991, 2001) and high-affinity nitrate transporters from plants and algae (Daniel-Vedele *et al.*, 1998).

Multiplicity of nitrate transporters is a common feature found in yeast, filamentous fungi, algae and plants. In *H. polymorpha*, low-affinity nitrate transport, unrelated to Ynt1, has been observed in mutants lacking *YNT1*. However, the gene(s) involved in this transport have not yet been cloned (Machín *et al.*, 2000). Two high-affinity nitrate transporters have been found in *A. nidulans* (Unkles *et al.*, 2001). In the alga *Chlamydomonas reinhardtii*, four systems for nitrate and nitrite transport have been characterized thus far. These systems present differential specificity for nitrate and nitrite affinity and transcriptional regulation (Galván *et al.*, 1996; Navarro *et al.*, 2000; Quesada *et al.*, 1994; Rexach *et al.*, 1999). Genes encoding high- and low-affinity nitrate transporters have been isolated in higher plants (Crawford and Glass, 1998; Daniel-Vedele *et al.*, 1998; Forde, 2000; Galván and Fernández, 2001).

The capacity of high-affinity nitrate transporters to transport nitrite and chlorite in fungi has been subject to controversy. Using *Xenopus* oocytes as a heterologous expression system, it was shown by electrophysiological methods that *A. nidulans* NrtA was able to transport nitrate, nitrite and chlorite (Zhou *et al.*, 2000). However, a mutant lacking both NrtA and NrtB nitrate transporters does not show any decrease in growth on nitrite, highlighting the limited participation, if any, of these transporters in nitrite transport (Unkles *et al.*, 2001). Transport of chlorate was not detected using the *Xenopus* oocytes heterologous expression system (Zhou *et al.*, 2000), whereas an *A. nidulans* mutant strain lacking NrtA showed resistance to chlorate (Unkles *et al.*, 2001).

The posttranslational responses of non-preferred nitrogen source permeases, such as Gap1, to the presence of preferred nitrogen sources, such as an ammonium compound or glutamine, have been widely documented in *Saccharomyces cerevisiae*, although they are not completely understood (Soetens *et al.*, 2001; Springael *et al.*, 1999; Springael and Andre, 1998). However, the response of a nitrate transporter to the presence of preferred nitrogen sources remains largely unexplored in yeast, filamentous fungi and plants.

In this work, the role of Ynt1 in nitrate, nitrite, chlorate and chlorite transport, its contribution to cell nitrate supply, and the response to the presence of reduced nitrogen are analysed. It is shown that Ynt1, a high-affinity nitrate transporter in fungi, is also able to transport nitrite with high affinity. The presence of glutamine in the medium triggers the inactivation of the nitrate and nitrite transport through Ynt1. Moreover, we show that at least one high-affinity nitrite-specific transporter, inducible by nitrate and not inactivated by glutamine, is present in *H. polymorpha*.

## Materials and methods

### Yeast strains

These are listed in Table 1.

### Growth conditions

Yeast were grown in liquid medium containing 0.17% yeast nitrogen base without ammonium sulphate and amino acids (YNB). The carbon source is indicated in each case. Ammonium chloride, sodium nitrate or sodium nitrite were commonly

**Table 1.** Yeast strains

| Strain        | Genotype  | Source                     |
|---------------|---|----------------------------|
| NCYC 495      | <i>leu1-1, ura3</i>   | Gröningen                  |
| NCYC495 (WT)* | <i>LEU2, URA3</i>   | La Laguna                  |
| MDP100        | $\Delta ynt1::URA3; P_{MOX1}::PET1$<br>( $P_{MOX1}-YNT1-T_{AMO1}$ ) | La Laguna                  |
| MDP 1995      | $\Delta ynr1::URA3$   | Avila <i>et al.</i> (1996) |
| NB 1996       | $\Delta yni1::URA3$   | Brito <i>et al.</i> (1998) |
| NB 1997       | $\Delta ynt1::URA3$   | Pérez <i>et al.</i> (1997) |
| NB 1998       | $\Delta yna1::URA3$   | Avila <i>et al.</i> (1998) |

\* This strain was obtained by transforming the NCYC495 *leu1.1, ura3* with plasmids linearized at the *LEU2* and *URA3* genes.

used as nitrogen sources. Cell growth was followed at OD<sub>660</sub>. Cell density is expressed as mg(wet weight)/ml.

#### Fusion of *YNT1* to *MOX1* promoter

The *H. polymorpha* expression plasmid pET1 (Tan *et al.*, 1995) was used to fuse the *YNT1* ORF to the *MOX1* gene promoter to construct plasmid pMDP16. This plasmid was linearized at the *StuI* site in the *MOX1* promoter to target its integration to the *MOX1* locus of the  $\Delta ynt1::URA3$  strain to obtain MDP100 ( $\Delta ynt1::URA3$ , *MOX1p-YNT1*). Most recombinant DNA procedures were performed according to Sambrook *et al.* (1989).

#### Nitrate and nitrite uptake rate assays

Nitrate or nitrite transport was inferred from extracellular nitrate or nitrite depletion. Nitrate or nitrite uptake rate experiments were carried out at the cell density indicated (1–50 mg/ml) at 37 °C in 0.17% YNB plus 2% glucose and the indicated concentrations of nitrate or nitrite. Modifications made to the media are always indicated. The uptake assay was initiated by the addition of either nitrate or nitrite to the cell suspension. At short time intervals (1–2 min), samples (usually 200  $\mu$ l) were withdrawn and the uptake activity stopped with 0.35 M HCl. Once the assay was finished, samples were neutralized with 1 M Tris and centrifuged. Extracellular nitrate was determined colorimetrically as nitrite, according to Snell and Snell (1948). Nitrate was enzymatically transformed into nitrite. The enzymatic reaction was carried out in a 96-well microtitre plate at 30 °C for 1 h in a 60  $\mu$ l reaction mixture containing: 50 mM Tris–HCl, pH 7.5;  $2.25 \times 10^{-3}$  U *Aspergillus* sp. NR (Roche); 75  $\mu$ M NADPH; and samples with no more than 20 nmol nitrate. The enzymatic nitrate determination method was validated by high performance liquid chromatography (HPLC), which was carried out using an anionic exchange column (IC-pack; A. Waters) and 2.5 mM lithium hydroxide as the eluant. Nitrate and nitrite were detected with an UV detector at 214 nm.

To determine extracellular nitrate at low concentration, samples were concentrated by freeze-drying. The nitrate/nitrite uptake assay conditions were modified to avoid interference with the determination of nitrate, either enzymatically or by

HPLC. Glucose and YNB were reduced to 0.2% and 0.017%, respectively, and cell concentration to 0.5–1 mg/ml. Samples were concentrated between 10- and 40-fold. In nitrite uptake assays, extracellular nitrite was determined directly according to Snell and Snell (1948).

In nitrate and nitrite uptake experiments the medium was buffered at different pH values: pH 3, 4 and 5 with 20 mM tartaric acid–Tris; pH 5.5 and 6 with 20 mM MES–Tris; and pH 7 and 8 with 50 mM Tris–MES. To determine whether any of the buffer solution components interfered with uptake assays, overlapping pH buffers using different components were used in nitrate and nitrite uptake experiments. No significant variations in nitrate and nitrite uptake were observed with regard to the buffer components used (data not shown).

#### Production of a polyclonal antiserum against Ynt1

The region of Ynt1 comprising part of the central hydrophilic loop and the seventh transmembrane span (residues 224–367) was used as an antigen. It was overexpressed in *E. coli* using the pRSETa plasmid (Invitrogen Corporation, San Diego, USA). The antiserum anti-Ynt1 was obtained from a New Zealand rabbit according to Harlow and David (1988).

#### Extraction of membrane proteins

50 mg cells were resuspended in 250  $\mu$ l 50 mM Tris–HCl, pH 7.4, plus protease inhibitor cocktail (Roche). The cell suspension, containing glass beads (0.5 mm diameter), was vortexed five times for 1 min, keeping it on ice for 1 min between each vortex. An additional 250  $\mu$ l resuspension buffer was added, and the sample was vortexed for 10 s. The glass beads and cell debris were discarded by two consecutive centrifugations (3200 rpm, 1 min) in a microfuge. The resulting supernatant was centrifuged at 15 000 rpm for 30 min at 4 °C, and the pellet obtained (particulate fraction) was resuspended in 70  $\mu$ l Triton X-100 0.25 mg/ml. Protein concentration was determined according to Bradford (1977), taking into account Triton X-100 interference.

### SDS-PAGE and Ynt1 immunodetection

20  $\mu$ l of 4 $\times$  sample buffer (12% SDS, 6% mercaptoethanol, 30% glycerol, 0.05% Serva blue G, 150 mM Tris-HCl, pH 7; Schägger; 1994) was added to 60  $\mu$ l particulate fraction resuspended in Triton X-100. Samples were heated to 40 °C for 30 min, 20  $\mu$ g protein per lane was loaded on 12% acrylamide gels. Proteins were transferred to a PVDF membrane (Hybond-P, Amersham Ibérica SA, Spain). Ynt1 immunodetection was carried out using the anti-Ynt1 antiserum (1/1500 dilution), and a goat anti-rabbit IgG antiserum (1/30 000 dilution) conjugated to peroxidase was used as a secondary antibody (Roche). Ynt1 was detected on the blot using the ECL chemiluminescent system (Amersham) according to the manufacturer's instructions.

### Electron microscopy

Whole cells were fixed and prepared for immunocytochemistry as described previously (Waterham *et al.*, 1994). Immunolabelling was performed on ultrathin sections of uncryl-embedded cells, using antiserum against Ynt1 and gold-conjugated goat-anti-rabbit (GAR-gold) antibodies, following the directions of the manufacturer (Amersham, Arlington Heights, IL).


### Other methods

NR activity was determined as described by González and Siverio (1992).

## Results

### Role of Ynt1 in nitrate uptake

To evaluate the contribution of Ynt1 to the total nitrate consumption of *H. polymorpha* cells, nitrate uptake experiments were carried out using wild-type (WT) and  $\Delta ynt1::URA3$  mutant strains. Under the experimental conditions used, nitrate uptake was directly proportional to the number of WT cells present in the assay medium (Figure 1). In contrast, no nitrate uptake was observed with  $\Delta ynt1::URA3$  mutants. Nitrate uptake rate in the WT was approximately 0.5 nmol/min/mg cells. The fact that no nitrate uptake was observed in the *ynt1::URA3* mutant led to the conclusion that at low nitrate




**Figure 1.** Nitrate uptake in WT and  $\Delta ynt1::URA3$  strains. Cells grown in 5 mM ammonium were resuspended at 5 mg/ml in 0.5 mM nitrate for 2 h. The cells were then washed and resuspended in YNB plus 2% glucose; WT at 10 mg/ml (○), 30 mg/ml (●) and at 50 mg/ml (□); and the  $\Delta ynt1::URA3$  mutant at 50 mg/ml (■). Nitrate uptake assays were triggered with 0.5 mM nitrate, samples for measurement of extracellular nitrate were taken at the appropriate times. The experiments were repeated three times without significant differences; the results shown are from a single experiment

concentration, nitrate uptake is mainly attributed to Ynt1. Therefore Ynt1 is the only high-affinity nitrate transporter in *H. polymorpha*. However, this strain is still able to grow in nitrate, although at a much lower growth rate than the WT. Moreover, low-affinity nitrate uptake has been detected in a  $\Delta ynt1::URA3$  strain (Machín *et al.*, 2000).

To ascertain whether Ynt1 is the rate-limiting step in the nitrate assimilation pathway, the relationship between nitrate uptake and Ynt1 levels was determined. To address this question, the MDP100 strain was used. In this strain *YNT1* transcription is under the control of the *MOX1* promoter, which is induced by methanol and repressed by glucose (Gödecke *et al.*, 1994). Ynt1 usually localises in the plasma membrane. It was thought that overexpression of Ynt1 from the *MOX1* promoter could result in erroneous localization of Ynt1 to other cellular compartments. This would lead to misinterpretation of results. To exclude this possibility, Ynt1 was immuno-located by electron microscopy in MDP100 cells. As shown in Figure 2, Ynt1 is located exclusively in the plasma membrane.


To induce *YNT1* independently from the rest of the nitrate assimilatory genes, MDP100 cells were



**Figure 2.** Membrane location of Ynt1 in a strain expressing *YNT1* under the *MOX1* promoter. (A) Survey of methanol-grown MDP100 cells overexpressing Ynt1, labelled with anti-Ynt1/GAR-gold. Labelling is exclusively located at the cell membrane. The high magnification (B) shows the labelled cell membrane in closer detail. M, mitochondrion; P, peroxisome; V, vacuole. The marker represents 0.5  $\mu\text{m}$

incubated in methanol plus ammonium, following incubation in glycerol. To produce increasing amounts of Ynt1, MDP100 cells were incubated in methanol for increasing time intervals. Subsequently, repression of the *MOX1* promoter and induction of NR were carried out by transferring cells to 4% glucose plus 1 mM nitrate for 1 h. Nitrate uptake and NR activity were determined in each sample, as shown in Figure 3. Nitrate uptake was directly proportional to the increasing levels of Ynt1 expressed during methanol induction. Moreover, the variation of the NR activity did not affect the nitrate uptake rate. Therefore, nitrate uptake rate is directly proportional to the amount of Ynt1, and independent of NR levels. A further conclusion made from this experiment, is that extracellular nitrate uptake depletion assays are a good method for determining nitrate transport in *H. polymorpha*.


With regards to Ynt1 affinity for nitrate, no differences in the nitrate depletion rate assays were found using extracellular nitrate concentrations of 25–250  $\mu\text{M}$  (data not shown). This suggests that the  $K_m$  of Ynt1 for nitrate could be lower than 25  $\mu\text{M}$ . The  $K_m$  determined by the Eadie–Hofstee plot was 2  $\mu\text{M}$  nitrate, while the  $V_{\text{max}}$  was around 0.5 nmol  $\text{NO}_3^-/\text{min}/\text{mg}$  cells (Figure 4).



**Figure 3.** Correlation of Ynt1 levels with nitrate uptake rate. In MDP100 cells grown in glucose plus ammonium, *MOX1-YNT1* was derepressed in 0.5% glycerol plus 1 mM ammonium for 3 h, and then induced in 0.5% methanol plus 1 mM ammonium for the indicated times. Samples were harvested by centrifugation, incubated for 1 h in 4% glucose and 0.5 mM nitrate to repress *MOX1-YNT1* and induce nitrate inducible genes, respectively. Subsequently, nitrate uptake (A), nitrate reductase activity (B) and Ynt1 levels (C), were determined for each sample. To measure Ynt1 levels, 10  $\mu\text{g}$  of the particulate fraction was subjected to SDS-PAGE and immunoblotted using anti-Ynt1 antiserum. The experiments were repeated three times; each value is expressed as the mean  $\pm$  SD

### In *H. polymorpha* nitrite is transported by Ynt1 and a nitrite-specific transport system(s)

To study whether Ynt1 was able to transport nitrite, nitrite uptake experiments were carried out using  $\Delta\text{ynr1}::\text{URA3}$  and  $\Delta\text{ynt1}::\text{URA3}$  mutants




**Figure 4.** Kinetic characteristics of Ynt1 nitrate uptake. Wild-type cells grown in 5 mM ammonium were washed and resuspended at 5 mg/ml in 0.5 mM nitrate for 90 min. Cells were then resuspended in 0.2% glucose, 0.017% YNB, 20 mM MES-Tris, pH 5.5, at a cell concentration of 1–10 mg/ml, depending on the extracellular nitrate concentration used. Uptake experiments were triggered with nitrate. Extracellular nitrate depletion was followed for 21 min. Samples were taken every 3 min. The experiments were repeated three times; each value is expressed as the mean  $\pm$  SD

(Figure 5). To avoid nitrite production and excretion (which could interfere with the nitrite uptake assay in the presence of nitrate) a  $\Delta ynr1::URA3$  mutant was used, which expresses WT *YNT1* but lacks NR. The fact that nitrite uptake was three times higher in a  $\Delta ynr1::URA3$  background than in a  $\Delta ynt1::URA3$  background led us to conclude that Ynt1 also transports nitrite. However, since  $\Delta ynt1::URA3$  mutants were shown to uptake nitrite, at least one additional transporter for nitrite is present in *H. polymorpha*.

Further evidence to corroborate the involvement of Ynt1 in nitrite transport was obtained by measuring the effect of nitrate on nitrite transport. The addition of 100  $\mu$ M nitrate inhibited nitrite uptake in YNR1-deleted cells, to the levels of nitrite uptake observed in YNT1-deleted cells in the absence of nitrate (Figure 5). In addition, no inhibition of nitrite uptake by nitrate was observed in  $\Delta ynt1::URA3$  mutants. This suggests specificity for nitrite of the Ynt1-independent nitrite uptake system (s). Our data allow us to conclude that Ynt1 transports nitrate and nitrite, and also strongly points towards the existence of a nitrite-specific uptake system (s).


With regard to the affinity of Ynt1 for nitrite, kinetic experiments showed that the  $K_m$  of Ynt1 for nitrite is in the  $\mu$ M range (Figure 6).




**Figure 5.** Ynt1 transports nitrite.  $\Delta ynt1::URA3$  and  $\Delta ynr1::URA3$  strains were grown in 5 mM ammonium, harvested, washed and resuspended for 2 h in 1 mM nitrate. To determine nitrite uptake, cells were resuspended at 10 mg/ml in a medium buffered with 10 mM MES-Tris, pH 5.5, in the absence and presence of 100  $\mu$ M and 1000  $\mu$ M nitrate. Nitrite uptake assays are triggered with 100  $\mu$ M nitrite.  $\Delta ynr1::URA3$  and  $\Delta ynt1::URA3$  strains are represented with black and white bars, respectively. The experiments were repeated three times; each value is expressed as the mean  $\pm$  SD

#### Characterization of nitrite transport

Figure 7 shows nitrite uptake rates measured at a pH range of 3–7 in WT and  $\Delta ynt1::URA3$  cells. We found that at low pH (3–5), both strains reach



**Figure 6.** Nitrite uptake in WT and  $\Delta ynt1::URA3$  strains at different extracellular nitrite concentrations. Cells grown in 5 mM ammonium were washed and resuspended in 1 mM nitrate, at 10 mg/ml, for 2 h. Cells were then resuspended at the appropriate cell density and nitrite uptake assays were triggered with 15, 25, 50, 100, 200 and 400  $\mu\text{M}$  nitrite. The experiments were repeated three times; each value is expressed as the mean  $\pm$  SD



**Figure 7.** Effect of external pH on nitrite uptake in WT and  $\Delta ynt1::URA3$  strains. Cells were grown in 5 mM ammonium, harvested, washed and resuspended in 1 mM nitrate for 2 h. To determine nitrite uptake, WT and  $\Delta ynt1::URA3$  cells were resuspended at 10 mg/ml in media buffered to pH 3–8. Nitrite uptake assays were triggered with 100  $\mu\text{M}$  nitrite.  $\Delta ynt1::URA3$  and WT strains are represented with black and white bars, respectively. The experiments were repeated three times; each value is expressed as the mean  $\pm$  SD

similar values of nitrite uptake, whereas at pH of around 6, only WT cells were able to transport nitrite efficiently. At pH 4, almost all nitrite transport occurs through the nitrite-specific system(s),

since no important differences between WT and the Ynt1-null mutant cells were observed. However, at pH 6 practically all nitrite transport occurs through Ynt1. The lack of transport observed at pH 3 is most likely due to the uncoupling effect of nitrous acid. However, the incapacity of the nitrite-specific system to take up the protonated form of nitrite cannot be discarded. The optimal pH for the nitrite-specific system (s), pH 4, could also be explained if nitrite uptake at this pH was due to a simple diffusion of the nitrous acid, since nitrous acid presents a pKa of 3.37. However, this possibility is discarded by the fact that, at pH 4, proton ionophores such as 2,4-dinitrophenol (DNP) and carbonyl cyanide *m*-chlorophenylhydrazone (CCCP) strongly impair the nitrite uptake in  $\Delta ynt1::URA3$  cells (Table 2), as expected for a proton symport system. Experiments designed to study the kinetic characteristics of the nitrite uptake system in  $\Delta ynt1::URA3$  mutants showed that at pH 4, the optimal pH for nitrite uptake, the  $K_m$  for nitrite is in the  $\mu\text{M}$  range (Figure 6). Likewise, these experiments also showed that the increase of nitrite concentration to over 100  $\mu\text{M}$  decreases nitrite uptake. This is inconsistent with simple diffusion of nitrite/nitrous acid into the cell. It is possible that at high nitrite concentrations the nitrous acid present at pH 4 could be at a concentration high enough to abolish nitrite transport due to its uncoupling effect.

To study the regulation of the nitrite uptake system(s) induction by nitrogen sources,  $\Delta ynt1::URA3$  cells were grown in ammonium and transferred to media containing either nitrate or ammonium. As shown in Figure 8, nitrite depletion was only observed in cells previously incubated in nitrate-containing medium, unlike those cells incubated for 2 h in 5 mM ammonium. No nitrite uptake at pH

**Table 2.** Effect of proton ionophores on nitrite uptake in the  $\Delta ynt1::URA3$  strain

|                     | Control      | CCCP       | 2,4 DNP   |
|---------------------|--------------|------------|-----------|
| $\Delta ynt1::URA3$ | 100 $\pm$ 16 | 10 $\pm$ 6 | 7 $\pm$ 4 |

The cells were grown in 5 mM ammonium and resuspended at 5 mg/ml in 1 mM nitrate for 2 h. Nitrite uptake assays were performed at pH 4 plus 0.1 mM CCCP or 0.1 mM 2,4-DNP. A control assay was carried out without ionophores. The cells were incubated in these media for 5 min before adding nitrite. The results are expressed as percentage of nitrite uptake rate with respect to the control. The experiments were repeated three times; each value is expressed as the mean  $\pm$  SD




4 was observed with *yna1* mutants, despite prior incubation in a nitrate-containing medium. This indicates that the induction of the nitrite uptake system (s) is also mediated by the  $Zn(II)_2Cys_6$  transcription factor *YNA1*, which is involved in nitrate and nitrite induction of the nitrate assimilatory genes (Avila et al., 1998).

#### Effects of chlorate and chlorite on nitrate/nitrite uptake systems

Due to the importance of chlorate as a tool to study the nitrate assimilation pathway, we have determined whether chlorate/chlorite can be transported by any of the nitrate/nitrite transport systems in *H. polymorpha*.

To address the question of whether Ynt1 was able to transport chlorate, nitrate uptake was assayed at extracellular nitrate concentrations of 10  $\mu M$  and 100  $\mu M$ . Assays were performed in the presence of chlorate at the same concentration, or five-fold higher than the extracellular nitrate concentration used. Under these conditions nitrate uptake was not affected by chlorate (results not



**Figure 8.** Nitrate induction and ammonium repression of the nitrite transport system.  $\Delta ynt1::URA3$  and  $\Delta yna1::URA3$  strains were grown in 5 mM ammonium. Cells were harvested, and resuspended at 10 mg/ml for 2 h;  $\Delta ynt1::URA3$  cells in 1 mM nitrate ( $\circ$ ), or 5 mM ammonium ( $\bullet$ ) and  $\Delta yna1::URA3$  cells in 5 mM ammonium ( $\blacklozenge$ ). Cells were then harvested and resuspended at 5 mg/ml in YNB plus glucose, buffered at pH 4 with 20 mM tartaric acid-Tris; nitrite uptake assays were triggered with 100  $\mu M$  nitrite. The experiments were repeated three times without significant differences. Results from a single experiment are shown


shown). Similar results were observed with WT and  $\Delta ynt1::URA3$  mutant cells in the presence of chlorate. As expected,  $\Delta yna1::URA3$  cells, in which the *YNR1* gene is not expressed, were much more resistant to chlorate (Figure 9). Furthermore, chlorate does not affect nitrite uptake at pH 4 in a  $\Delta ynt1::URA3$  background (data not shown). This suggests that the nitrite-specific system does not transport chlorate.

The effect of chlorite on Ynt1 nitrate/nitrite transport and on the nitrite-specific system was assayed. In addition, the effect of chlorite on the growth of WT cells, *ynt1* and *yna1* mutant cells cultured on nitrate-containing media over a range pH was measured. It was observed that all strains show a very similar sensitivity to chlorite, unlike chlorate, thus indicating that Ynt1, nitrite and NR are not involved in the sensitivity to chlorite. Furthermore, no differences between WT and  $\Delta ynt1::URA3$  strains were found at either pH 4 or pH 6 (Figure 9). Chlorite inhibition assays carried out on Ynt1 nitrite/nitrate uptake at pH 6, and in the *YNT1*-null mutant strain at pH 4, showed a significant drop in nitrite uptake in both strains (data not shown). This indicates that both systems (Ynt1 and the nitrite-specific system) might transport chlorite. However, the non-specific toxic effect of chlorite on the cells, consistent with that observed in the growth experiments, casts doubt on the capacity of Ynt1 and the nitrite-specific system to transport chlorite.

#### Glutamine inactivates Ynt1 nitrate/nitrite uptake but not nitrite-specific uptake system(s)

To study further the characteristics of nitrate and nitrite transport in *H. polymorpha*, the posttranslational response to the presence of reduced nitrogen sources was explored. We studied the effect of glutamine on cells grown in nitrate. Glutamine brought about a rapid inactivation of nitrate and nitrite uptake, and the degradation rate of Ynt1 increased notably (Figure 10A, B). Thus far, it is not clear whether the inactivation of the nitrate/nitrite uptake is a direct consequence of the proteolytic degradation of Ynt1, or whether covalent modifications of Ynt1 are also involved. To confirm that the inactivation of nitrate uptake is targeted to Ynt1, NR activity was measured during the experiment. No NR inactivation was observed (Figure 10A).

The effect of glutamine on the nitrite-specific system was also studied (Figure 10C). The  $\Delta ynt1::$



**Figure 9.** Effect of chlorate and chlorite on the growth of WT,  $\Delta ynt1::URA3$  and  $\Delta yna1::URA3$  strains. Cells grown in 5 mM ammonium were washed thoroughly and  $10^7$ ,  $10^6$ ,  $10^5$  and  $10^4$  cells were spotted onto solid medium containing 2% glucose, 0.17% YNB, 50 mM sodium nitrate, 0.2 mM urea and either chlorate or chlorite at the indicated concentrations. Chlorite-containing media were buffered at pH 5 with 20 mM tartaric acid-Tris and at pH 6.5 with 20 mM MES-Tris. The cells were grown for 3 days


*URA3* mutant was grown in ammonium, and the nitrite-specific system was induced with nitrate. Glutamine did not inactivate the nitrite-specific system. This suggests that the nitrite-specific system is not post-translationally regulated. This is in contrast to the Ynt1 transporter, at least in the presence of glutamine.

## Discussion

In this work the role of Ynt1 in nitrate and nitrite transport has been evaluated. Nitrate uptake was measured in WT and  $\Delta ynt1::URA3$  strains. It was concluded that the uptake of nitrate takes place mainly through Ynt1. The dependence of nitrate uptake on Ynt1 and NR levels was also studied. As shown in Figure 2, nitrate uptake is dependent on the levels of Ynt1, and is independent of NR levels. Therefore, under the experimental conditions

employed here, extracellular nitrate depletion can be considered as a measure of Ynt1-dependent nitrate transport. However, this method does not allow us to study nitrate transport for short time intervals. Using the developed method, a  $K_m$  for nitrate in the  $\mu M$  range was found for Ynt1. This allows us, based on kinetic data, to include Ynt1 in the high-affinity nitrate transporter group found in fungi, algae and plants (Forde, 2000).

A property of Ynt1 that also required investigation is its ability to transport nitrite, chlorate and chlorite. Two lines of evidence were considered, and it was concluded that Ynt1 transports nitrite. The first piece of evidence was the decrease of nitrite uptake in  $\Delta ynt1::URA3$  cells when compared with the WT. The second is the inhibition of nitrite uptake to those levels observed in the  $\Delta ynt1::URA3$  strain by nitrate. The reduction of nitrite uptake in the  $\Delta ynt1::URA3$  strain could also



**Figure 10.** Effect of glutamine on nitrate and nitrite uptake in WT and  $\Delta ynt1::URA3$  strains. (A) WT cells grown in 5 mM ammonium were resuspended at 5 mg/ml in YNB plus 2% glucose and 0.5 mM nitrate for 2 h, and were then subjected to nitrogen starvation for 1 h. The cells were then split into two aliquots and nitrate uptake was triggered with 0.25 mM nitrate. 10 min after nitrate uptake assay was initiated, 5 mM glutamine was added ( $\downarrow$ ) to one of the cultures. Nitrate depletion was followed in the presence ( $\bullet$ ) and absence ( $\circ$ ) of glutamine. Nitrate reductase activity in nmol nitrite/min/mg protein was determined at 15, 30 and 45 min after starting the nitrate uptake assay. (B) Ynt1 levels in the presence and absence of glutamine. 10  $\mu$ g protein from the particulate fraction was subjected to SDS-PAGE and immunoblot. (C)  $\Delta ynt1::URA3$  cells were treated as in (A), except that the cell suspension was buffered at pH 5 and nitrite uptake was determined. The assay was triggered with 100  $\mu$ M nitrite. The experiments were repeated three times without significant differences. Results from a single experiment are shown

be explained if the Ynt1-independent nitrite transporter were inefficiently induced by nitrate in a *ynt1* mutant, as previously shown for NR (Pérez *et al.*, 1997). However, no increase of nitrite consumption in the  $\Delta ynt1::URA3$  mutant was observed when higher nitrate concentrations were used in the medium to induce equivalent levels of NR between this strain and the WT (results not shown). Therefore, our results strongly suggest that Ynt1 transports nitrite, in addition to nitrate. This is the case for *C. reinhardtii* nitrate/nitrite transporters: CrNRT2.1 and CrNRT2.2 (Galván *et al.*, 1996). The capacity of Ynt1 to transport nitrate and nitrite is consistent with that shown for *A. nidulans* NrtA in *Xenopus* oocyte experiments (Zhou *et al.*, 2000). However, it has been claimed that NrtA and NrtB do not transport nitrite because an *A. nidulans* mutant strain lacking NrtA and NrtB is able to grow as efficiently in nitrite as the WT (Unkles *et al.*, 2001). This does not rule out a role for NrtA and NrtB in nitrite transport, as the ability of these

transporters to transport nitrite could be substituted for by additional nitrite transporters.

The presence of nitrite uptake in the  $\Delta ynt1::URA3$  strain leads us to conclude that at least one additional nitrite transport system is present in this yeast. In contrast to the observations made with the Ynt1 transporter, no inhibition of nitrite uptake by nitrate is observed with the Ynt1-independent transporter. Therefore, it seems likely that the Ynt1-independent transporter is specific for nitrite. At pH 4, nitrite uptake occurs mainly via the nitrite-specific transporter(s), whereas at pH 6 nitrite transport occurs mainly through Ynt1. Proton ionophores such as DNP and CCCP inhibit nitrite uptake in *ynt1* mutants. This highlights the proton symport nature of the nitrite-specific transporter(s) and discounts the hypothesis for simple diffusion of nitrous acid across the plasma membrane at pH 4. The nature of the transporter(s) involved in nitrite uptake is unknown. However, the gene(s) encoding this transport system have no significant homology

with *YNT1*, since attempts made to isolate genes with *YNT1* homology by either DNA hybridization or PCR were unsuccessful.

Chlorate has been widely used to obtain mutants of the nitrate assimilation pathway, mainly in NR, since this enzyme is able to reduce chlorate to the very toxic compound chlorite. It has been proposed, in at least in two cases, that chlorate is transported into the cell through nitrate transporters, first by CHL1, a low-affinity nitrate transporter from *A. thaliana* (Tsay *et al.*, 1993), and second by NrtA from *A. nidulans* (Unkles *et al.*, 1991, 2001). However, Ynt1 does not seem to be involved in chlorate transport. Two facts support this conclusion: first, chlorate does not inhibit nitrate uptake; and second, there are no differences in growth between WT and  $\Delta ynt1$  mutants when chlorate is present in the growth medium. In contrast to the *H. polymorpha ynt1* mutant described in this study, an *A. nidulans* mutant lacking NrtA showed resistance to chlorate, thus suggesting the capacity of NrtA to transport chlorate (Unkles *et al.*, 2001). However, expression of NrtA in *Xenopus* oocytes suggests that NrtA is involved in chlorite transport instead of chlorate (Zhou *et al.*, 2000). As a result, we also decided to investigate the effects of chlorite on *H. polymorpha*. Our results show that chlorite, or any compound present as a contaminant, inhibited growth regardless of the strain examined (WT,  $\Delta ynt1::URA3$  or  $\Delta yna1::URA3$ ) (Figure 9). Although nitrate and nitrite uptake are decreased in the presence of chlorite, no conclusions can be drawn.

The effects of glutamine on nitrate and nitrite uptake by Ynt1-dependent, and nitrite-specific transport systems were investigated. Our results clearly show a distinctive response of Ynt1-dependent uptake to reduced nitrogen sources. This system is inactivated by glutamine. However, the nitrite-specific system(s) is not. A possible explanation for this is that the nitrite system is only subject to transcriptional, and not post-translational, regulation, unlike the Ynt1-dependent system. With regard to the inactivation of nitrate and nitrite uptake by Ynt1, it could be argued that the observations could be due to the inactivation of NR. However, neither ammonium or glutamine inactivate NR (Navarro *et al.*, 2003). Further support for inactivation of Ynt1, but not NR by glutamine, is the observation that a strain expressing *YNRI* under the control of the *MOXI* promoter shows

NR activity, even in an ammonium-containing medium (Navarro *et al.*, 2003). In conclusion, our results strongly suggest that Ynt1 undergoes inactivation in response to reduced nitrogen sources. Total inactivation of nitrate uptake by ammonium or glutamine has been reported in the yeasts *Sporobolomyces roseus* and *Rhodotorula glutinis*. In contrast, partial inactivation has been found in *Candida nitratophila* and *C. utilis*. However, the nitrate assimilation pathway component responsible for inactivation of nitrate uptake has not been investigated further in these yeasts (Hipkin, 1989).

The results presented here go some way to bridge the gaps in our understanding between the physiological studies and the molecular basis of nitrate and nitrite transport in yeast. However, among the elements still unknown are genes that encode components of the nitrite-specific uptake system and the low-affinity nitrate uptake system. The discovery and subsequent cloning and investigation of these genes will broaden our overall understanding of nitrate and nitrite transport in yeast. The importance of post-translational regulation of nitrate and nitrite transport in both yeast and plants remains unclear. *H. polymorpha* could play an important role as a model system for understanding these processes.

#### Acknowledgements

We thank Carlos Gancedo (IIB, Madrid) for many fruitful discussions. This work was supported by grants from the Ministerio de Ciencia y Tecnología (BMC2001.3719) and Gobierno de Canarias (PI2001/050) to J.M.S. F.J.N. and P.T. were recipients of predoctoral fellowships from the Ministerio de Educación Cultura y Deporte (Spain) and F.M. and B.M. from the Gobierno de Canarias.

#### References

- Avila J, González C, Brito N, Machín F, Pérez MD, Siverio JM. 2002. A second  $Zn(II)_2Cys_6$  transcriptional factor encoded by the *YNA2* gene is indispensable for the transcriptional activation of the genes involved in nitrate assimilation in the yeast *Hansenula polymorpha*. *Yeast* **19**: 537–544.
- Avila J, González C, Brito N, Siverio JM. 1998. Clustering of the *YNA1* gene encoding a  $Zn(II)_2Cys_6$  transcriptional factor in the yeast *Hansenula polymorpha* with the nitrate assimilation genes *YNT1*, *YNI1* and *YNRI*, and its involvement in their transcriptional activation. *Biochem J* **335**: 647–652.
- Avila J, Pérez MD, Brito N, González C, Siverio JM. 1995. Cloning and disruption of *YNRI* gene encoding the nitrate reductase apoenzyme of the yeast *Hansenula polymorpha*. *FEBS Lett* **366**: 137–142.

- Bradford MM. 1977. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248–254.
- Brito N, Avila J, Pérez MD, González C, Siverio JM. 1996. The genes *YNII* and *YNRI*, encoding nitrite reductase and nitrate reductase respectively in the yeast *Hansenula polymorpha*, are clustered and coordinately regulated. *Biochem J* **317**: 89–95.
- Crawford NM, Glass AD. 1998. Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant Sci* **3**: 389–395.
- Daniel-Vedele F, Filleur S, Caboche M. 1998. Nitrate transport: a key step in nitrate assimilation. *Curr Opin Plant Biol* **1**: 235–239.
- Forde B. 2000. Nitrate transporters in plants: structure, function and regulation. *Biochim Biophys Acta* **1465**: 219–235.
- Galván A, Fernández E. 2001. Eukaryotic nitrate and nitrite transporters. *Cell Mol Life Sci* **58**: 225–233.
- Galván A, Quesada A, Fernández E. 1996. Nitrate and nitrite are transported by different specific transport systems and by a bispecific transporter in *Chlamydomonas reinhardtii*. *J Biol Chem* **271**: 2088–2092.
- Gödecke S, Eckart M, Janowicz ZA, Hollenberg CP. 1994. Identification of sequences responsible for transcriptional regulation of the strongly expressed methanol oxidase encoding gene in *Hansenula polymorpha*. *Gene* **139**: 35–42.
- González C, Siverio JM. 1992. Effect of nitrogen source on the levels of nitrate reductase in the yeast *Hansenula anomala*. *J Gen Microbiol* **138**: 1445–1451.
- Harlow E, David L. 1988. *Antibodies. A Laboratory Manual*. Cold Spring Harbor Laboratory Press: New York.
- Hipkin CR. 1989. Nitrate assimilation in yeast. In *Molecular and Genetic Aspects of Nitrate Assimilation*, Wray JL, Kinghorn JR (eds). Oxford Scientific: Oxford; 51–68.
- Machín F, Perdomo G, Pérez MD, Brito N, Siverio JM. 2000. Evidence for multiple nitrate uptake systems in *Hansenula polymorpha*. *FEMS Microbiol Lett* **194**: 171–174.
- Navarro F, Perdomo G, Tejera P, et al. 2003. The role of the nitrate reductase in the regulation of the nitrate assimilation pathway in the yeast *Hansenula polymorpha*. *FEMS Yeast Res* **4**: 149–155.
- Navarro MT, Guerra E, Fernández E, Galván A. 2000. Nitrite reductase mutants as an approach to understanding nitrate assimilation in *Chlamydomonas reinhardtii*. *Plant Physiol* **122**: 283–290.
- Pao SS, Paulsen IT, Saier MH. 1998. Major facilitator superfamily. *Micobiol Molec Biol Rev* **62**: 1–34.
- Pérez MD, González C, Brito N, Siverio JM. 1997. The *YNTI* gene encoding the nitrate transporter in the yeast *Hansenula polymorpha* is clustered with genes *YNII* and *YNRI* encoding nitrite reductase and nitrate reductase, and its disruption causes inability to grow in nitrate. *Biochem J* **321**: 397–403.
- Quesada A, Galván A, Fernández E. 1994. Identification of nitrate transporter genes in *Chlamydomonas reinhardtii*. *Plant J* **5**: 407–419.
- Rexach J, Montero B, Fernández E, Galván A. 1999. Differential regulation of the high affinity nitrite transport systems III and IV in *Chlamydomonas reinhardtii*. *J Biol Chem* **274**: 27 801–27 806.
- Sambrook J, Fritsch EF, Maniatis T. 1989. *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbor Laboratory Press: New York.
- Schägger H. 1994. Denaturing electrophoretic techniques. In *A Practical Guide to Membrane Protein Purification*, von Jagow G, Schägger H (eds). Academic Press: San Diego, CA; 59–79.
- Snell FD, Snell CT. 1948. *Colorimetric Methods of Analysis*. Van Nostrand: New York.
- Soetens O, De Craene JO, Andre B. 2001. Ubiquitin is required for sorting to the vacuole of the yeast general amino acid permease, Gap1. *J Biol Chem* **276**: 43 949–43 957.
- Springael JY, Galan JM, Haguenaer-Tsapis R, Andre B. 1999.  $\text{NH}_4^+$ -induced downregulation of the *Saccharomyces cerevisiae* Gap1p permease involves its ubiquitination with lysine-63-linked chains. *J Cell Sci* **112**: 1375–1383.
- Springael JY, Andre B. 1998. Nitrogen-regulated ubiquitination of the Gap1 permease of *Saccharomyces cerevisiae*. *Mol Biol Cell* **9**: 1253–1263.
- Tan X, Waterham HR, Veenhuis M, Cregg JM. 1995. The *Hansenula polymorpha* *PER8* gene encodes a novel peroxisomal integral membrane protein involved in proliferation. *J Cell Biol* **128**: 307–319.
- Tsay Y-F, Schroeder JI, Feldman KA, Crawford NM. 1993. The herbicide-sensitive gene *CHL1* of *Arabidopsis* encodes a nitrate-inducible nitrate transporter. *Cell* **72**: 705–713.
- Unkles SE, Zhou D, Siddiqi MY, Kinghorn JR, Glass ADM. 2001. Apparent genetic redundancy facilitates ecological plasticity for nitrate transport. *EMBO J* **20**: 6246–6255.
- Unkles SE, Hawker KL, Grieve C, et al. 1991. *crnA* encodes a nitrate transporter in *Aspergillus nidulans*. *Proc Natl Acad Sci USA* **88**: 204–208.
- Waterham HR, Titorenko VI, Haima P, et al. 1994. The *Hansenula polymorpha* *PER1* gene is essential for peroxisome biogenesis and encodes a peroxisomal matrix protein with both carboxy- and amino-terminal targeting signals. *J Cell Biol* **127**: 737–749.
- Zhou J-J, Trueman LJ, Boorer KJ, et al. 2000. A high affinity fungal nitrate carrier with two transport mechanisms. *J Biol Chem* **275**: 39 894–39 889.