





The Influence of High Load Training on Reaction Time in Cyclists

Nederhof, Esther; Lemmink, Koen A.P.M.; Zwerver, Johannes; Meeusen, Romain

Published in: MEDICINE AND SCIENCE IN SPORTS AND EXERCISE

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version Early version, also known as pre-print

Publication date: 2006

Link to publication in University of Groningen/UMCG research database

Citation for published version (APA): Nederhof, E., Lemmink, K. A. P. M., Zwerver, H. J., & Meeusen, R. (2006). The Influence of High Load Training on Reaction Time in Cyclists. MEDICINE AND SCIENCE IN SPORTS AND EXERCISE, 38(5), S380-S380.

Copyright Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): http://www.rug.nl/research/portal. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Medicine & Science in Sports & Exercise

Issue: Volume 38(5) Supplement, May 2006, p S380

Copyright: ©2006The American College of Sports Medicine

Publication Type: [Thursday Morning Poster Presentations: Posters displayed from 7:30 a.m.-12:30 p.m.: One-hour author presentation times are staggered from 8:30–9:30 a.m. and 9:30–10:30 a.m.: D-26 Free Communication/Poster – Reaction Time: THURSDAY, JUNE 1, 2006 2:00 PM – 5:00 PM ROOM: Hall B]

ISSN: 0195-9131

Accession: 00005768-200605001-02482

[Thursday Morning Poster Presentations: Posters displayed from 7:30 a.m.–12:30 p.m.: One-hour author presentation times are staggered from 8:30–9:30 a.m. and 9:30–10:30 a.m.: D-26 Free Communication/Poster – Reaction Time: THURSDAY, JUNE 1, 2006 2:00 PM – 5:00 PM ROOM: Hall B]

The Influence of High Load Training on Reaction Time in Cyclists: 2148: Board #85 2:00 PM – 3:00 PM

Nederhof, Esther¹; Lemmink, Koen A.P.M.¹; Zwerver, Hans J.²; Meeusen, Romain FACSM³

Author Information

¹Center for Human Movement Sciences, University Medical Center Groningen, University of Groningen, The Netherlands.

²Sports Medicine, University Medical Center Groningen, Groningen, The Netherlands.

³Department of Human Physiology and Sports Medicine, Free University, Brussels, Belgium. Email: e.nederhof@rug.nl

Overtraining syndrome (OTS) is seen as a serious threat by athletes and coaches because performance is deteriorated in OTS. Central fatigue has also been mentioned as an accompanying symptom of OTS which could possibly be measured through tasks of psychomotor speed. Since the only cure to OTS is (relative) rest, it is of utmost importance to prevent it.

RESULTS: of studies into markers have been contradictory, partly because researchers who study the effects of high load training claim to study OTS whereas these athletes were at most functional overreached (FO; Meeusen et al., In Press).

PURPOSE: The purpose of the present study is investigate whether changes in psychomotor speed are already present in early stages of overtraining.

METHODS: Fourteen cyclists have completed an incremental exercise test, two questionnaires and the finger pre-cueing task (FPT) three times: before, right after and two weeks after a training camp. Maximal work load, heart rate, oxygen uptake and mood states were used to determine training status. The FPT is a complex four-choice reaction time task in which pre-cues reduce the task to a two-choice reaction time task in three out of four conditions (Miller, 1982). A control group of fourteen age and gender matched active individuals completed the FPT at the same time as the cyclists.

RESULTS: Five out of fourteen cyclists showed performance decrements and worsened mood states and were classified as overreached. Because performance and mood states had improved two weeks after the training camp, their status was specified as FO. Seven athletes did not show differences on performance or mood states after the training camp and were classified as well-trained (WT). Two athletes showed disturbed mood states before but not after the training camp and were excluded from analysis. A repeated measures ANO VA showed no significant difference between the FO, the WT and the control group. However, the interaction between time and group showed a trend (F = 2,30, p = .079). The FO group showed longer reaction times compared to the control group right after the training camp.

CONCLUSIONS: The RESULTS show that psychomotor slowness as an indication of central fatigue is not present in FO. This could be because FO is part of normal training and signs and

symptoms of OTS are still mild. Future research should point out whether psychomotor slowness is present in non-functional overreaching and/or OTS. Meeusen R, Duclos M, Gleeson M, et al. Prevention, diagnosis and treatment of the overtraining syndrome. Eur J Sport Sci; In Press.Miller J. Discrete versus continuous stage models of human information processing: In search of partial output. J Exp Psychol1982;8: 273–96.

Copyright (c) 2000-2011 <u>Ovid Technologies, Inc.</u> <u>Terms of Use Support & TrainingAbout UsContact Us</u> Version: OvidSP_UI03.04.02.112, SourceID 54875