

University of Groningen

The Photosystem II Light-Harvesting Protein Lhcb3 Affects the Macrostructure of Photosystem II and the Rate of State Transitions in Arabidopsis

Damkjaer, Jakob T.; Kereiche, Sami; Johnson, Matthew P.; Kovacs, Laszlo; Kiss, Anett Z.; Boekema, Egbert J.; Ruban, Alexander V.; Horton, Peter; Jansson, Stefan; Kereiche, Sami

Published in:
 Plant Cell

DOI:
[10.1105/tpc.108.064006](https://doi.org/10.1105/tpc.108.064006)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Damkjaer, J. T., Kereiche, S., Johnson, M. P., Kovacs, L., Kiss, A. Z., Boekema, E. J., ... Kereiche, S. (2009). The Photosystem II Light-Harvesting Protein Lhcb3 Affects the Macrostructure of Photosystem II and the Rate of State Transitions in Arabidopsis. *Plant Cell*, 21(10), 3245-3256. DOI: 10.1105/tpc.108.064006

Copyright


Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.


Supplemental Data Damkjær et al (2009). The Photosystem II light harvesting protein Lhcb3 affects the macrostructure of photosystem II and the rate of state transitions in *Arabidopsis*



Supplemental Figure 1: NPQ induction and dark relaxation in wild type and *koLhcb3* *Arabidopsis* plants. NPQ was measured at 1000 microE actinic light treatment in wild type (closed symbols, full line) and *koLhcb3* (open symbols, dashed line) *Arabidopsis* plants. Error bars show SE, n=5. The grey bars show when the actinic light is on and black bars show when actinic light is off.




Supplemental Figure 2. State 1 to State 2 transition induced by PSII light treatment after 15 minutes PSI light (far red) illumination (n=6). The upper (black) trace shows wild type, the two lower (grey) trace shows koLhcb3 lines N520342 and N661731.




Supplemental Figure 3. In vivo LHCII phosphorylation in wild type (white bars) and koLhcb3 lines N520342 and N661731 (grey bars) *Arabidopsis* thylakoids isolated following State 1 and State 2 inducing light treatments. Results are normalised to wild type state 1 phosphorylation, corrected relative to the CP47 phosphorylation signal. Error bars show SE, n= 13

Supplemental Figure 4. Linear range of the Lhcb1 antibody signal.



Supplemental Figure 5. Linear range of the Lhcb2 antibody signal.



Supplemental Table 1: ANOVA analysis of the significance of the interaction between genotype (Wild type or *koLhcb3*) and treatment (State 1 or State 2 light) with regard to PSII antennae size measured using maximal fluorescence and 15 minute light treatments. ** indicate statistically significant differences ($p < 0.01$) and *** statistically significant differences ($p < 0.001$) using the ANOVA test for variance, n=13.

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
Genotype	1	0.00119	0.00119	0.5855	0.451053	
State	1	0.097667	0.097667	48.0731	2.318E-07	***
Sample date	3	0.037241	0.012414	6.1102	0.002741	**
Genotype:state	1	0.001824	0.001824	0.8979	0.352082	
Residuals	26	0.052822	0.002032			

Supplemental Table 2: ANOVA analysis of the significance of the interaction between genotype (wild type or *koLhcb3*) and treatment (State 1 or State 2 light) with regard to LHCI phosphorylation. ** indicate statistically significant differences ($p < 0.01$) and *** statistically significant differences ($p < 0.001$) using the ANOVA test for variance, $n=27$.

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
Genotype	1	4.8327E+11	4.8327E+11	46.539	5.572E-10	***
State	1	2.0593E+12	2.0593E+12	198.3116	< 2.2e-16	***
Prep Date	1	6.6701E+12	6.6701E+12	642.3288	< 2.2e-16	***
Sample Date	2	5.1757E+11	2.5878E+11	24.9209	1.303E-09	***
Genotype:State	1	8.9476E+10	8.9476E+10	8.6165	0.004078	**
Residuals	107	1.1111E+12	1.0384E+10			

Supplemental Table 3. Seed set of wild type and *koLhcb3* Arabidopsis plants in the field.

	Wildtype	<i>koLhcb3</i>	t-test p-value
Mean siliques per plant	149 ± 13	152 ± 14	0.8884
Mean seeds per siliques	52.1 ± 0.9	49.3 ± 1.2	0.0623
Mean seeds per plant	7783 ± 682	7683 ± 807	0.9226