

University of Groningen

Peroxisomes are required for efficient penicillin biosynthesis in *Penicillium chrysogenum*

Meijer, Wiebe H.; Gidijala, Loknath; Fekken, Susan; Kiel, Jan; van den Berg, Marco A.; Lascaris, Romeo; Bovenberg, Roelof; van der Klei, Ida

Published in:
Applied and Environmental Microbiology

DOI:
[10.1128/AEM.02327-09](https://doi.org/10.1128/AEM.02327-09)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Meijer, W. H., Gidijala, L., Fekken, S., Kiel, J. A. K. W., van den Berg, M. A., Lascaris, R., ... van der Klei, I. J. (2010). Peroxisomes are required for efficient penicillin biosynthesis in *Penicillium chrysogenum*. *Applied and Environmental Microbiology*, 76(17), 5702-5709. DOI: 10.1128/AEM.02327-09

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

