

University of Groningen

Aluminium tolerance of Mucuna

Hairiah, Kurniatun

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1992

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hairiah, K. (1992). Aluminium tolerance of Mucuna: A tropical leguminous cover crop s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUMMARY

In the humid tropics leaching of N and other nutrients to the subsoil may occur throughout the growing season. Typically, soils in this zone have a low soil pH, a high Al saturation of the cation exchange complex and low levels of Ca and P in the subsoil. Efficiency of N-use under such conditions generally is very low, due to a combination of high leaching rates and shallow root development. The ability to develop a deep root system is important in this situation for recovery of nutrients leached early on.

From a number of leguminous cover crops tested, the velvet bean *Mucuna pruriens* var *utilis* gave the highest biomass production on acid soils in Nigeria and Indonesia, and it was the most suitable in control of the weed *Imperata cylindrica*. However, it had a very superficial root system. Inhibition of root growth on acid soil is usually caused by Al toxicity, but other soil factors, such as mechanical impedance or poor aeration might be involved as well.

Laboratory experiments showed that the response of roots of *Mucuna pruriens* var. *utilis* to Al depended on the duration of exposure and the type of experiment. In a short-term (24-48 hours) study, the usual reduction of root elongation rate by Al was found. In contrast, solution culture studies (pH 4.2; NO_3^- as N source) over a longer period of time (4 weeks) showed that *M. p. utilis* and the related species *M. deerin-giana*, had an increased root dry weight at 110 μM or 185 μM Al^{3+} , compared to controls without Al. The moderate Al-tolerance found in solution culture contradicted with the shallow root development of *Mucuna* in acid subsoil.

Field tests in Lampung (Indonesia), showed that if both *Mucuna* species was sown directly into the subsoil, after removal of the topsoil, a large root system was formed in the subsoil layers which hardly contained any roots when topsoil was present in the soil profile. Topsoil and subsoil were placed in mesh bags at various positions in the soil profile and various bulk density; *Mucuna* roots avoided the subsoil regardless of position and bulk density. P fertilization and higher liming rates applied in the mesh bag had a positive effect on root length density. Such field evidence suggested a combination of acid-subsoil tolerance (when roots had no choice) and acid-subsoil avoidance (when roots could choose between sub- and topsoil).

The hypothesis that subsoil avoidance was based on an Al avoidance acting at the level of the intact root system rather than individual roots, was tested in split-root

experiments on a source, at pH 4.2, sides, was compared μM applied to both reduced shoot/root containing solution growth to the control Al avoidance reaction to increased branched the plant indeed

In conclusion, strong avoidance a deep root development heterogeneous intact root system

In a split-root preference after containing solution

Total P concentration while the H_2PO_4^- grown on NO_3^- ionic strength of Al content could continuous pH containing Al was less pH tolerance of *Mucuna* about 19 % of Al as Al-PO_4 , this was however, no evidence concentration (with control plants. The solution can be between NO_3^- a major Al tolerance

experiments on a circulated nutrient solution. In such an experiment with NO_3^- as N source, at pH 4.2, the growth of *M. p. utilis* with Al (+/+) or without Al (0/0) in both sides, was compared with that of plants with a choice (0/+). An Al concentration 185 μM applied to both sides of the root system (+/+), increased root dry weight and reduced shoot/root ratio compared to the control (0/0). Application of such an Al-containing solution to half of the root system (0/+), led to a significant shift in root growth to the control side. In further experiments the hypothesis was tested that the Al avoidance reaction was related to the local response of plants under P stress leading to increased branch root development close to a P source. Increasing the P supply to the plant indeed resulted in the disappearance of the Al avoidance.

In conclusion, Al tolerance in a homogeneous medium can be accompanied by a strong avoidance of Al in a heterogeneous root environment. To obtain cultivars with a deep root development in acid subsoils, it is necessary to test for Al-avoidance in a heterogeneous medium as a natural soil. Screening for Al tolerance of single roots or intact root systems in a homogeneous seems inadequate.

In a split-root experiment with NH_4NO_3 (1 : 1) as N source, *Mucuna* roots had no preference after 2 weeks, but after 4 weeks a preferential root development in Al containing solution was found, in contrast to Al avoidance found with NO_3^- nutrition.

Total P concentration in the roots was increased in the presence of Al (185 μM), while the H_2PO_4^- concentration in the root tissue was reduced, when *Mucuna* was grown on NO_3^- nutrient solution with 150 μM P, daily pH adjustment to pH 4.2 and an ionic strength of 8 - 10 mM. It was estimated that at maximum 40 - 60 % of the root Al content could be present as Al- PO_4 precipitates. In subsequent experiments with continuous pH control, the increase in total P content of the roots in solutions containing Al was less pronounced. N form (NO_3^- and NH_4^+) had little effect on the Al tolerance of *Mucuna* in nutrient solution. It was estimated that with NO_3^- nutrition about 19 % of Al in the root might be bound as Al- PO_4 . If all non-ionic P was bound as Al- PO_4 , this would represent 32 % of the Al in the roots. With NH_4^+ -nutrition, however, no evidence of Al- PO_4 precipitation was obtained, as the non-ionic P concentration (with or without Al in the solutions) was similar to that of the NO_3^- -fed control plants. Therefore, Al- PO_4 precipitation in the root of plants grown in NH_4^+ solution can be excluded; as the Al tolerance of *Mucuna* did not differ drastically between NO_3^- and NH_4^+ nutrition, it was concluded that Al- PO_4 precipitation is not a major Al tolerance mechanism for *Mucuna*.

Primary root damage due to Al toxicity occurred in the cell elongation zone, 2-3 mm behind the root tip. Measurements of the external pH on the root surface showed a slight alkalization in the subsoil (from 4.7 to 5.2) and an acidification in the topsoil (from 5.8 to 5.5). The pH around the root tip was independent of the overall external pH, which depended on the $\text{NO}_3^-/\text{NH}_4^+$ uptake ratio. In the (partial) NH_4^+ nutrition, the Al toxicity was low even though the external pH was not increased. Al detoxification by increasing the pH of the external medium was not important for *Mucuna*. Precipitation of polymeric Al forms in the apoplast with a pH of 5.5 or higher may be important.

Suppression of the grass *Imperata* by *Mucuna* mainly depends on intensity and duration of shading. Leaf area index (LAI) of *Mucuna* at 6 weeks after planting was estimated at about 0.8 which corresponds with 50 % light interception. The highest LAI of *M. p. utilis* obtained in 6 weeks was 1.3, and provided an almost complete cover of the soil. It probably reduced light intensity at ground level by less than 80 %. A LAI of 2.6 is required for 90 % light interception. Even for the best growing *Mucuna* it would require more than 6 weeks before sufficient shading effects on *Imperata* would start. As *M. p. utilis* normally dies off after about 4 months, the shading would affect vigor of *Imperata* but certainly not eliminating it. *M. deeringiana*, may be more useful in *Imperata* control because of its long growth cycle. Growth of *Mucuna* in the presence of topsoil was only just enough to affect *Imperata* and any reduction in growth rate due to erosion of the topsoil indicates that *Mucuna* will not be sufficiently effective in *Imperata* control.

In de humide tr
uitspoelen naar
een hoge Al be
heid in de onde
combinatie van
met een diepe t
ingespoelde nu

Van een aar
pruriens var. *ut*
bodem in Niger
alang gras, *Imp*
oppervlakkig w
een gevolg van
zoals een hoge
spelen.

Laboratoriu
Mucuna pruriens
proef van korte
van wortels doe
(pH 4.2; NO_3^- al
soort *M. deering*
 $\mu\text{M Al}^{3+}$, vergel
in voedingsplo
van *Mucuna* in

Veldproever
ondergrond we
groot wortelste
slechts enkele w
ook in een net
op grondsoort e
ondergrond te