

University of Groningen

Transformation of carbon tetrachloride in an anaerobic packed-bed reactor without addition of another electron donor

de Best, JH; Hunneman, P; Doddema, HJ; Janssen, Dick; Harder, W; Doddema, Hans J.

Published in:
Biodegradation

DOI:
[10.1023/A:1008309003059](https://doi.org/10.1023/A:1008309003059)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1999

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

de Best, J. H., Hunneman, P., Doddema, H. J., Janssen, D. B., Harder, W., & Doddema, H. J. (1999). Transformation of carbon tetrachloride in an anaerobic packed-bed reactor without addition of another electron donor. *Biodegradation*, 10(4), 287-295. DOI: 10.1023/A:1008309003059

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Transformation of carbon tetrachloride in an anaerobic packed-bed reactor without addition of another electron donor

Jappe H. de Best^{1*}, Peter Hunneman¹, Hans J. Doddema¹, Dick. B. Janssen² & Wim Harder³

¹TNO Environmental Technology and Process Engineering Division, Department of Environmental Biotechnology, Apeldoorn, The Netherlands; ²Department of Biochemistry, University of Groningen, Groningen, The Netherlands; ³Department of Microbiology, University of Groningen, Groningen, The Netherlands (*author for correspondence: current address: Grontmij A&T, PO Box 119, 3990 DC, Houten, The Netherlands, e-mail: jappe.debest@grontmij.nl)

Accepted 13 July 1999

Key words: anaerobic, carbon tetrachloride, electron donor, metabolic transformation, transformation products

Abstract

Carbon tetrachloride (52 μM) was biodegraded for more than 72% in an anaerobic packed-bed reactor without addition of an external electron donor. The chloride mass balance demonstrated that all carbon tetrachloride transformed was completely dechlorinated. Chloroform and dichloromethane were sometimes also found as transformation products, but neither accumulated to significant levels in comparison to the amount of carbon tetrachloride transformed. Transformation of carbon tetrachloride in the absence of an added electron donor suggests that carbon tetrachloride itself is the source of energy for the biological reaction observed, and possibly the source of carbon for cell growth. No such mechanism is yet known. The pathway of carbon tetrachloride transformation is not clear; it may be dehalogenated by hydrolytic reduction to carbon monoxide or formic acid which are electron demanding transformations. Carbon monoxide or formic acid may be further utilized and serve as electron donor. Complete dechlorination of carbon tetrachloride according to this pathway is independent of a second electron donor or electron acceptor, as with a fermentation process. Vancomycin, an inhibitor of gram positive eubacteria, severely inhibited carbon tetrachloride transformation in batch incubations with an enrichment culture from the reactor, indicating that gram positive eubacteria were involved in carbon tetrachloride transformation. Batch experiments with bromoethanesulfonic acid, used to inhibit methanogens, and molybdate, an inhibitor of sulfate reduction in sulfate reducing bacteria, demonstrated that neither methanogens nor sulfate reducers were involved in the complete dechlorination of carbon tetrachloride.

Introduction

Carbon tetrachloride (CT) is a common water pollutant due to widespread use as an industrial solvent. CT resists aerobic biotransformation, but several authors have described transformation of CT under anaerobic conditions in both continuous flow studies (Bae et al. 1990, Bouwer and McCarty 1983a, Bouwer and Wright 1988, Cobb and Bouwer 1991, Petersen et al. 1994, de Best et al. 1998) and batch culture studies (Bouwer and McCarty 1983b, Criddle et al. 1990, Egli et al. 1987 1988, Gälli and McCarty 1989, Picardal et al. 1995). In all of these studies, CT trans-

formation was described as a cometabolic process. During cometabolic transformation, dehalogenation is not coupled to energy conservation and occurs due to a fortuitous reaction with certain enzymes or coenzymes (Holliger and Schumacher 1994). Cometabolic dehalogenation processes require both an external electron donor and electron acceptor to support growth of the active organisms.

Another microbial process for dehalogenation is dehalorespiration. In this process reductive dehalogenation is coupled to energy conservation. The halogenated compound serves as a terminal electron acceptor and only an external electron donor is required (Holl-

ger and Schumacher 1994, Máymo-Gatell et al. 1997, Mohn and Tiedje 1990). Theoretically, CT can also serve as terminal electron acceptor since the redox potential of redox couples for CT transformation are positive and higher than the redox potential for couples of common electron acceptors such as nitrate, sulfate or carbon dioxide (Vogel et al. 1987).

A third anaerobic dehalogenation process, described for dichloromethane (DCM) (Mägli et al. 1996) and chloromethane (Traunecker et al. 1991), is fermentation. Fermentative dehalogenations, are processes that are carried out in the absence of an externally supplied electron acceptor and with an internally balanced oxidation-reduction reaction.

The aim of this work was to investigate the requirement of CT transformation for electron donors and electron acceptors. CT transformation was studied in an anaerobic continuous-flow packed-bed reactor at different electron donor (acetate) concentrations and in the absence of an additional electron donor and electron acceptor. To obtain more information about the pathway and products of CT transformation, ^{13}C -CT transformation was studied. Finally, the role of methanogenic, acetogenic and sulfate reducing microorganisms in CT transformation is discussed.

Material and methods

Packed-bed reactor studies

The experiments were performed in an upflow packed-bed reactor (glass; height 40 cm; inside diameter 4.6 cm; volume 665 ml) (de Best et al. 1997) packed with polyurethane foam (PUR) particles ($5 \times 5 \times 6$ mm, Bayer B.V., Mijdrecht, the Netherlands) mixed with digested sludge from the wastewater treatment plant Kralingseveer (Rotterdam, The Netherlands). The biomass content added to the reactor was about 0.8 g volatile suspended solids (VSS). The packed-bed reactor was wrapped with aluminum foil to prevent growth of phototrophs.

The reactor was continuously fed over a period of 5 years with an anaerobic non-sterile phosphate and bicarbonate buffered mineral medium that contained no organics or yeast extract (de Best et al. 1997). The medium was continuously purged with oxygen free N_2/CO_2 (99.5/0.5%) to remove all oxygen. The medium (pH 7.3 ± 0.2) was pumped into the reactor by means of a peristaltic pump with marprene tubing. All other tubing was either viton or Teflon. CT, acetate

(CH_3COOH) and Na_2S ($42 \mu\text{M}$, to maintain reducing conditions) were added to the medium as a concentrated solution at the influent of the reactor with a syringe pump. Although the medium contained no added sulfate, about 0.15 mM sulfate was present in the influent of the reactor, probably due to oxidation of sulfide by oxygen permeating the tubing. The hydraulic retention time in the packed-bed was 24 h, based on void volume (flow rate 0.411/day). All experiments were carried out at 25°C .

^{13}C -carbon tetrachloride studies

Experiments with ^{13}C -CT were done in the packed-bed reactor under the same conditions as described above. ^{13}C -CT ($56 \mu\text{M}$), together with acetate ($30 \mu\text{M}$) and Na_2S ($42 \mu\text{M}$), was added to the medium from a concentrated stock solution at the influent of the reactor with a syringe pump. After two weeks of adaptation, daily samples (5 ml) were taken at the influent and effluent (40 cm) of the reactor and at two sample ports at different heights along the reactor (20, 30 cm) for a period of 20 days. Samples were frozen until used.

Immediately after the 20 daily samples taken at the same sample port were pooled, the pH was adjusted to a value below the pKa (4.77) of acetic acid. Each sample was saturated with NaCl and extracted with diethylether (1 : 1). The extract was concentrated by a factor 10 and analyzed for acetic acid and formic acid. CT was analyzed according to the method described at the analytical methods.

Batch culture studies

Batch incubations were done using an anaerobic minimal medium containing (per liter of demineralized water) 80.1 mg $(\text{NH}_4)_2\text{HPO}_4$, 200 mg $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1 mg resazurin and 5 ml trace element solution (de Best et al. 1997). The medium was purged with a mixture of CO_2 and N_2 (0.5 : 99.5 v/v%, 700 ml/min) for 45 min. After purging, $\text{Na}_2\text{S} \cdot 9\text{H}_2\text{O}$ (91 mg/l) and NaHCO_3 (100 mg/l) were added.

The medium was transferred to 120 ml bottles (brown glass) in an anaerobic glove-box. Each bottle contained 60 ml of medium and was closed with Teflon-lined butyl rubber stoppers and aluminum crimp seals. After sterilization, CT ($6.3 \mu\text{M}$) and acetate ($100 \mu\text{M}$) were added as concentrated solutions. All cultures were inoculated with 2 ml of the liquid phase taken from the packed-bed reactor. The cultures were incubated on a shaker (100 rpm) in a canted

position at 25 °C and analyzed regularly for chlorinated compounds (taking into account the partitioning), acetate, sulfate, CO₂, CO, CH₄ and chloride.

To investigate the role of methanogens and gram positive eubacteria in the transformation of CT, inhibitors were added to some batch cultures at $t = 0$. Vancomycin (0.14 mM), 2-bromoethane sulfonic acid (Bres; 6 mM) and molybdate (2 mM) were used as inhibitors.

Analytical methods

CT, CF, DCM and chloromethane were quantified by headspace gas chromatography. Liquid samples (100–1000 μ l) were injected in 10 ml headspace autosampler vials with Teflon-lined butyl rubber stoppers and aluminum crimp seals. The final volume was adjusted to 2 ml with demineralized water. The vials were analyzed using a Hewlett Packard 19395A headspace sampler connected to a gas chromatograph equipped with an electron capture detector and a CP-Sil 5CB reactor (de Best et al. 1997). Calibration samples were analyzed according to the same method to adjust for air/water partitioning. A four-point curve was used for calibration.

Carbon dioxide and methane concentrations were determined after separation on a CarboPlot P7 column using a gas chromatograph equipped with FID and a methanizer (de Best et al. 1997). Liquid samples (2 ml) from the packed-bed reactor were injected in 10 ml headspace autosampler vials with Teflon-lined butyl rubber stoppers and aluminum crimp seals and equilibrated at 80 °C for 45 min. An amount of 50 μ l of the headspace was injected into the GC by hand with a 100 μ l Hamilton gas and liquid-tight syringe. For batch cultures, 50 μ l of the headspace was injected into the GC. A four point calibration curve was used for quantification.

Samples from ¹³C-CT experiments were analyzed for peaks of molecular weights of 60 (¹²C-CH₃COOH), 61 and 62 (¹³C-CH₃COOH) with a gas chromatograph equipped with a mass selective detector (Hewlett Packard, Wilmington, USA) in the selected-ion-monitoring (SIM) mode. The presence of ¹³C-CH₃COOH besides ¹²C-CH₃COOH was demonstrated from an increase in the 61/60 amu isotopic intensity ratio (All-ratio). The natural ratio between ¹³C and ¹²C is 1.1%. This means that unlabeled CH₃COOH will give a peak in the mass spectrum at 60 amu (¹²C-CH₃COOH) and a smaller one at 61 amu (¹³C-CH₃COOH). The intensity of the peak at 61 amu

will be about 1.1% of the peak at 60 amu. An increase in this intensity ratio demonstrates the presence of ¹³C-CH₃COOH in a sample. The detection limit for unlabeled ¹²C-CH₃COOH was 1.67 μ M.

Sulfate and chloride were determined after separation on an IONPAC AG9-SC guard column and IONPAC AG9-SC anion column (Dionex, Breda, The Netherlands) (de Best et al. 1997) with an ion chromatograph equipped with a conductivity detector, thermal stabilizer and suppressor (Dionex, Breda, The Netherlands).

Acetate concentrations were determined with an enzymatic test-combination based on the formation of NADH (Boehringer, Mannheim, Germany). NADH formation was measured by the increase in absorbance at 340 nm on a JASCO UV/VIS spectrophotometer (JASCO, The Netherlands). The detection limit was 3 μ M.

Chemicals

All chemicals were obtained from commercial suppliers. Carbon tetrachloride was obtained from Baker. Chloroform (CF) and dichloromethane were purchased from Rathburn. ¹³C-carbon tetrachloride was purchased from C.N. Schmidt B.V. (Amsterdam, The Netherlands) as a 99% pure solution. Vancomycin and sodium molybdate were purchased from Sigma. Chloromethane and 2-bromo-ethanesulfonic acid were obtained from Aldrich. Calibration gases were obtained from AGA (carbon dioxide, methane).

Results

Transformation of CT was studied in a packed-bed reactor under conditions of simultaneous methanogenesis and sulfate reduction to investigate the rate and extent of biological mineralization. Acetate (1 mM) served as a substrate.

First the packed-bed reactor was operated without CT. Acetate was completely utilized by methanogenic and sulfate reducing bacteria. Methane production (0.85 ± 10.05 mM) indicated that 85% of the added acetate was converted by methanogens. Methanogenic bacteria use acetate both as electron acceptor and donor and convert acetate to methane and carbon dioxide. The presence of methanogens was confirmed by fluorescence microscopy (Doddema and Vogels 1978). Sulfate reducing bacteria utilized 15% of the added acetate for the complete reduction of all available sulfate (0.15 ± 0.1 mM).

Four days after CT ($1.3 \mu\text{M}$) was first added to the influent, complete transformation of CT occurred. Most of the CT transformation ($>85\%$) took place within the first 10 cm of the packed-bed reactor. Chloroform ($0.30 \mu\text{M}$) was found as a transformation product at the first sample port. Further along the reactor chloroform (CF) was completely degraded.

Since CT is often found in contaminated groundwater's at higher concentrations, CT transformation in the packed-bed reactor was studied at different concentrations. The CT concentration in the reactor was increased stepwise to $60 \mu\text{M}$ over a period of 15 weeks. Up to a concentration of $40 \mu\text{M}$ CT was completely transformed. Besides CF ($<0.9 \mu\text{M}$), dichloromethane ($<0.8 \mu\text{M}$) was found as a transformation product in the effluent of the reactor. At a CT concentration of $60 \mu\text{M}$, low concentrations of CT ($0.9 \pm 0.2 \mu\text{M}$) were detected in the effluent of the reactor showing that CT was transformed for more than 98%.

Acetate (1 mM) was completely metabolized up to a CT concentration of $40 \mu\text{M}$. Sulfate reducing microorganisms accounted for about 15% of the removal of added acetate as calculated from the amount of sulfate removed, while methanogens converted about 85% to methane according to the amount of methane produced. When the CT concentration was further increased to $60 \mu\text{M}$, acetate utilization decreased until only about 65% of the added acetate was metabolized (Table 1). Sulfate reducing bacteria still utilized 0.15 mM of the added acetate for the complete reduction of sulfate. Methane production by methanogens decreased from 0.85 to about 0.55 mM .

Effect of the acetate concentration on carbon tetrachloride transformation

To determine the acetate concentration that is sufficient for complete transformation of CT, the acetate concentration was decreased stepwise from 1000 to $0 \mu\text{M}$ at an influent CT concentration of about $60 \mu\text{M}$. At each acetate concentration the reactor was operated for at least 60 days.

Down to an acetate concentration of $8 \mu\text{M}$, the amount of CT transformed slowly decreased (Figure 1). This was in great part caused by a decrease in CT mineralization. The amount of CT converted to CF increased. The concentration of dichloromethane (DCM) in the effluent of the reactor was always low ($< 0.6 \mu\text{M}$) and further decreased during the course of the experiment until DCM was no longer found




Figure 1. Transformation of carbon tetrachloride in an anaerobic packed-bed reactor at different acetate concentrations. Symbols: CCl_4 transformed (\square); CCl_4 mineralized (\blacksquare); CHCl_3 formed (\bullet).




Figure 2. Transformation of carbon tetrachloride in an anaerobic packed-bed reactor without addition of an electron donor. Symbols: CCl_4 (\blacksquare); Cl^- (\square), CH_3COOH (\bullet).

as a transformation product. At acetate concentrations below $30 \mu\text{M}$, the percentage of CT mineralized no longer decreased and remained constant at about 73% of the CT added.

Carbon tetrachloride transformation in the absence of acetate

The continuous absence of acetate in the influent of the reactor had no significant effect on CT transformation. After running the packed-bed reactor in the absence of acetate for 115 days (CT was continuously added at a concentration of about $53 \mu\text{M}$) still $37.5 \mu\text{M}$ (72%) of CT was degraded (Figure 2, Table 1), mainly to non-chlorinated products as indicated by the chloride production ($141 \mu\text{M}$). CF was no longer found as a transformation product.

Table 1. Mass balance of an anaerobic carbon tetrachloride degrading packed-bed reactor. CH₃COOH (different concentrations), CCl₄ (56.8 ± 4 μM), SO₄²⁻ (0.15 ± 0.1 mM) and CO₂ (1.1 ± 0.05 mM) were present in the influent of the reactor. CHCl₃, CH₂Cl₂ or CH₄ were not present in the influent of the reactor. The mass balance was calculated from the difference between influent and effluent concentrations

Influent (μM)		Transformation/formation ^e (μM)							
CH ₃ COOH	CCl ₄	CCl ₄ transformed	CHCl ₃ formed	CH ₂ Cl ₂ formed	Cl ⁻ formed	CH ₃ CCOOH transformed	SO ₄ ²⁻ transformed	CH ₄ formed	CO ₂ formed
1030	60.6	59.4	0.3	0.2	- ^a	630	128	570	420
484	56.6	55.3	2.1	0.2	-	319	134	268	224
237	56.3	53.9	3.5	0.1	-	160	105	71	149
132	61.3	58.8	4.3	< ^b	-	101	110	30	115
53	59.1	54.2	5.7	0.5	204	46	46	3	73
33	55.8	49.0	7.2	<	157	15	11	<	-
15	53.5	47.3	6.5	<	161	8	-	<	-
8	59.4	44.1	2.9	<	178	6	<	<	-
0	51.9	37.5	<	<	141	-4 ^d	<	<	-
0 ^c	53.5	37.3	<	<	143	-5	<	<	-

^a not determined; ^b <, below detection limit; ^c no Na₂S (42 μM) added to the influent of the reactor; ^d negative sign means production in the reactor; ^e calculated from difference between influent and effluent concentrations.

Table 2. Transformation of ¹³C-labeled carbon tetrachloride in an anaerobic packed-bed reactor

Sample port	Distance from reactor inlet (cm)	CT (μM)	C (μM)	¹² C-CH ₃ COOH (μM)	¹³ C-CH ₃ COOH (μM)	61/60 All-ratio ^a (%)
1	0 (=influent)	56.2	0	33	< ^b	1.3 ± 0.1
3	20	33.6	ND ^c	28	0.1	1.6 ± 0.1
4	30	18.6	ND	1.7	<	1.6 ± 0.1
5	40 (=effluent)	15.2	157	<	<	- ^d
-	standard ^e	-	33.3	-	1.3 ± 0.1	

^a All-ratio = amu isotopic intensity ratio; ^b <, below detection limit; ^c ND not determined; ^d cannot be determined; ^e standard = 33.3 μM (2.0 mg/l) unlabeled CH₃COOH.

The complete dechlorination of CT in the reactor was not an abiotic (chemical) conversion for two reasons. First, in a sterile control reactor, filled with PUR particles and autoclaved digested sludge (three times for 20 min at 120 °C) no transformation of CT was detected. This indicated that abiotic hydrolysis of CT to CO₂ (Figure 3, pathway 1), as described by Kriegman-King and Reinhard (1992), was not a significant process in our reactor. Secondly, removal of sulfide, a potential chemical reductant of CT (Figure 3, pathway 5) (Curtis and Reinhard 1994, Kriegman-King and Reinhard 1992), from the influent of the reactor had no significant effect on the transformation of CT (Table 1, bottom line).

Mass balance for carbon tetrachloride transformation

For a better understanding of the mechanism of microbial CT transformation in the reactor, mass balances were determined at different acetate concentrations. Table I shows mass balances of the packed-bed reactor at nine different acetate concentrations. The mass balance at each acetate concentration is the average of at least two steady states. A steady state was based on constant performance for 28 days.

With a decrease in acetate concentration, the production of methane proportionally decreased until no methane production occurred. Sulfate was nearly completely reduced down to an acetate concentration of 132 μM. At acetate concentrations below 132 μM, sulfate reduction gradually decreased until sulfate reduction no longer occurred. Since CT transformation




Figure 3. Proposed pathways for the anaerobic transformation of carbon tetrachloride (deduced from Criddle et al. 1991).

still occurred in the absence of methane production and sulfate reduction, these results suggest that neither methanogenic nor sulfate reducing bacteria were involved in CT transformation.

When acetate was no longer added to the influent of the reactor, acetate was still found at different heights in the reactor (Figure 2). A net acetate production of 4 up to 20 μM was also measured at acetate concentrations in the influent of the reactor between 8 and 250 μM (data not shown). Acetate production usually indicates activity by acetogenic bacteria. Besides CT added to the influent, viable and non-viable biological solids present in the reactor are the only possible carbon source which theoretically could serve as a substrate for acetogens. When CT was removed

from the influent of the reactor, acetate production no longer occurred. This indicates that viable and non-viable biological solids probably did not serve as a substrate for acetogens. To demonstrate transformation of CT to CH_3COOH in our reactor we carried out experiments with ^{13}C labeled carbon tetrachloride (^{13}C -CT).

Transformation of ^{13}C -CT

The transformation of ^{13}C -CT was studied at a concentration of unlabeled ^{12}C - CH_3COOH in the influent of the reactor of 30 μM . Addition of unlabeled ^{12}C - CH_3COOH to the influent of the reactor was necessary because the concentration of ^{13}C - CH_3COOH was de-

terminated from the 61/60 amu isotopic intensity ratio (All-ratio) of unlabeled $^{12}\text{C-CH}_3\text{COOH}$ (see material and methods). Unlabeled $^{12}\text{C-CH}_3\text{COOH}$ was added to the influent at a low concentration to prevent interference of the peak of unlabeled $^{12}\text{C-CH}_3\text{COOH}$ with the peak of $^{13}\text{C-CH}_3\text{COOH}$ in the mass spectrum. Besides $^{12}\text{C-CH}_3\text{COOH}$, no other electron donor was added to the influent of the reactor.

When $^{13}\text{C-CT}$ ($60\ \mu\text{M}$) was added, the conversion of CT did not alter. During the 35 days of the experiment, about 80% of the CT added was transformed. CF and DCM were not found as transformation products. The liquid samples taken at the influent (first sample port) and effluent of the reactor and at sample ports at 20 and 30 cm from the inlet of the reactor, were analyzed for $^{13}\text{CH}_3\text{COOH}$ and $^{12}\text{CH}_3\text{COOH}$. The results are shown in Table 2.

The 61/60 amu ratio of the standard of unlabeled $^{12}\text{C-CH}_3\text{COOH}$ ($33.3\ \mu\text{M}$) was $1.3 \pm 0.1\%$. This ratio differs from the theoretical ratio of 1.1%. This is probably caused by an interference of the solvent used for extraction with the intensity of molecular mass 60 and seemed to have no effect on the determination of $^{13}\text{C-CH}_3\text{COOH}$.

At the first sample port of the reactor, $33\ \mu\text{M}$ of unlabeled acetate was found. The All-ratio was similar to the standard, indicating that no additional $^{13}\text{C-CH}_3\text{COOH}$ is present. At sample ports 3 and 4, part of the added acetate was utilized, and an increase in the All-ratio from 1.3 ± 0.1 detected. This increase indicated that $^{13}\text{C-CH}_3\text{COOH}$ was a product of CT transformation in the reactor. Egli et al. (1988) also found transformation of $^{14}\text{C-CT}$ to $^{14}\text{CH}_3\text{COOH}$. $^{14}\text{CH}_3\text{COOH}$ accounted for 38% of the products formed by *Acetobacterium Woodii*.

From the increase in the All-ratio and the unlabeled $^{12}\text{C-CH}_3\text{COOH}$ ($28\ \mu\text{M}$) concentration, a concentration of $^{13}\text{C-CH}_3\text{COOH}$ of $0.1\ \mu\text{M}$ was calculated for sample port 3, resulting from $^{13}\text{C-CT}$ transformation. The $^{12}\text{C-CH}_3\text{COOH}$ concentration of sample port 4 is too low to calculate a reliable $^{13}\text{C-CH}_3\text{COOH}$ concentration. These very low concentrations of $^{13}\text{C-CH}_3\text{COOH}$ in the reactor were probably caused by an immediate utilization of $^{13}\text{C-CH}_3\text{COOH}$ formed. The product of $^{13}\text{C-CH}_3\text{COOH}$ transformation, probably $^{13}\text{C-CO}_2$, could not be detected.

Effect of inhibitors on CT transformation

To determine whether acetogenic bacteria could be involved in the transformation of CT ($6.3\ \mu\text{M}$), batch

experiments with an enrichment culture obtained from the reactor were performed in the presence of vancomycin ($0.14\ \text{mM}$). Vancomycin, an inhibitor of cell wall synthesis in gram positive eubacteria (Distefano et al. 1992) and used to inhibit acetogenic bacteria, partly inhibited transformation of CT for 14 days (Table 3). In the absence of vancomycin, transformation of CT started immediately. These results indicated that gram positive bacteria, possibly acetogens, were in some way involved in the transformation of CT.

At acetate concentrations in the reactor below $53\ \mu\text{M}$, no methane production occurred while about 74% of the CT added to the influent of the reactor was removed (Table 1). This suggested that methanogenic bacteria were not involved in the transformation of CT. This was confirmed by batch experiments in the presence of 2-bromoethane sulfonic acid (Bres; $6\ \text{mM}$). Bres, an inhibitor of methanogenesis (Distefano et al. 1992), only reduced the transformation of CT by about 10%, whereas methane production was completely inhibited (Table 3).

At acetate concentrations in the reactor below $8\ \mu\text{M}$ (Table 1) and batch experiments in the absence of inhibitors (Table 3), no sulfate reduction occurred while CT was still transformed. This suggested that sulfate reducing bacteria were also not involved in the transformation of CT. This was confirmed by batch experiments in the presence of molybdate ($2\ \text{mM}$). Molybdate, a specific inhibitor of sulfate reducing bacteria (Smith and Klug 1981), only reduced the transformation of CT by about 10% while no sulfate reduction occurred.

In sterile control batches there was no transformation of CT and CH_3COOH (data not shown).

Discussion

Here we report complete biological dechlorination of CT ($37.5\ \mu\text{M}$) in an anaerobic packed-bed reactor without addition of an electron, suggesting that CT degradation in the reactor was independent of an added electron donor. It is also likely that CT served as an electron acceptor since CT is a much stronger electron acceptor than carbon dioxide, the only other potential electron acceptor present in the reactor.

Two possible pathways for the microbial transformation of CT can be distinguished where degradation is independent of a (second) electron donor or electron acceptor, as with a fermentation process. Both pathways start with a 2-electron reduc-

Table 3. Effect of inhibitors on transformation of carbon tetrachloride with 100 μM of acetate and with or without and inhibitor (batch experiments)

Inhibitor	CCl_4 degraded (μM)	CHCl_3 formed (μM)	CH_3COOH degraded (μM)	CH_4 formed (μM)	SO_4^{2-} reduced (μM)
None	6.29 (100%)	0.69	37.5	36.5	<
Bres	5.30 (88%)	0.69	33.8	<	<
Vancomycin	2.31 (38%)	< ^a	6.5	4.0	<
Molybdate	5.42 (86%)	0.29	35.5	5.8	<

^a Below detection limit.

Table 4. Gibbs free energy changes (ΔG°) of carbon tetrachloride transformation

Reaction	ΔG° (kJ/reaction) ^a
1. $\text{CCl}_4 + \text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{CO} + 2\text{H}^+ + 4\text{Cl}^-$	-528.3
2. $\text{CCl}_4 + 2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{HCOOH} + 2\text{H}^+ + 4\text{Cl}^-$	-544.8
3. $\text{CO} + \text{H}_2\text{O} \rightarrow \text{CO}_2 + 2\text{H}^+ + 2\text{e}^-$	-91.4
5. $\text{HCOOH} \rightarrow \text{CO}_2 + 2\text{H}^+ + 2\text{e}^-$	-74.9
6. $\text{CO} + 1/2\text{H}_2\text{O} \rightarrow 1/4\text{CH}_3\text{COOH} + 1/2\text{CO}_4$	-39.5
7. $\text{HCOOH} \rightarrow 1/4\text{CH}_3\text{COOH} + 1/2\text{CO}_2 + 1/2\text{H}_2\text{O}$	-23.0
8. $1/4\text{CH}_3\text{COOH} + 1/2\text{H}_2\text{O} \rightarrow 1/2\text{CO}_2 + 2\text{H}^+ + 2\text{e}^-$	-51.9
9. $\text{CCl}_4 + 2\text{H}_2\text{O} \rightarrow \text{CO}_2 + 4\text{H}^+ + 4\text{Cl}^-$	-619.7

^a Calculated on the basis of data from Thauer et al. (1977) and Dolfig and Janssen (1994), assuming the following conditions: H^+ , 10^{-7} M; Cl^- , 10^{-3} M.

tion of CT, which formally leads to the formation of dichlorocarbene (Figure 3, pathway 3 and 4). Dichlorocarbene is further transformed to either CO or HCOOH through substitutive dehalogenation. CO and HCOOH can be further utilized and serve as electron donor and source of carbon resulting in the formation of carbon dioxide (Figure 3, Table 4). Overall, for the complete dechlorination of CT to CO_2 no net electrons, thus no addition of an electron donor or electron acceptor would be needed. Transformation of CT to CO_2 via HCOOH or CO are also energetically very favorable reactions with an overall ΔG° of -619.7 kJ/reaction.

Both biotic transformation of CT to HCOOH and CT transformation to CO have been demonstrated (Hashsham et al. 1995, Dybas et al. 1995, Stromeyer et al. 1992). The mechanism of this conversion is not yet clear, but Egli et al. (1988, 1990) proposed that CT transformation may involve the acetyl-CoA pathway and that cobamides could play a role. There are several reports about the stimulation of CT transformation in the presence of cobamides (Chiu and Reinhard 1996, Hashsham et al. 1995, Stromeyer et al. 1992). It is not clear whether CT was degraded via HCOOH

or CO in our reactor. Both the transformation of CT to CO and the transformation of CT to HCOOH are energetically very favorable reactions (Table 4). The samples of the ^{13}C -CT experiment were screened for ^{12}C - and ^{13}C -HCOOH but neither of these compounds were detected in any of the samples, indicating that, at most, less than $2.2 \mu\text{M}$ was present (detection limit). We were unable to detect for ^{12}CO and ^{13}CO since the mass of ^{12}CO is similar to that of N_2 .

The observed formation of ^{13}C - CH_3COOH from ^{13}C -CT indicated that CH_3COOH could be a (minor) product of CT transformation in the reactor. Formation of CH_3COOH would result from the utilization of CO or HCOOH by acetogens (Ljungdahl 1986) (Table 4). CH_3COOH , when formed, again could be oxidized and serve as an electron donor in the reactor, yielding eight electrons per CH_3COOH oxidized (Table 4). Batch experiments with vancomycin strongly suggested that acetogens could be involved in CT transformation in our reactor.

Only isolation of the CT degrading microorganism can give a definitive answer whether acetogenic bacteria or a new physiological type of microorgan-

ism were involved in CT transformation in our open microbial system.

Acknowledgments

This work was financed by a grant No. IOP91225 from the Dutch IOP Environmental Biotechnology program.

References

- Bae W, Odencrantz J, Rittmann BE & Valocchi A (1990) Transformation kinetics of trace-level halogenated organic contaminants in a biologically active zone (BAZ) induced by nitrate injection. *J. Contamin. Hydrol.* 6: 53–68
- Bouwer EJ & McCarty PL (1983a) Transformations of 1- and 2-carbon halogenated aliphatic organic compounds under methanogenic conditions. *Appl. Environ. Microbiol.* 45: 1286–1294
- Bouwer EJ & McCarty PL (1983b) Transformations of halogenated organic compounds under denitrification conditions. *Appl. Environ. Microbiol.* 45: 1295–1299
- Bouwer EJ & Wright JP (1988) Transformations of trace halogenated aliphatics in anoxic biofilm reactor. *J. Cont. Hydrol.* 2: 155–169
- Chiu P-C & Reinhard M (1996) Transformation of carbon tetrachloride by reduced vitamin B₁₂ in aqueous cysteine solution. *Environ. Sci. Technol.* 30: 1882–1889
- Cobb GD & Bouwer EJ (1991) Effects of electron acceptors on halogenated organic compound biotransformation in a biofilm reactor. *Environ. Sci. Technol.* 25: 1068–1074
- Criddle CS, DeWitt JT, Grbic-Galic D & McCarty PL (1990) Transformation of carbon tetrachloride by *Pseudomonas* sp. strain KC under denitrification conditions. *Appl. Environ. Microbiol.* 56: 3240–3246
- Criddle CS & McCarty PL (1991) Electrolytic model system for reductive dehalogenation in aqueous environments. *Environ. Sci. Technol.* 25: 973–978
- Curtis GP & Reinhard M (1994) Reductive dehalogenation of hexachloroethane, carbon tetrachloride and bromoform by anthrahydroquinone disulfonate and humic acid. *Environ. Sci. Technol.* 28: 2393–2401
- De Best JH, Jongema H, Weijling A, Doddema HJ, Janssen DB & Harder W (1997) Transformation of 1,1,1-trichloroethane in an anaerobic packed-bed reactor at various concentrations of 1,1 1-trichloroethane, acetate and sulfate. *Appl. Microbiol. Biotechnol.* 48: 417–423
- De Best JH, Salminen E, Doddema HJ, Janssen DB & Harder W (1998) Transformation of carbon tetrachloride under sulfate reducing conditions. *Biodegradation* 8: 429–436
- Distefano TD, Gossett JM & Zinder SH (1992) Hydrogen as an electron donor for dechlorination of tetrachloroethene by an anaerobic mixed culture. *Appl. Environ. Microbiol.* 58: 3622–3629
- Doddema HJ & Vogels GD (1978) Improved identification of methanogenic bacteria by fluorescence microscopy. *Appl. Environ. Microbiol.* 36: 752–754
- Dolfing J & Janssen DB (1994) Estimates of Gibbs free energies of formation of chlorinated aliphatic compounds. *Biodegradation* 5: 21–28
- Dybas MJ, Tataru GM & Criddle CS (1995) Localization and characterization of the carbon tetrachloride transformation activity of *Pseudomonas* sp. strain KC. *Appl. Environ. Microbiol.* 61: 758–762
- Egli C, Scholtz R, Cook A & Leisinger T (1987) Anaerobic dechlorination of tetrachloromethane and 1,2-dichloroethane to degradable products by pure cultures of *Desulfobacterium* sp. and *Methanobacterium* sp. *FEMS Microbiol. Lett.* 43: 257–261
- Egli C, Tschan T, Scholtz R, Cook A & Leisinger T (1988) Transformation of tetrachloromethane to dichloromethane and carbon dioxide by *Acetobacterium woodii*. *Appl. Environ. Microbiol.* 54: 2819–2824
- Egli C, Stromeyer S, Cook A & Leisinger T (1990) Transformation of tetrachloromethane and chloroform to CO₂ by anaerobic bacteria is a non-enzymic process. *FEMS Microbiol. Lett.* 68: 207–212
- Gälli R & McCarty PL (1989) Biotransformation of 1,1,1-trichloroethane, trichloromethane, and tetrachloromethane by a *Clostridium* sp. *Appl. Environ. Microbiol.* 55: 837–844
- Hashsham S, Scholze R & Freedman DL (1995) Cobalamin-enhanced anaerobic transformation of carbon tetrachloride. *Environ. Sci. Technol.* 29: 2856–2863
- Holliger C & Schumacher W (1994) Reductive dehalogenation as a respiratory process. *Antonie van Leeuwenhoek* 66: 239–246
- Kriegman-King MR & Reinhard M (1992) Transformation of carbon tetrachloride in the presence of sulfide, biotite and vermiculite. *Environ. Sci. Technol.* 26: 2198–2206
- Ljungdahl LG (1986) The autotrophic pathway of acetate synthesis in acetogenic bacteria. *Ann. Rev. Microbiol.* 40: 415–450
- Mägli A, Wendt M & Leisinger T (1996) Isolation and characterization of *Dehalobacterium formicoaceticum* gen. nov. sp. nov., a strictly anaerobic bacterium utilizing dichloromethane as source of carbon and energy. *Arch. Microbiol.* 166: 101–108
- Mäymo-Gatell X, Chien YT, Gossett JM & Zinder SH (1997) Isolation of a bacterium that reductively dechlorinates tetrachloroethene to ethene. *Science* 276: 1568–1571
- Mohn WW & Tiedje JM (1990) Strain DCB-1 conserves energy for growth from reductive dechlorination coupled to formate oxidation. *Arch. Microbiol.* 153: 267–271
- Petersen JN, Skeen RS, Amos KM & Hooker BS (1994) Biological destruction of tetrachloromethane: I Experimental design and data. *Biotechnol. & Bioengin.* 43: 521–528
- Picardal FW, Arnold RG & Huey BB (1995) Effects of electron donor and acceptor conditions on reductive dehalogenation of tetrachloromethane by *Shewanella putrefaciens* 200. *Appl. Environ. Microbiol.* 61: 8–12
- Smith RL & Klug MJ (1981) Electron donors utilized by sulfate-reducing bacteria in eutrophic lake sediments. *Appl. Environ. Microbiol.* 42: 116–121
- Stromeyer SA, Stumpf K, Cook AM & Leisinger T (1992) Anaerobic transformation of tetrachloromethane by *Acetobacterium woodii*: separation of dechlorinative activities in cell extracts and roles for vitamin B₁₂ and other factors. *Biodegradation* 3: 113–123
- Thauer RK, Jungermann K & Decker K (1977) Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol. Rev.* 41: 100–180
- Trautnecker J, Preuß A & Diekert G (1991) Isolation and characterization of a methyl chloride utilizing, strictly anaerobic bacterium. *Arch. Microbiol.* 156: 416–421
- Vogel TM, Criddle CS & McCarty PL (1987) Transformations of halogenated aliphatic compounds. *Environ. Sci. Technol.* 21: 722–736