

University of Groningen

1.4-GHz polarimetric observations of the two fields imaged by the DASI experiment

Bernardi, G.; Carretti, E.; Sault, R. J.; Cortiglioni, S.; Poppi, S.

Published in:
Monthly Notices of the Royal Astronomical Society

DOI:
[10.1111/j.1365-2966.2006.10649.x](https://doi.org/10.1111/j.1365-2966.2006.10649.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bernardi, G., Carretti, E., Sault, R. J., Cortiglioni, S., & Poppi, S. (2006). 1.4-GHz polarimetric observations of the two fields imaged by the DASI experiment. *Monthly Notices of the Royal Astronomical Society*, 370(4), 2064-2070. <https://doi.org/10.1111/j.1365-2966.2006.10649.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

1.4-GHz polarimetric observations of the two fields imaged by the DASI experiment

G. Bernardi,^{1,2*} E. Carretti,^{1†} R. J. Sault,³ S. Cortiglioni¹ and S. Poppi⁴

¹INAF–IASF Bologna, Via Gobetti 101, Bologna I-40129, Italy

²Kapteyn Astronomical Institute, University of Groningen, PO Box 800, 9700 AV Groningen, The Netherlands

³CSIRO–ATNF, PO Box 76, Epping, NSW 1710, Australia

⁴INAF–IRA Bologna, Via Gobetti 101, Bologna I-40129, Italy

Accepted 2006 May 31. Received 2006 April 28; in original form 2006 March 6

ABSTRACT

We present results of polarization observations at 1.4 GHz of the two fields imaged by the DASI experiment ($\alpha = 23^{\text{h}}30^{\text{m}}$, $\delta = -55^\circ$ and $\alpha = 00^{\text{h}}30^{\text{m}}$, $\delta = -55^\circ$, respectively). Data were taken with the Australia Telescope Compact Array with 3.4-arcmin resolution and $\sim 0.18\text{-mJy beam}^{-1}$ sensitivity. The emission is dominated by point sources, and we do not find evidence for diffuse synchrotron radiation even after source subtraction. This allows to estimate an upper limit of the diffuse polarized emission. The extrapolation to 30 GHz suggests that the synchrotron radiation is lower than the polarized signal measured by the DASI experiment by at least two orders of magnitude. This further supports the conclusions drawn by the DASI team itself about the negligible Galactic foreground contamination in their data set, improving by a factor of ~ 5 the upper limit estimated by Leitch et al.

The dominant point-source emission allows us to estimate the contamination of the cosmic microwave background (CMB) by extragalactic foregrounds. We computed the power spectrum of their contribution, and its extrapolation to 30 GHz provides a framework where the CMB signal should dominate. However, our results do not match the conclusions of the DASI team about the negligibility of point-source contamination, suggesting taking into account a source subtraction from the DASI data.

Key words: polarization – radiation mechanisms: non-thermal – cosmic microwave background – diffuse radiation – radio continuum: ISM.

1 INTRODUCTION

The key role played by the cosmic microwave background polarization (CMBP) in modern cosmology is now well established. The cosmic microwave background (CMB) *E* mode carries information on the reionization (e.g. Zaldarriaga, Spergel & Seljak 1997), and joined to the temperature anisotropies, helps constraining the cosmological parameter estimation (e.g. Zaldarriaga et al. 1997). The CMB *B* mode looks interesting for its connection with the inflationary era (e.g. Kosowsky 1999) and its capability to measure the amount of primordial gravitational waves.

On the experimental hand, the search for the CMBP is underway. Detections have been claimed by the DASI (Leitch et al. 2005), CAPMAP (Barkats et al. 2004), CBI (Readhead et al. 2004) and *BOOMERanG* (Montroy et al. 2005) teams. However, we are far

from a complete characterization of the *E*-mode power spectrum whereas the *B*-mode detection is still missing.

The tiny level of the CMB polarized component makes the contamination by foreground astrophysical sources even more severe than for the anisotropy temperature term.

At frequencies lower than 100 GHz, the foreground contribution is expected to be dominated by both the Galactic synchrotron emission and the extragalactic radiosources.

Several theoretical works (Mesa et al. 2002; Tucci et al. 2005; de Zotti et al. 2005) studied the contamination due to point-source emission on the microwave background using data coming from low-frequencies surveys. In particular, Mesa et al. (2002) and Tucci et al. (2005) achieve similar results pointing out that the contamination to the CMB *E* mode is not serious in the 70–100 GHz frequency range. However, both of them find that extragalactic sources could be a considerable contaminant up to 44 GHz. In addition, observational efforts are underway in order to establish the properties of the radiosources directly at high frequency. The results from 18-GHz observations of the Kuhr sample (Ricci et al. 2004a) and preliminary results from an 18-GHz survey by Ricci et al. (2004b) show a

*E-mail: bernardi@iasfbo.inaf.it (GB)

†Present address: INAF–IRA Bologna, Via Gobetti 101, Bologna I-40129, Italy.

substantial agreement with the predictions by Mesa et al. (2002). Given this, the estimate of the E -mode contamination by point sources is still open for frequencies up to 44 GHz.

About the Galactic synchrotron emission, template maps (e.g. Giardino et al. 2002; Bernardi et al. 2004) suggest that the CMBP E -mode signal should be higher than the synchrotron emission on large angular scales and for frequencies around 90 GHz.

On degree and subdegree scales, polarization observations of the diffuse synchrotron radiation in selected low-emission regions have recently begun with the aim of estimating the possible contamination on the CMBP. The field imaged by the *BOOMERanG* experiment was observed with both the Australia Telescope Compact Array (ATCA) at 1.4 GHz (Bernardi et al. 2003, hereafter B03; Carretti et al. 2005a) and the Parkes telescope at 2.3 GHz (Carretti et al. 2005b) providing the first characterization of the diffuse radiation in a low-emission area. They find that the Galactic signal should not prevent the detection of the CMB E mode in that area at frequencies ≥ 30 GHz. A similar analysis has been conducted in another area in the Northern sky providing similar conclusions (Carretti et al. 2006). Although these results can be used as indicators of conditions in low-emission regions, every area has its own specific features and requires dedicated observations, especially if representing a target for CMBP experiments.

Among the sky regions imaged by CMB experiments, direct observations of the synchrotron emission only exist for *BOOMERanG* area (B03) to date. Barkats et al. (2004) assessed that the CAPMAP field is not contaminated at 90 GHz since they checked that the total intensity synchrotron emission in their area is below the detected polarized signal. On the other hand, DASI and CBI operated at ~ 30 GHz, where the synchrotron emission can still play a relevant role. In particular, the DASI team itself pointed out that no polarization observations are available for the two sky patches they observed (Kovac et al. 2002).

In this paper, we present results of the first deep polarization observations of the two fields imaged by the DASI experiment at a frequency of 1.4 GHz, where the synchrotron emission is dominant.

This paper is organized as follows. In Section 2, we describe the observations while in Section 3, we present the power spectrum analysis. Finally, in Section 4 we discuss the results in the framework of CMBP measurements.

2 OBSERVATIONS AND RESULTS

Observations of the two DASI fields (hereafter named as field 1 and field 2, see Table 1 for their coordinates) were performed in 2003 July and 2004 September with the ATCA. The EW214 configuration was used to sample the spatial frequencies between 3 and 30 arcmin. Each field has a $3^\circ \times 3^\circ$, size and the same amount of time (~ 72 h) was spent in each of them. A sensitivity of ~ 0.18 mJy beam $^{-1}$ has been achieved in both fields in the inner $2^\circ \times 2^\circ$ square, correspond-

ing to ~ 3.2 mK. Both of them were observed with the same system configuration whose characteristics are reported in Table 1. A standard procedure for the ATCA data reduction, as described in B03, was followed to obtain Stokes I , Q , U and V images.

Figs 1 and 2 show the results for the fields 1 and 2, respectively. It is evident that the cleaning procedure did not manage to completely remove the side lobes for the brightest sources even in polarization.

The emission is dominated by point sources both in total intensity and in polarization. Besides this, no evident diffuse polarized emission rises up the noise level in both fields. This is in contrast with that found in the B03 area observed with the same array (same configuration and sensitivity). In order to make this contrast evident, we plot the map of the Stokes parameter U measured in the B03 field for a direct comparison in Figs 1 and 2 (similar results hold for the Q images). All the structures present in the DASI fields seem to be associated to point-like sources and no extended structures are evident like in the B03 field, where a polarized diffuse emission of ~ 11 mK was detected spread throughout the area. Besides intrinsic differences due to different sky positions, the lower emission found there has been explained by the action of Faraday effects. These, by transferring power from large to small angular scales, enhance the polarized emission on the 3–30 arcmin scales to which the observations are sensitive (Carretti et al. 2005b).

The DASI fields, however, are at higher Galactic latitudes ($b \sim -58^\circ$ and -62° for the two areas, respectively) and Carretti et al. (2005a) found that the emission is unlikely to be affected by Faraday rotation effects at 1.4 GHz at $|b| \geq 50^\circ$. This seems to be confirmed also by the recent polarization map at 1.4 GHz of the Northern celestial hemisphere (Wolleben et al. 2005).

We identify the point sources present in the two fields by setting a detection threshold of 10σ in order to avoid false and spurious candidates due to the presence of residual non-white noise. The analysis provides a catalogue complete down to a polarized flux $I_p = 2$ mJy (see Tables 2 and 3).

We look for counterparts and find that every source, but the number six of field 1, has a radio counterpart in the Parkes survey at 408 MHz (PKS; Wright 1994), in the Parkes–MIT–NRAO survey at 4.85 GHz (PMN; Griffith & Wright 1993) or in the Sydney University Molonglo Sky Survey at 843 MHz (SUMSS; Mauch et al. 2003). In the case of source number six, we find a counterpart in the optical/infrared Automated Plate Measuring survey (APM; Maddox et al. 1990). When a source is present in more than one survey, the strongest one within the beam area is listed.

Actually, we cannot justify the absence of the source number six in the field 1 in the SUMSS survey. In fact, if we consider a power-law behaviour for the flux density $F_\nu \propto \nu^{\alpha_\nu}$ as a function of the frequency ν and assume a spectral index $\alpha_\nu = -0.7$, this source would have a flux of ~ 195 mJy at 843 MHz whereas the sensitivity of the SUMSS is 8 mJy. Even in the case of a source fully self-absorbed, the spectral index would be $\alpha_\nu = 2.5$ and the flux at 843 MHz would be ~ 40 mJy, at the limit of a 5σ detection in the SUMSS. It is worth noting that the source is unlikely to be in a complete self-absorbed regime because of its polarized emission; then the spectral slope should be flatter.

Given the limited sample, no general conclusion can be drawn but just a few qualitative remarks. We note that there are almost the same number of sources in the two fields (18 and 15, respectively) at the flux limit $I_p = 2$ mJy and almost all the sources show a polarization degree of a few per cent whereas only three sources are more than 10 per cent polarized. The mean polarization percentage is 6.2 per cent for the first field and 3.8 per cent for the second one.

Table 1. Main characteristics of the observations.

Central frequency	1380 MHz
Effective bandwidth	205 MHz
Field 1 (J2000)	$\alpha = 23^{\text{h}}30^{\text{m}}$, $\delta = -55^\circ$
Field 2 (J2000)	$\alpha = 00^{\text{h}}30^{\text{m}}$, $\delta = -55^\circ$
Area size for each field	$2^\circ \times 2^\circ$
Sensitivity (flux)	0.18 mJy beam $^{-1}$
Sensitivity (temperature)	3.2 mK
Gain	17.3 K Jy $^{-1}$ beam $^{-1}$




Figure 1. DASI field 1: total intensity I (top left hand panel), polarized intensity $I_p = \sqrt{Q^2 + U^2}$ (top right hand panel). The U image (bottom right hand panel) is also reported for comparison with the diffuse emission (point sources subtracted) detected in B03 (bottom left hand panel).

3 POWER SPECTRUM ANALYSIS

We compute the power spectra of the E and B modes in the inner $2^\circ \times 2^\circ$ square of both fields, where the sensitivity is higher. We use the Fourier transform-based method by Seljak (1997) already applied in previous works (e.g. Tucci et al. 2002; Carretti et al. 2005a). It can be applied as dealing with small-sky areas where the flat geometry approximation can be assumed. E - and B -mode spectra are computed from the Fourier transforms of the Q and U images as follows (Seljak 1997):

$$C_{E\ell} = \left\{ \frac{\Omega}{N_\ell} \sum_{\mathbf{l}} |Q(\mathbf{l}) \cos(2\phi_l) + U(\mathbf{l}) \sin(2\phi_l)|^2 - w_p^{-1} \right\} b^{-2}(\ell) \quad (1)$$

$$C_{B\ell} = \left\{ \frac{\Omega}{N_\ell} \sum_{\mathbf{l}} |-Q(\mathbf{l}) \sin(2\phi_l) + U(\mathbf{l}) \cos(2\phi_l)|^2 - w_p^{-1} \right\} b^{-2}(\ell), \quad (2)$$

where Ω is the sky fraction, N_ℓ is the number of modes around the multipole ℓ , ϕ_l is the angle associated to the vector \mathbf{l} , w_p^{-1} is the pixel noise, $b^{-2}(\ell)$ is the power spectrum of the beam window function [see Seljak (1997) for a more detailed discussion].

The calculated power spectra are plotted in Fig. 3 together with the noise contribution defined by (Tegmark 1997)

$$C_\ell^{\text{noise}} = \frac{\Omega w_p^{-1}}{B_\ell^2}. \quad (3)$$

They fit the power-law equation

$$C_\ell^X = C_{2000}^X \left(\frac{\ell}{2000} \right)^{\beta_X}, \quad X = E, B \quad (4)$$

as a function of the multipole ℓ . Best-fitting parameters are shown in Table 4. The slopes of the spectra for the field 2 are compatible with a flat $\beta = 0$ spectrum within 2σ CL, which is typical of the emission dominated by point sources. A similar result holds for the E mode of the field 1, whereas the B mode looks steeper and compatible with a point-source power spectrum only at $\sim 3\sigma$ CL. Also the spectra, therefore, support an emission dominated by point sources.

To estimate an upper limit of the diffuse emission contribution, we subtracted all the bright sources listed in Tables 2 and 3. The resulting power spectra are shown in Fig. 4.

A remarkable lowering of the flux and a certain steepening of the slopes can be seen, in particular, for field 2. This is clearly shown by the best-fitting parameters quoted in Table 5.

Although these spectra certainly exhibit a lower contribution by point sources, they cannot be considered as representative of the diffuse emission. In fact, the spectra match the expected noise level at the high ℓ -range tail, so that the power on the largest scales can likely be due to residual non-white noise. As a consequence, we prefer to consider these spectra as upper limits of the diffuse emission.




Figure 2. DASI field 2: total intensity I (top left hand panel), polarized intensity I_p (top right hand panel). The U image (bottom panel) is also reported.

Table 2. Position, total intensity (I) and polarized intensity (I_p) of the sources detected in the first DASI field. The rms error is the same for both the two intensities and corresponds to the beam-sensitivity ($0.18 \text{ mJy beam}^{-1}$) but a 5 per cent error due to accuracy has to be added. Polarization angle (ϕ) and polarization degree ($\Pi = I_p/I$) are also reported. The last column provides the distance from the counterpart.

Source	RA J2000	Dec. J2000	I (mJy)	I_p (mJy)	ϕ	Π (per cent)	Counterpart	Distance (arcmin)
1	23 ^h 22 ^m 07 ^s .1	−54°45′29″.1	1914.00	50.06	−33°9 ± 0°:1	2.6	PKS 2319-55	0.0
2	23 ^h 33 ^m 07 ^s .4	−54°16′12″.0	296.10	20.76	−38°6 ± 0°:3	7.0	PKS 2330-545	0.1
3	23 ^h 29 ^m 34 ^s .5	−54°25′25″.2	125.90	6.99	39°0 ± 0°:7	5.6	SUMSS J232942−542524	1.2
4	23 ^h 28 ^m 04 ^s .8	−55°41′11″.2	99.84	6.73	11°9 ± 0°:8	2.4	SUMSS J232806−554110	0.2
5	23 ^h 28 ^m 35 ^s .4	−54°41′15″.0	55.91	6.30	−27°6 ± 0°:8	11.27	SUMSS J232835−544125	0.2
6	23 ^h 27 ^m 27 ^s .2	−54°42′31″.3	136.70	5.50	−2°4 ± 0°:9	4.0	APMUKS(BJ) B232445.84-545734.8	1.7
7	23 ^h 33 ^m 29 ^s .3	−54°09′25″.4	57.30	4.76	−10°2 ± 1°:1	8.3	SUMSS J233329−540934	0.1
8	23 ^h 27 ^m 24 ^s .8	−54°51′01″.0	105.60	4.45	−26°5 ± 1°:2	4.2	SUMSS J232724−545114	0.2
9	23 ^h 29 ^m 25 ^s .4	−54°54′22″.5	209.00	4.31	−8°7 ± 1°:2	2.1	SUMSS J232925−545434	0.2
10	23 ^h 32 ^m 07 ^s .6	−54°44′06″.1	63.00	4.27	35°0 ± 1°:2	6.8	SUMSS J233207−544406	0.1
11	23 ^h 26 ^m 14 ^s .4	−54°03′07″.7	57.60	3.70	−22°3 ± 1°:4	6.4	SUMSS J232614−540321	0.2
12	23 ^h 30 ^m 33 ^s .1	−54°32′06″.4	24.02	3.63	14°3 ± 1°:4	15.0	SUMSS J233033−543141	0.4
13	23 ^h 34 ^m 44 ^s .3	−54°19′00″.7	43.37	3.32	34°2 ± 1°:6	7.7	SUMSS J233445−541910	0.2
14	23 ^h 27 ^m 20 ^s .3	−55°09′16″.9	104.20	2.86	−31°6 ± 1°:8	2.7	SUMSS J232718−550936	0.5
15	23 ^h 31 ^m 20 ^s .1	−55°27′06″.4	12.84	2.67	−26°3 ± 1°:9	20.8	SUMSS J233119−552702	0.1
16	23 ^h 32 ^m 53 ^s .9	−55°11′16″.2	106.10	2.31	−34°5 ± 2°:2	2.2	SUMSS J233253−551055	0.4
17	23 ^h 25 ^m 20 ^s .1	−55°02′17″.5	148.20	2.21	6°5 ± 2°:3	1.5	SUMSS J232520−550228	0.2
18	23 ^h 34 ^m 05 ^s .1	−54°11′00″.6	131.20	2.13	25°3 ± 2°:4	1.6	SUMSS J233404−541058	0.0

Table 3. As for Table 2 but for field 2.

Source	RA J2000	Dec. J2000	I (mJy)	I_p (mJy)	ϕ	Π (per cent)	Counterpart	Distance (arcmin)
1	00 ^h 31 ^m 02 ^s .8	−55°21′12″.5	264.10	15.66	26°0 ± 0°3	5.9	PKS 0028-556	0.1
2	00 ^h 30 ^m 33 ^s .6	−54°38′01″.2	608.50	15.54	−9°3 ± 0°3	2.6	PKS 0028-549	0.2
3	00 ^h 32 ^m 26 ^s .3	−54°54′55″.3	493.00	11.71	6°1 ± 0°4	2.4	PKS 0030-552	0.2
4	00 ^h 36 ^m 54 ^s .0	−55°36′28″.9	196.90	11.25	15°3 ± 0°5	5.7	SUMSS J003653−553632	0.1
5	00 ^h 26 ^m 14 ^s .7	−55°23′38″.4	105.50	8.51	−21°3 ± 0°6	8.1	SUMSS J002615−552337	0.1
6	00 ^h 35 ^m 57 ^s .3	−54°03′23″.0	177.40	8.27	−4°7 ± 0°6	4.7	SUMSS J003558−540324	0.1
7	00 ^h 32 ^m 55 ^s .3	−54°56′28″.0	1381.00	7.09	7°6 ± 0°7	0.5	SUMSS J003254−545614	0.3
8	00 ^h 33 ^m 20 ^s .3	−55°37′54″.9	134.10	6.61	−39°6 ± 0°8	4.9	SUMSS J003319−553757	0.1
9	00 ^h 23 ^m 36 ^s .9	−54°31′13″.2	98.59	6.26	−15°8 ± 0°8	6.3	SUMSS J002340−543151	0.9
10	00 ^h 22 ^m 40 ^s .9	−55°03′58″.4	120.60	4.51	−23°3 ± 1°1	3.7	SUMSS J002241−550400	0.0
11	00 ^h 35 ^m 53 ^s .1	−54°16′47″.1	146.50	4.31	28°7 ± 1°2	2.9	SUMSS J003554−541634	0.3
12	00 ^h 25 ^m 10 ^s .7	−54°27′39″.9	131.40	3.56	−39°1 ± 1°5	2.7	PMN J0025−5427	0.0
13	00 ^h 28 ^m 15 ^s .2	−55°26′43″.6	138.70	3.36	−40°8 ± 1°5	2.4	SUMSS J002814−552701	0.3
14	00 ^h 23 ^m 03 ^s .4	−55°42′48″.3	159.80	3.32	8°4 ± 1°6	2.1	PMN J0023−5542	0.3
15	00 ^h 33 ^m 44 ^s .8	−54°56′24″.8	152.70	3.23	−37°1 ± 1°6	2.1	SUMSS J003344−545619	0.1



Figure 3. Power spectra of field 1 (left-hand panel) and field 2 (right-hand panel) with the respective 1σ error bars. Solid line represents the noise (see text).

Table 4. Fit parameters for E - and B -mode spectra plotted in Fig. 3. The fit has been performed in the $800 < \ell < 2800$ multipole range.

Spectrum	Number of field	C_{2000}^X (μK^2)	β_X
C_ℓ^E	1	183 ± 9	-0.19 ± 0.15
C_ℓ^B	1	177 ± 9	-0.44 ± 0.13
C_ℓ^E	2	345 ± 17	-0.20 ± 0.10
C_ℓ^B	2	340 ± 16	$+0.34 \pm 0.16$

4 DISCUSSION

The spectra reported in Fig. 3 allow an estimate of the contamination to the CMB due to point-source emission. Then, we use the E -mode power spectrum (Fig. 3) of the field 2 that, having the highest signal, represents the worst case. The frequency extrapolation is not trivial because of the large range of possible spectral slopes. In low-frequency surveys, the radio population is a mixture of steep ($\alpha_\nu < -0.5$) and flat-spectrum ($\alpha_\nu > -0.5$) sources. In particular, the steep-spectrum sources have been found for the 87 per cent of the population in the NRAO VLA Sky Survey (NVSS; Condon et al. 1998), whereas the remaining 13 per cent are flat-spectrum sources. Therefore, we take the 13 per cent of the E -mode power spectrum and scale it up to 30 GHz with a brightness temperature spectral in-

dex $\alpha = -2.25$, which is an average value in the interval $-2.5 < \alpha < -2.0$ quoted by Toffolatti et al. (1998) for flat-spectrum sources. Then, we take the other 87 per cent of the E -mode power spectrum and scale it up with $\alpha = -2.75$, as reported by Peacock & Gull (1981) for steep-spectrum sources. The two resulting contributions are added together to give the estimate of the E -mode power spectrum at 30 GHz. Since the angular behaviour is compatible with a point-source power spectrum, we extend it down to $\ell = 200$. The result is shown in Fig. 5 together with the DASI measurements and the CMBP E -mode power spectrum expected according to cosmological parameters measured after *Wilkinson Microwave Anisotropy Probe* (WMAP) data (Spergel et al. 2003).

The situation represented in Fig. 5 does not exclude a point-source contamination to the CMB E -mode power spectrum. At $\ell = 300$, we find $\ell(\ell + 1)C_\ell^E/2\pi \simeq 1.3 \mu\text{K}^2$, which is just slightly greater than the value quoted as upper limit by Leitch et al. (2005, $0.98 \mu\text{K}^2$ at 2σ CL). At higher multipoles, the estimated point-source emission is increasing and, compared to the CMB peaks, ranges from a factor of 10 to a factor of 3 lower than the CMB spectrum (3.2 and 1.7 in signal, respectively). Although within the limits of the uncertainties in the frequency extrapolation, our results suggest that the point-source contamination in the DASI data is not negligible. In the light of this, a re-analysis of the data considering a subtraction of the detected sources could allow a better estimate of the extragalactic signal component.




Figure 4. As for Fig. 3 but after point-sources subtraction.

Table 5. As for Table 4 but after point-sources subtraction.

Spectrum	Number of field	C^X_{2000} (μK^2)	β_X
C^E_{ℓ}	1	73 ± 4	-1.29 ± 0.10
C^B_{ℓ}	1	80 ± 4	-0.63 ± 0.13
C^E_{ℓ}	2	32 ± 2	-1.09 ± 0.20
C^B_{ℓ}	2	36 ± 3	-0.58 ± 0.25




Figure 5. The CMB E -mode power spectrum (solid line) computed with the *WMAP* first year cosmological parameters (Spergel et al. 2003), the measurements reported by the DASI team (asterisks; Leitch et al. 2005), the extrapolation to 30 GHz of the measured power spectra in the second DASI field (solid straight line) and its extension down to $\sim 1^\circ$ angular scale (dashed line).

Power spectra of Fig. 4 allow us to estimate an upper limit on the diffuse synchrotron emission as contaminant of the CMBP. We use the result of the field 1, which represents the worst case. The power spectrum of the field 1 is scaled up to 30 GHz using a brightness temperature spectral index $\alpha = -3.1$ (Bernardi et al. 2004), and the result is shown in Fig. 6. Since the multipole range accessible by our and DASI data is only marginally overlapped, we perform an extrapolation in the ℓ -space. Since the spectrum we measure is an upper limit and prevents us to use its slope for such extrapolation, we consider a pessimistic and a more realistic case.

We use a slope $\beta = -2.7$ that is the steepest slope measured so far for the synchrotron E mode (Tucci et al. 2002), as a worst case. This value measured on the Galactic plane at 1.4 GHz is likely to be significantly altered by Faraday effects. However, at the latitudes




Figure 6. As Fig. 5 but for the diffuse synchrotron emission: the extrapolation to 30 GHz of the upper limit on the synchrotron power spectrum obtained by our 1.4-GHz observations (solid straight line) is plotted with two possible extrapolations of the synchrotron power spectrum up to $\sim 1^\circ$ angular scale with a slope $\beta = -2.7$ (dashed line) and a slope $\beta = -1.6$ (dot-dashed line, see text for details). The arrows indicate that our extrapolation represents an upper limit.

of the DASI fields, a value $\beta = -1.6 \pm 0.2$ is more likely expected (Bruscoli et al. 2002), which is more typical where Faraday rotation effects are weak. Therefore, we perform a second extrapolation using $\beta = -1.6$.

The results (Fig. 6) show that the expected theoretical CMBP E -mode power spectrum is not contaminated down to the peak at $\ell \sim 300$. Considering the most likely slope of $\beta = -1.6$, we see that the theoretical peak at $\ell \sim 120$ would not be contaminated by the synchrotron emission. The average value of the synchrotron power spectrum over the $200 < \ell < 1050$ multipole range is $\sim 0.19 \mu\text{K}^2$, which is a factor of ~ 5 lower than the upper limit quoted by the DASI team ($0.91 \mu\text{K}^2$; see Leitch et al. 2005). This further supports the evaluation of negligible synchrotron contamination done by the DASI team itself.

In the light of the data presented in this paper, a twofold picture is emerging. Even if we did not detect the synchrotron diffuse emission at 1.4 GHz, the upper limit we set is robust enough to conclude that this component is not a serious contaminant of the CMB E mode at 30 GHz. For this reason, our observations support the conclusions inferred by Kovac et al. (2002) and Leitch et al. (2005).

On the other hand, we argue that the problem regarding the point-source contamination is still open. Leitch et al. (2005) find an upper

limit of $0.98 \mu\text{K}^2$ and exclude a contamination by point sources. From the data presented here, we retrieve a value very close to it but we cannot exclude that the CMBP power spectrum is not contaminated at $\ell > 300$ (Fig. 5). Our data suggest the CMBP signal should dominate over the point-source emission at 30 GHz but a reappraisal of their contamination is recommended.

ACKNOWLEDGMENTS

We thank Marijke Haverkorn for useful discussions and an anonymous referee for suggestions which improved this paper. This work has been carried out in the framework of the SPORt experiment, a programme funded by Italian Space Agency (ASI). The ATCA is part of the Australia Telescope, which is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. We acknowledge the use of the CMBFAST package. This research has made use of the NASA/IPAC Extragalactic Database which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Barkats D. et al., 2004, *ApJ*, 619, L127
 Bernardi G., Carretti E., Cortiglioni S., Sault R. J., Kesteven M. J., Poppi S., 2003, *ApJ*, 594, L5 (B03)
 Bernardi G., Carretti E., Fabbri R., Sbarra C., Cortiglioni S., Poppi S., Jonas J. L., 2004, *MNRAS*, 351, 436
 Bruscoli M., Tucci M., Natale V., Caretti E., Fabbri R., Sbarra C., 2002, *MNRAS*, 330, 43
 Carretti E., Bernardi G., Sault R. J., Cortiglioni S., Poppi S., 2005a, *MNRAS*, 358, 1
 Carretti E., McConnell D., McClure-Griffiths N. M., Bernardi G., Cortiglioni S., Poppi S., 2005b, *MNRAS*, 360, L10
 Carretti E., Poppi S., Reich W., Reich P., Fürst E., Bernardi G., Cortiglioni S., Sbarra C., 2006, *MNRAS*, 367, 132
 Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693
 de Zotti G., Ricci R., Mesa D., Silva L., Mazzotta P., Toffolatti L., González-Nuevo J., 2005, *A&A*, 431, 893
 Giardino G., Banday A. J., Górski K. M., Bennett K., Jonas J. L., Tauber J., 2002, *A&A*, 387, 82
 Griffith M., Wright A., 1993, *A&A*, 105, 1666
 Kosowsky A., 1999, *New Astron. Rev.*, 43, 157
 Kovac J. M., Leitch E. M., Pryke C., Carlstrom J. E., Halverson N. W., Holzzapfel W. L., 2002, *Nat*, 420, 772
 Leitch E. M., Kovac J. M., Halverson N. W., Carlstrom J. E., Pryke C., Smith M. W. E., 2005, *ApJ*, 624, 10
 Maddox S. J., Sutherland W. J., Efstathiou G., Loveday J., 1990, *MNRAS*, 243, 692
 Mauch T., Murphy T., Buttery H. J., Curran J., Hunstead R. W., Piestrzynski B., Robertson J. G., Sadler E. M., 2003, *MNRAS*, 342, 1117
 Mesa D., Baccigalupi C., De Zotti G., Gregorini L., Mack K.-H., Vigotti M., Klein U., 2002, *A&A*, 396, 463
 Montroy T. E. et al., 2005, preprint (astro-ph/0507514)
 Peacock J. A., Gull S. F., 1981, *MNRAS*, 196, 611
 Readhead A. C. S. et al., 2004, *Sci*, 306, 836
 Ricci R., Prandoni I., Gruppioni C., Sault R. J., De Zotti G., 2004a, *A&A*, 415, 549
 Ricci R. et al., 2004b, *MNRAS*, 354, 305
 Seljak U., 1997, *ApJ*, 482, 6
 Spergel D. N. et al., 2003, *ApJ*, 148, 175
 Tegmark M., 1997, *Phys. Rev. D.*, 56, 4514
 Toffolatti L., Argueso Gomez F., de Zotti G., Mazzei P., Franceschini A., Danese L., Burigana C., 1998, *MNRAS*, 297, 117
 Tucci M., Carretti E., Cecchini S., Fabbri R., Orsini M., Pierpaoli E., 2000, *New Astron.*, 5, 181
 Tucci M., Carretti E., Cecchini S., Nicastro L., Fabbri R., Gaensler B. M., Dickey J. M., McClure-Griffiths N. M., 2002, *ApJ*, 579, 607
 Tucci M., Martínez-González E., Vielva P., Delabrouille J., 2005, *MNRAS*, 360, 935
 Wolleben M., Landecker T. L., Reich W., Wielebinski R., 2005, *A&A*, 448, 411
 Wright A. E., 1994, *AuJPh*, 47, 585
 Zaldarriaga M., 1997, *PRD*, 55, 1822
 Zaldarriaga M., Spergel D. N., Seljak U., 1997, *ApJ*, 488, 1

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.