

University of Groningen

## On the three-dimensional structure of edge-on disc galaxies

Pohlen, Michael; Zaroubi, Saleem; Peletier, Reynier F.; Dettmar, Ralf-Juergen

*Published in:*  
Monthly Notices of the Royal Astronomical Society

*DOI:*  
[10.1111/j.1365-2966.2007.11790.x](https://doi.org/10.1111/j.1365-2966.2007.11790.x)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2007

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Pohlen, M., Zaroubi, S., Peletier, R. F., & Dettmar, R-J. (2007). On the three-dimensional structure of edge-on disc galaxies. *Monthly Notices of the Royal Astronomical Society*, 378(2), 594-616.  
<https://doi.org/10.1111/j.1365-2966.2007.11790.x>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# On the three-dimensional structure of edge-on disc galaxies

Michael Pohlen,<sup>1,2\*</sup> Saleem Zaroubi,<sup>1</sup> Reynier F. Peletier<sup>1</sup> and Ralf-Jürgen Dettmar<sup>2</sup>

<sup>1</sup>*Kapteyn Astronomical Institute, University of Groningen, PO Box 800, NL-9700 AV Groningen, the Netherlands*

<sup>2</sup>*Astronomisches Institut, Ruhr-Universität Bochum, Universitätsstrasse 150, D-44780 Bochum, Germany*

Accepted 2007 March 28. Received 2007 March 28; in original form 2006 September 25

## ABSTRACT

A simple algorithm is employed to deproject the two-dimensional images of a pilot sample of 12 high-quality images of edge-on disc galaxies and to study their intrinsic three-dimensional (3D) stellar distribution. We examine the radial profiles of the stars as a function of height above the plane and report a general trend within our sample of an increasing radial scalelength with height outside of the dust lane. This could be explained by the widespread presence of a thick disc component in these galaxies. In addition, the 3D view allows the study of the vertical distribution of the outer disc, beyond the break region, where we detect a significant increase in scalelength with vertical distance from the major axis for the truncated discs. This could be regarded as a weakening of the ‘truncation’ with increasing distance from the plane. Furthermore, we conclude that the recently revised classification of the radial surface brightness profiles found for face-on galaxies is indeed independent of geometry. In particular, we find at least one example of each of the three main profile classes as defined in complete samples of intermediate to face-on galaxies: not-truncated, truncated and antitruncated. The position and surface brightness that mark the break location in the radial light distribution are found to be consistent with those of face-on galaxies.

**Key words:** galaxies: evolution – galaxies: formation – galaxies: fundamental parameters – galaxies: photometry – galaxies: structure.

## 1 INTRODUCTION

To study galaxy formation and evolution it is often helpful to imagine that galaxies are made up of a small number of separate components. Surface photometry of galaxies provides a very simple way of parametrizing these various components. By modelling the surface-brightness distribution of a galaxy and thereby parametrizing the individual components, we obtain a common ground to measure, compare and sort large samples of galaxies.

Owing to their flat geometry the random orientation of a specific galaxy disc provides us with different projected views of the, presumably, common underlying three-dimensional (3D) distribution. In particular, one gets complementary views from observing face-on or edge-on galaxies. In the face-on view, one studies the radial distribution of the starlight discussing, e.g. non-axisymmetric features such as bars, rings or spiral arms. In the edge-on view on the other hand, one is able to study the vertical distribution, e.g. disentangle thick disc, thin disc and stellar haloes. Because of a longer line-of-sight projection, the edge-on view allows in principle the exploration of the radial profile to larger distances.

Early on, from face-on galaxy observations, several authors (e.g. Patterson 1940; de Vaucouleurs 1959) noticed that the radial light distribution of stellar discs is well described by a simple exponential, albeit the lack of a clear physical explanation for this fact. Later on, Freeman (1970) called these discs Type I – as opposed to some other galaxies called Type II – showing a break in their radial profile with a downbending steeper outer region, or equivalently, a flatter inner region.

Using the edge-on view, van der Kruit (1979) discovered that the exponential decline does not continue to the last measured point, but is *truncated* after several radial scalelengths. Afterwards, with the advent of CCD technology, several groups (Barteldrees & Dettmar 1994; Pohlen et al. 2000b; de Grijs, Kregel & Wesson 2001; Florido et al. 2001; Pohlen 2001; Kregel, van der Kruit & de Grijs 2002) studied larger samples of edge-on galaxies to find out if and at what radius this truncation occurs.

More recently, Erwin, Beckman & Pohlen (2005), Pohlen & Trujillo (2006) and Erwin et al. (in preparation) have analysed surface brightness profiles of face-on to intermediate inclined galaxies and revised our view on disc truncations. They identified three basic classes of surface brightness profiles depending on the apparent break features or lack of them. (1) The well-known Type I that has an exponential profile, with no break. (2) Type II with a ‘downbending break’. (3) A completely new class, called Type III, also

\*E-mail: pohlen@astro.rug.nl

described by a broken exponential but with an upbending profile beyond the break. In this view the *classical truncations* described initially in the pioneering work of van der Kruit (1979) and van der Kruit & Searle (1981) are in fact part of a generalized Freeman Type II class (Freeman 1970). In this picture truncated galaxies show a surface brightness profile with a shallow inner and a steeper outer exponential region separated at a relatively well-defined break radius.

These results are in agreement with those reported by Pohlen et al. (2002a), who obtained deep photometry of three face-on galaxies and clearly demonstrated that the sharp cut-offs<sup>1</sup> fitted to profiles of edge-on galaxies (e.g. by van der Kruit & Searle 1981; Barteldrees & Dettmar 1994; Pohlen et al. 2000b) are in fact not complete but better described by a broken exponential. Even before, de Grijs et al. (2001) and Pohlen (2001) noted that for edge-on galaxies the surface brightness distribution does not disappear asymptotically ('vertically') into the background noise. Interestingly, Hunter & Elmegreen (2006) find a similar broken exponential shape for many objects in their large sample of irregular galaxies.

The redefinition of the truncation as a point at which the two exponential fits meet solves, among other things, the apparent contradiction between simulations by Holley-Bockelmann & Mihos (2001) and the edge-on observations of Barteldrees & Dettmar (1994) and Pohlen, Dettmar & Lütticke (2000a). The latter authors find that 'truncations' are as 'close' as twice the radial scalelength from the centre in contradiction with the simulations. Furthermore, 'early' truncations (i.e. at  $R \lesssim 2h$ ) of edge-on galaxies have been always questioned by the 'face-on viewers', who had never observed a face-on galaxy with a sharp truncation at a distance of twice the scalelength.

Despite the line-of-sight enhancement of the surface brightness level in edge-on galaxies, star counts in nearby galaxies are able to provide significantly deeper profiles with much higher signal-to-noise ratio (S/N). Using this completely independent method, Ferguson et al. (2006) followed the profile of the face-on galaxy M33 down to  $\mu_{\text{lim}} \sim 30 I\text{-mag/arcsec}^2$ , which is currently unreachable with surface photometry. Their result shows that the broken exponential description is valid down to the last bin above the noise level and an actual *edge* of the disc is not found, at least for this galaxy. Likewise, Bland-Hawthorn et al. (2005) have made comparably deep star counts for the Scd galaxy NGC 300 and also do not find an outer edge. Its surface brightness profile continues as an unbroken single exponential (Type I) to the limit of their star counts.

Nevertheless, there still seem to be some open issues for a complete reconciliation of the renewed face-on with the 'old' edge-on view (de Zeeuw 2007) as to provide a consistent picture of the 3D structure, examples are the following.

(i) Is the broken exponential structure, now well observed for face-on galaxies, quantitatively consistent with the truncated profiles observed for edge-on galaxies?

(ii) Do we see a different kind of truncation in the deeper views provided by the edge-on galaxies?

(iii) Can we identify the edge-on counterparts of the three main types of surface brightness profiles (Type I, Type II and Type III) observed for face-on galaxies by Pohlen & Trujillo (2006) and Erwin et al. (in preparation)?

(iv) Does the frequency of different profile classes found for the face-on galaxies match the edge-on results?

(v) Why did the former edge-on surveys, with large and deep samples, not discover the new class of Type III profiles?

To answer these questions and disentangle the crucial problems with the line-of-sight integration (see e.g. Pohlen et al. 2004a) we decided to deproject a sample of edge-on galaxies. This can only be done using some geometric assumption. Here we choose the assumption of axial symmetry. This will allow us to gain insight into the 3D galactic stellar disc structure by bringing together the classical face-on and edge-on views.

The vertical structure may well be the key to understand the nature of the different breaks in the surface brightness profiles, by setting additional constraints on the various physical explanations put forward. For example, some recent star formation models (Elmegreen & Hunter 2006; Li, Mac Low & Klessen 2006) are now able to produce the observed broken exponential structure, but, alternatively, Debattista et al. (2006) also find downbending breaks using purely collisionless  $N$ -body simulations. Once the nature of the galactic stellar disc is determined it will be fully justified to utilize these breaks for comparison of galaxies at various redshifts as done by Pérez (2004), Trujillo & Pohlen (2005) or Tamm & Tenjes (2006), and explain their appearance in cosmological simulations as shown by Governato et al. (2007).

The remainder of this paper is organized as follows. Section 2 describes our sample selection and data. In Section 3 we introduce our method to deproject the edge-on galaxies. The results are given in Section 4, we discuss them in Section 5 and conclude in Section 6. In Appendix A we give detailed comments for all our galaxies and show in Appendix B radial surface brightness profiles and isophote maps for all galaxies.

## 2 DATA

All optical images used in this paper were taken as part of a PhD study on the radial structure of galactic stellar discs by Pohlen (2001). Originally, the disc galaxies ( $-2 \leq T \leq 7$ ) were chosen according to the allocated observing time, the observatory (north/south), and the available field of view (FOV) ( $D_{25} \gtrsim 2$  arcsec) to meet the following morphological selection criteria verified by using images from the Digitized Sky Survey (DSS). (1) The galaxies are edge-on with  $i \gtrsim 86^\circ$ . (2) Undisturbed with smooth photometric features (see below). (3) They look similar to some prototypical cases like NGC 4565 or IC 2531, this is done to make it possible to consistently fit a simple disc model.

Galaxies with the following characteristics were rejected. (1) Apparent spiral arms, indicating a lower inclination. (2) A significantly asymmetric or disturbed disc, indicating strong interaction. (3) Two-sided or significantly one-sided warped discs. (4) Galaxies apparently dominated by the light of their bulge component (i.e. there are almost no Sa/Sab galaxies, but several lenticulars). (5) Galaxies that showed only a faint, patchy, not well-defined disc. For more details of the original sample we refer the reader to Pohlen (2001) or Pohlen et al. (2004b).

The present sample of 11 galaxies is drawn from this base sample of 72 galaxies as a pilot sample covering a wide range of Hubble types and apparent vertical distributions (thin and thick) while trying to pick galaxies with very few disturbing fore- or background objects. We add an additional near-infrared (NIR) image of one of the galaxies to observationally assess the influence of the dust. Global properties of the finally chosen galaxies are given in Table 1.

<sup>1</sup> It is interesting to note that in this concept the truncation radius is often equal to the last measured point and therefore much further out compared to the break radius of a broken exponential fit.

**Table 1.** The 11 galaxies of our pilot survey.

Galaxy	RA	Dec.	RC3	$T$	Diam.	$v_{\odot}$	$v_{\text{vir}}$	$D$	$v_{\text{rot}}$	$M_B^0$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
NGC 522	01 24 45.9	+09 59 40	.S..4*/	3.7	2.6	2727	2729	37.9	168	-20.6
NGC 1596	04 27 38.1	-55 01 40	.LA..*/	-2.0	4.1	1508	1227	17.0	98	-19.3
NGC 3390	10 48 04.5	-31 32 02	.S..3./	3.0	3.6	2850	2696	37.4	210	-21.4
NGC 4179	12 12 52.6	+01 17 47	.L.../	-2.0	3.9	1239	1269	17.6	-	-19.6
ESO 380-019	12 22 02.1	-35 47 32	.S..6*/	6.0	3.6	2937	2793	38.8	240	-21.2
NGC 5290	13 45 19.1	+41 42 45	.S..4*/	3.7	4.0	2588	2817	39.1	219	-20.8
NGC 5981	15 37 53.4	+59 23 29	.S..5\$/	5.0	2.5	2528	2813	39.1	251	-20.6
UGC 10459	16 35 07.9	+40 59 29	.S..6*.	6.0	1.5	9001	9252	128.5	218	-21.5
IC 4871	19 35 42.4	-57 31 06	.SXS7?/	6.7	3.1	1927	1726	24.0	115	-19.6
FGC 2339	21 44 39.5	-06 41 17	.S..6*/	6.0	1.8	3097	3096	43.0	85	-19.5
ESO 404-018	22 01 10.2	-32 34 44	.7?/	7.0	2.6	2275	2144	29.8	73	-19.1

Notes. (1) Principal name, (2) right ascension, (3) declination, (4) RC3 coded Hubble type and the (5) Hubble parameter  $T$  (all taken from de Vaucouleurs et al. 1991), (6) diameter in arcmin, (7) heliocentric radial velocities, (10) maximum deprojected rotation velocity of the gas, and the  $B$ -band absolute magnitude (11) [taken from Lyon/Meudon Extragalactic Database (LEDA)]. According to the heliocentric radial velocities corrected for the Local Group infall into the Virgo cluster (8) from LEDA, we estimated the (9) distances following the Hubble relation with the Hubble constant from the *Hubble Space Telescope* (*HST*) Key Project of  $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$  (Freedman et al. 2001).

The images were obtained in six different observing runs between 1998 and 2004. Two runs (E1 and E2) at the Danish 1.54-m telescope and one (E3) at the 3.6-m New Technology Telescope (NTT) telescope, both at the European Southern Observatory in Chile. One run (C1) at the 1.23-m telescope and one (C2) at the 3.5-m telescope both at Calar Alto in Spain. Finally, one run (L1) at the Hall 42-inch (1.06-m) telescope of the Lowell Observatory at Anderson Mesa, USA.

During the two runs with the 1.54-m Danish telescope it was equipped with the Danish Faint Object Spectrograph and Camera (DFOSC) and the C1W7/CCD which is a  $2\text{k} \times 2\text{k}$  LORAL chip providing a field size of  $\approx 13$  arcmin and a scale of  $0.39 \text{ arcsec pixel}^{-1}$ . At the NTT we used ESO Multi-Mode Instrument (EMMI), equipped with a  $2\text{k} \times 2\text{k}$  Tektronix chip providing a field size of  $\approx 9$  arcmin and a scale of  $0.27 \text{ arcsec pixel}^{-1}$ . The run at the Calar Alto 1.23-m telescope was done in service mode with the Site#18b chip, a  $2\text{k} \times 2\text{k}$  SITE CCD with  $24 \mu\text{m}$  pixel size, providing an unvignetted field of  $\approx 10$  arcmin and a scale of  $0.5 \text{ arcsec pixel}^{-1}$ . At the 3.5-m telescope we used the wide-field near-infrared camera OmegaPrime with a  $1\text{k} \times 1\text{k}$  pixel HAWAII

HgCdTe array by Rockwell, providing a field of  $\approx 7$  arcmin at a scale of  $0.4 \text{ arcsec pixel}^{-1}$ . During the run at Anderson Mesa the 42-inch telescope was equipped with the NSF CCD used with  $2 \times 2$  binning providing a field size of  $\approx 4.9$  arcmin and a scale of  $\approx 0.73 \text{ arcsec pixel}^{-1}$ . For the optical imaging the available standard Bessel  $R$  and  $V$  filter at ESO, and Johnson  $R$  at Calar Alto and Anderson Mesa are used. The NIR image with OmegaPrime is in the  $K'$  band. Table 2 summarizes the detailed observational parameters for each image.

For more details about the data reduction (i.e. flat-fielding, mosaicing and sky subtraction) and photometric calibration we refer to Pohlen (2001) and Pohlen et al. (2004b). Details of the NIR data reduction can be found in van den Berg et al. (in preparation).


## 2.1 Preparation of the images

Before deprojecting our galaxy images we first have to prepare the images, here we describe the procedure followed in this work (see Fig. 1).

**Table 2.** Observing log for the individually combined images.

Galaxy	Filter	Date	Site	Exp. time	Seeing	Co-adds
(1)	(2)	(3)	(4)	(5)	(6)	(7)
NGC 522	$R$	1998 November	L1	60.0	3.0	$6 \times 600$
NGC 1596	$R$	1998 January	E1	45.0	1.3	$8 \times 300, 2 \times 150$
NGC 3390	$R$	1998 January	E1	51.7	1.6	$3 \times 600, 3 \times 300, 2 \times 200$
NGC 4179	$V$	1999 May	E2	65.0	1.7	$1 \times 600, 1 \times 480, 1 \times 360, 1 \times 300, 9 \times 240$
ESO 380-019	$V$	1999 May	E2	60.0	1.6	$6 \times 600$
NGC 5290	$R$	1999 June	C1	60.0	2.0	$3 \times 600$
NGC 5290	$K'$	2004 February	C2	38.0	1.2	$38 \times 60$
NGC 5981	$R$	1999 June	C1	60.0	1.7	$6 \times 600$
UGC 10459	$R$	1999 June	C1	60.0	1.6	$6 \times 600$
IC 4871	$V$	1999 May	E2	60.0	1.4	$6 \times 600$
FGC 2339	$R$	2000 July	E3	30.0	0.9	$3 \times 600$
ESO 404-018	$V$	1999 May	E2	60.0	1.3	$6 \times 600$

Notes. (1) Galaxy name, (2) filter, (3) observing date, (4) site, (5) the total co-added on-source exposure time, (6) seeing, as measured on the final combined image, (7) the number of individual images with their individual exposure times.



**Figure 1.** Preparation of the images. Contour maps of NGC 5981 showing the four main steps in image preparation. The upper left-hand panel shows the rotated, original image. The upper right-hand panel shows the image after masking of fore- and background sources and interpolating across the masks. The lower left-hand panel shows the galaxy after averaging the four quadrants. The lower right-hand panel shows the galaxy after Gaussian smoothing. The equidistant contour levels plotted are in all cases the same, except for the smoothed version where two additional outer contours are added.

As a first step we mask all the foreground stars and the background galaxies. To do this semi-automatically, we use SExtractor (Bertin & Arnouts 1996) to create an object catalogue. Depending on the measured ellipticity and the provided star/galaxy classifier we separated stars from galaxies, which in turn get assigned circular masks with radii depending on their measured flux. The output is a DS9 region file which is overlaid on the galaxy image and visually inspected for each of the masked objects. For highly elliptical background galaxies the mask is changed to an ellipse. In addition, we have to adapt the radii of the masks. Typically one opts for a conservative masking (i.e. prefer to mask more than necessary and leave ‘larger holes’) to make sure that all non-galaxy features are removed.

However, since our deprojection method cannot deal with masked areas (actual holes) we have to interpolate the flux across our mask. By doing this, one is forced to choose the mask radius as small as possible in order to not introduce too much unwanted structure. This approach clearly leaves residuals on the masked, interpolated image which will be discussed later (cf. Section 3.7).

In a second step the galaxy image is rotated to align the galactic plane parallel to the major axis using the smallest angle of rotation according to their true position on the sky (chip). The rotation and centring is done by blinking the adjacent sides and minimizing the difference by eye. Depending on the intrinsic symmetry this can be done down to an accuracy of  $\pm 0.5^\circ$ .

The third step is to cut the galaxy into four quadrants and average them. This procedure increases the S/N and minimizes the residuals from the interpolated, masked regions and gives a more regular averaged galaxy. The influence of the dust lane and intrinsic asymmetries will be discussed later (cf. Section 3.7).

The final step is smoothing the image. Since our deprojection code only handles one radial cut at a time, we choose the kernel size

for the Gaussian smoothing to match the vertical increment between successive deprojections in order to get fully independent cuts but using all available rows.

### 3 METHOD

In this section we describe and test the method used to recover the 3D structure of the available disc galaxies. We start with describing the deprojection algorithm (Section 3.1) and discuss the noise properties of the reconstructed 3D distribution, especially after the original image has been smoothed with a Gaussian kernel (Section 3.2). To further reduce the noise we applied an additional denoising (Section 3.3). In Section 3.4 we discuss in detail our choice of fitting function for the radial surface brightness distribution. Then we describe how the radial profiles of the recovered distribution for a given  $z$  are actually fitted (Section 3.5). In Section 3.6 we apply the method and the fitting procedure on a large sample of artificial galaxy images in order to test the method’s ability to recover the underlying 3D distribution. In this set of artificial images the influence of the inclination of the galaxy and the existence of dust on the recovery of the 3D distribution are tested. Finally, we estimate the effects of our image preparation on the resulting profiles using real galaxies in Section 3.7.

#### 3.1 Deprojection

The deprojection of axially symmetric quantities is a classical problem in astronomy (e.g. Lucy 1974). It is considered for determining the 3D stellar orbit structure in elliptical galaxies (e.g. Binney, Davies & Illingworth 1990; Dehnen & Gerhard 1993, 1994), and in determining cluster 3D structure from X-ray observations (Fabricant, Rybicki & Gorenstein 1984) and more generally from

X-ray, Sunyaev–Zeldovich and cluster weak-lensing data (Zaroubi et al. 1998, 2001). In the case of edge-on galaxies, Florido et al. (2001, 2006) have applied an inversion method to obtain deprojected profiles before, but without studying a dependence with the vertical distance from the plane. Here we apply a simple analytical method that employs the so-called Fourier slice theorem in order to recover the intrinsic 3D stellar distribution from our sample of high-quality two-dimensional (2D) edge-on disc galaxy images.

The disc galaxy images are deprojected as follows. We adopt the convention that boldface italic symbols denote 3D quantities [e.g.  $\mathbf{k} = (k_x, k_y, k_z)$ ]. Let the observer’s coordinate system be defined with the Cartesian axes  $(x, y, z)$ , with the  $y$ -axis aligned with the line-of-sight. The disc galaxy axis of symmetry lies along the  $z$ -axis which is perpendicular to the line-of-sight. Let  $I(x, z) = \int \lambda(x) dy$  denote a projected quantity (image) of the source function  $\lambda$ . The 3D Fourier transform of the source function is related to the image by

$$\begin{aligned} \tilde{\lambda}(k_x, 0, k_z) &= \int e^{-i(k_x x + k_z z)} I(x, z) dx dz \\ &= \tilde{I}(k_x, k_z). \end{aligned} \quad (1)$$

Note that functions marked with  $\tilde{\phantom{x}}$  are in Fourier space.

This relation could be easily understood since the galaxy has a rotational symmetry, i.e.  $\lambda(\mathbf{r}) = \lambda(r, z)$  and its Fourier transform is  $\tilde{\lambda}(\mathbf{k}) = \tilde{\lambda}(k, k_z)$ , where  $k = \sqrt{k_x^2 + k_y^2}$ . The deprojection is then obtained by inverse Fourier transform of  $\tilde{I}(k_x, k_z)$  and replacing  $k_x$  with  $k$ , namely by using the following equation:

$$\lambda(r, z) = \frac{1}{(2\pi)^2} \int e^{ik_z z} \tilde{I}(k, k_z) J_0(kr) k dk dk_z. \quad (2)$$

Here  $J_0$  is the zero order Bessel function.

Notice that for a specific radial slice, i.e. a cut at a given  $z_0$  and parallel to the major axis, the integral (2) can be written simply as


$$\lambda(r, z_0) = \frac{1}{2\pi} \int \hat{\mathcal{I}}(k, z_0) J_0(kr) k dk, \quad (3)$$

where  $\hat{\mathcal{I}}(k, z_0)$  is the Fourier transform over the image  $x$  direction. This way of writing the equation enables us to easily deproject certain radial slices of the image, or collapsing the whole  $z$  direction of the image to recreate the equivalent of a face-on view of the galaxy.

In principle, it is easy to obtain an equation similar to (2) for the general case of a disc with an arbitrary inclination angle. However, one can show that for such cases there is a cone in  $k$  space inside which there is no information on the 3D structure in the image. The general case is beyond the scope of this paper and is not needed for the range of inclinations we restrict our selves to i.e. to  $i \gtrsim 86^\circ$  (see Section 3.6). The interested reader is referred to Zaroubi et al. (1998, 2001).

Furthermore, one can show that the Fourier slice theorem approach for inverting edge-on galaxies is actually another way of writing the inverse Abel transform (Binney & Tremaine 1987). The advantage of the Fourier space language here is that it avoids taking direct derivatives of noisy data, hence, making the inversion more stable.

The applied fast Fourier transform (FFT) assumes periodic boundary conditions which has to be satisfied by our images. Therefore, one has to pad the image with at least twice the number of the image pixels along each axis. However, since FFT of sharp transitions will give a strong aliasing signal in the deprojected slice, we do not apply zero padding. Rather we padded the image with white noise background with the same rms as the noise in the observed



**Figure 2.** Test of deprojection method for NGC 5981. In the upper frame the radial profiles of the symmetrized image (dashed lines) and those of the deprojected–reprojected image (solid lines) are shown. The uppermost profile is the vertically integrated profile, while each of the others represents a cut at increasing vertical distance from the major axis. The lower frame shows the percentage error one obtains from this procedure.

image. The residual small differences at the edge changing from the artificial white noise to the actual noise in the background of the image are washed out by the final wavelet denoising and Gaussian smoothing of the image (see Sections 3.2 and 3.3). Further tests with more careful tapering of the image’s edge with a Fermi–Dirac function without adding noise showed that there is no significant influence on the deprojected profiles.

As a zero-order test, we deproject the image of NGC 5981 and then reproject it back. The result is that we obtain profiles that are identical to those of the original image (cf. Fig. 2) down to  $<0.5$  per cent error level. Most of this error actually comes from the reprojection – not the deprojection – procedure. The reason is simply that the deprojection is a differential quantity that depends on the intensity of the image at a given radius, whereas the reprojection is an integration procedure that will depend on the intensity of galaxy at every radius including radii beyond those explored in the image (i.e. after the noise dominates over the signal). A more thorough testing of the method in which we compare the deprojected 3D structure with the original one will be presented in Section 3.6.

### 3.2 Noise properties and smoothing

To explore the noise properties, let us suppose that the image only contains white noise, with rms  $\sigma$ , i.e.  $\langle I(x_1, z_1) I(x_2, z_2) \rangle = (2\pi)^2 \delta(x_1 - x_2) \delta(z_1 - z_2) \sigma^2$ . The noise correlation function in the deprojected image is given by

$$\langle \lambda(r_1, z_1) \lambda(r_2, z_2) \rangle = \sigma^2 \frac{\delta(z_1 - z_2)}{(2\pi)} \int k^2 J_0(kr_1) J_0(kr_2) dk. \quad (4)$$

Not surprisingly, the high-frequency components dominate and hence some smoothing must be employed. The analytic nature of the correlation matrix is an attractive feature of the method.

Another nice feature of the method is that any isotropic smoothing function applied to the image will depend only on the modulus of the radial wavevector,  $|k|$ , in the deprojection. This can easily be seen from the previous equations. A simple and practical Gaussian smoothing filter, with smoothing scale  $R_s$ , gives an autocorrelation of the noise of the form

$$\begin{aligned} \mathcal{N}(r) &\equiv \langle \lambda(r, z)^2 \rangle \\ &= {}_2F_2\left(\frac{1}{2}, \frac{3}{2}; 1, 1; -\frac{r^2}{R_s^2}\right) / (8\sqrt{\pi}R_s^5), \end{aligned} \quad (5)$$

where  ${}_2F_2$  is a hypergeometric function. This has a particularly simple asymptotic form

$$\lim_{r \rightarrow \infty} \mathcal{N}(r) \propto r^{-1}. \quad (6)$$

Since we are interested in the large-scale features of the reconstruction a Gaussian smoothing is applied on the image. The choice of the full width at half-maximum (FWHM) of the smoothing kernel is discussed in Section 2.1.

Despite these simple analytic features of the noise properties, we sample the noise properties of the smoothed 3D structure created by the method with deprojecting noisy slices of the image (with a given  $z$  well outside the galaxy). For each image we deproject some 10 randomly selected lines, average their rms and use that as our noise estimate. The profile of the reconstructed noise rms is shown in every set of reconstructed profiles (see Fig. 5).

The noise is dominated not by pixel-to-pixel variations, but by any residual large-scale asymmetries in the background as well as residuals from the masking (cf. Section 2.1). The latter cause the somewhat bumpy structure of the resulting deprojected profiles (see Fig. 5).

The seeing acts in effect as an additional Gaussian smoothing of the image. This type of smoothing flattens the brightness profile around the centre but has very little influence on the scalelength measurements.

### 3.3 Wavelet denoising

In order to reduce the apparent bumpy structures along the radial profiles we apply an extradenosing algorithm based on wavelet expansion. The employment of this algorithm is not necessary and we tested that it does not change the quantitative results of the deprojection in any significant way, it however reduces the apparent ‘bumpiness’ in the radial profiles.

The denoising scheme adopted here follows the method proposed by Donoho (1992) (see also Donoho & Johnstone 1994; Zaroubi & Goelman 2000) where the wavelet coefficients are reduced with a constant equal to the rms of the white noise. This procedure is known as the soft thresholding of wavelet coefficients. The wavelet functions used here are the Daubechies’ discrete wavelets (Daubechies 1988) with 12 parameters (DAUB12).

### 3.4 Profile shape and fitting function

The recent studies of profile shapes for face-on disc galaxies (Erwin et al. 2005; Pohlen & Trujillo 2006; Erwin et al., in preparation) and irregular galaxies (Hunter & Elmegreen 2006) suggest that the *broken exponential* is a good fitting function for the radial light distribution (cf. Section 1). Technically, this results in a discontinuity between the inner and outer slope, which is unphysical and of course not implied. Pohlen & Trujillo (2006) find an extended *transition or break region*, which is not as sharp as measured before (e.g. by

Pohlen et al. 2002a). From their sample only a few clean and undisturbed truncated cases (e.g. NGC 5300 or 7437) exhibit a very sharp transition between the inner and outer disc region. Most often their Type II breaks appear with a more gradual transition zone of a size up to  $\sim 4$  kpc. Despite the fact that the exponential function in itself has no clear physical explanation, the broken exponential is a reasonable fitting function that describes the shape of the profiles in a simple and uniform way.

The key point of this description is that the *outer discs* of truncated galaxies – regardless of any transition zone and the exact shape of the outer profile – makes up a significant fraction of the physical size of the stellar discs. For example, the ratio between the inner and the outer disc is typically 2 to 1 in radius (Pohlen & Trujillo 2006), therefore, in terms of area the outer disc is even slightly larger. In some cases authors ignore the outer region altogether (e.g. van der Kruit & Searle 1981; Pohlen et al. 2000b) and use an exponential with a sharp *cut-off* as their fitting function of choice. However, this is not the case for edge-on galaxies in general as clearly apparent from the examination of the radial profiles of e.g. NGC 5907 in van der Kruit & Searle (1981) or IC 4393 in Pohlen et al. (2000b). We see the same for several galaxies used in our study. For example, the major axis profiles of ESO 380–019, ESO 404–018, NGC 5981, UGC 10459 and NGC 522 (see Fig. B1 in Appendix B) are obviously better described by a broken exponential fit (Pohlen 2001; Pohlen et al. 2002b). Many further examples in the literature (e.g. see Jensen & Thuan 1982; Sasaki 1987; Näslund & Jörsäter 1997; Byun 1998; de Grijs et al. 2001; Pohlen 2001) show that the broken exponential shape is indeed a very good fitting function even for edge-on galaxies. However, we should mention that some authors (e.g. Florido et al. 2001, 2006) still prefer the description of a smooth, but complete truncation.

To summarize, since the profiles of face-on galaxies are well parametrized by a broken exponential model, and those of edge-on galaxies seem to be, it is useful to apply the same parametrization for the profiles of the edge-on galaxies studied here.

### 3.5 Parameter fitting

In order to quantify the behaviour of the recovered 3D stellar distribution of the galaxies, we fit the deprojected radial profiles, for a given  $z$ , with a simple broken exponential function of the form

$$\mu(R) = \begin{cases} \mu_{0,\text{in}} + 1.086 R/h_{\text{in}} & \text{if } R \leq R_{\text{br}}, \\ \mu_{0,\text{out}} + 1.086 R/h_{\text{out}} & \text{if } R > R_{\text{br}}. \end{cases}$$

In principle, the model includes five free parameters, the inner and outer scalelengths,  $h_{\text{in}}$  and  $h_{\text{out}}$ , the inner and outer central surface brightnesses,  $\mu_{0,\text{in}}$  and  $\mu_{0,\text{out}}$ , and the location of the break radius,  $R_{\text{br}}$ , which marks the boundary between the inner and outer regions. However, by using the downhill simplex method (Nelder & Mead 1965) with initial starting values of only the scalelength and central brightness parameters we avoid using the break radius as a free parameter. This is achieved by deciding a priori that the inner slope is either shallower or steeper than the outer one. This enables the break radius to be determined by the crossing point of two exponential fits found by the algorithm. We find that in all cases, except NGC 1596, the inner profile is shallower than the outer profile. In addition, when the ratio between the two fitted radial scalelengths is about  $1 \pm 0.2$ , we fit it with a single exponential profile of the form  $\mu(R) = \mu_0 + 1.086 R/h$ .

In order to avoid including the bulge and/or bar regions in the fitting procedure, i.e. avoid a detailed bulge to disc decomposition, we



used the ‘marking the disc’ method with fixed inner,  $b_1$ , and outer,  $b_2$ , boundaries (see e.g. Pohlen & Trujillo 2006). The inner boundary is chosen to exclude the region that is obviously dominated by the bulge plus bar (shoulder) component. The outer boundary is set at the point at which the surface brightness profile is comparable to the noise level (see Section 3.2). In the case of an extended bump clearly visible in the profile, caused for example by a residual from the masking, the boundary is moved further inside. Sometimes a sharp rise or drop appears beyond the outer boundary of the recovered profiles (see, e.g. the face-on-equivalent profiles of IC 4871 or NGC 5290 *R* band in Fig. 5). This is caused by the FFT operation used in the method. The FFT procedure requires periodic boundary conditions. To avoid contamination of the profile at hand by these conditions, we increase the size of the image by a certain factor. However, this periodicity still shows up in the noise-dominated regions and sometimes close enough to, but never within, the region of interest.

The five starting values for the downhill simplex minimization are obtained by a single exponential fit to the inner or outer region with an approximate break radius for each galaxy.

### 3.6 Tests with artificial data

#### 3.6.1 Model grid

To study the influence of the dust and possible non-90° inclination on the scalelength measurements in our deprojected profiles, we used a grid of 96 mock images (400 × 400 pixel) kindly provided by Simone Bianchi. Using a Monte Carlo radiative transfer method (Bianchi, Ferrara & Giovanardi 1996), he calculated single disc models at a range of edge-on inclinations (80°–90°) with different input parameters for the stellar and dust distribution. A similar approach is adopted by Möllenhoff, Popescu & Tuffs (2006) to measure the influence of dust and inclination for less inclined galaxies.

The chosen inclination angles are 80°, 85°, 86°, 87.5, 89° and 90°. The stellar disc is a 3D single disc model with exponential profiles both along the radial and vertical direction. The radial scalelength is  $h = 33.4$  pixel, and the vertical scaleheight is  $z_0 = 0.125 h$ . The only parameter of the stellar distribution that is varied is the position of an outer (sharp) truncation of the light distribution. The truncation parameter has four values,  $10h$ ,  $5.4h$ ,  $3.4h$  and  $1.4h$ . The first value mimics an infinite exponential profile, and the other three values represent a long truncated, an intermediate truncated and a short truncated stellar disc.

For the dust distribution a set of four different cases with increasing influence on the measured *R*-band light distribution is adopted. The parameters for these cases are within the range measured by Xilouris et al. (1999). The first case is a dust free (transparent) model which we use as our reference case. The second case, which we call the ‘best case’, has the following dust parameters:  $\tau_R = 0.20$ ,  $h_d/h = 1.0$  and  $z_d/z_0 = 0.30$ . Here  $\tau_R$  is the *R*-band optical depth,  $h_d$  is the dust scalelength,  $z_d$  is the dust vertical scaleheight and  $z_0$  is the stellar scaleheight. The parameters of the third case, called the ‘intermediate case’, are  $\tau_R = 0.54$ ,  $h_d/h_* = 1.4$  and  $z_d/z_* = 0.55$ . Finally, the parameters of the fourth case, which we call the ‘worst case’ are  $\tau_R = 0.65$ ,  $h_d/h_* = 1.5$  and  $z_d/z_* = 0.8$ . For the sake of simplicity the dust disc is truncated at the same radius as that of the stars. In addition, we have tried several other cases with variable dust truncations, however, those did not show any significant difference with respect to the ones presented here.

The number of images used to test the dust and inclination influence on the reconstruction is then doubled (except in the 90° cases) by using two different methods of symmetrizing the far and near side of the galaxy, insofar as the dust lanes are concerned. In the first case we average all four quadrants, whereas in the second we only average over the two far side quadrants which have less dust influence and ignore the two near side quadrants with the dust lane.

The artificial images are  $3 \times 3$  pixels carbox smoothed. The deprojection is carried out on the vertically integrated profile, hereafter, the face-on-equivalent view, as well as on the major axis and two higher cuts at  $z = 1.7z_0$  and  $3.4z_0$ . Since the input profile in this case is sharply truncated, we apply only simple exponential radial fits ranging from the centre to three times the radial scalelength, except for the short truncated disc where we go out to one scalelength. From these fits we obtain the scalelength and central surface brightness of the deprojected 3D distribution.

#### 3.6.2 Influence on the measured parameters

To visualize the influence of the various dust distributions we plotted the relative deviation in the recovered scalelength as a function of the inclination angle (Fig. 3). Here the relative deviation is defined explicitly as  $100(h_{\text{true}} - h_{\text{recovered}})/h_{\text{true}}$ .

From Fig. 3, it is obvious that using only the two dust-free quadrants, shown as dashed–crossed lines, improves the quality of the fit. However, above 88° the gain is only minimal and countered by the advantages of an increased S/N – especially important to reach large vertical distances – and by minimizing the residuals from masked objects. Therefore, for the image preparation of the real data we average over all four quadrants, even in the clear presence of dust lanes.


As expected our deprojected, face-on-equivalent, scalelengths are very close to the true, 3D input scalelength in the absence of dust. There is almost no influence of the non-edge-on geometry (down to 80°) of the actual model galaxy which we treat in our deprojection to be always at 90°. For the model galaxies with dust, we see that the error on the scalelength is almost always systematically smaller for galaxies with increasing deviation from the perfect edge-on geometry.

We also notice, from the dust-free case, that for the short-truncated disc the deprojection improves the fits quite noticeably, since it disentangles the truncated part of the profile and avoids fitting for it, as is the case with the projected edge-on view. For the latter we know that they result, slightly depending on the region used for fitting, in a systematic overestimation of the scalelength in the range of 20 per cent (e.g. Pohlen 2001; Kregel et al. 2002; Kregel 2003).

However, we note that our face-on-equivalent scalelengths are still systematically overestimated by typically 15–30 per cent due to neglecting the existence of the dust component. Naturally the effect is far the worse on the individually deprojected major axes ( $h_{\text{in}}$  overestimated by more than 50 per cent), which probably make only sense for dust-deficient S0 galaxies or in the dust-reduced NIR bands. However, the narrowness of the dust lane works in our advantage, so that above 1.4 times the vertical scaleheight the error in the measured scalelength already drops to less than 10 per cent (for  $i \gtrsim 87^\circ$ ). This means the scalelength measured on the individually deprojected higher vertical cuts is much better.

The discussion of the results that we obtain from the observed galaxies takes the effects of the dust and inclination, that we learned from this section, into account.





**Figure 3.** The systematic deviation of the measured scalelength with inclination and dust influence. Plotted is the deviation of the measured from the known input scalelength (in per cent) from fitting the 96 mock images with the same set-up compared to the real galaxies. From left to right the four columns are the results for the face-on-equivalent profiles, for the major axis, and for two cuts at 1.7 and 3.4 $z_0$ . The four rows correspond to the different sharply truncated stellar input models, from top to bottom: not truncated (mimicked by truncating at 10 $h$ ), long truncated (truncated at 5.4 $h$ ), intermediate truncated (truncated at 3.4 $h$ ) and finally short truncated (truncated at 1.4 $h$ ). The different symbols represent the two different ways to obtain an average quadrant: filled circles for the applied mean average out of four, crosses for the average out of two quadrants, excluding the dusty side. The different lines connecting the points in each panel are associated to the different dust distributions (ordered according to influence on the stellar disc). From bottom to top we have the transparent case, the best case, an intermediate case and finally the worst case. See text for more details.

### 3.7 Influence of intrinsic asymmetries

#### 3.7.1 Dust lanes

In the last section we have estimated the influence of the dust on the measured flattening of the inner scalelength using our artificial images. Aside from a solely quantitative correction to be made to the measured scalelength in this case, one could argue that the dust in addition might be responsible for the observed broken exponential structure. In this picture the break would correspond to the point at which the dust lane becomes less important. However, in almost all cases the location of the break does not coincide with the, admittedly difficult to estimate, location of the visible end of the dust lane on the image. An obvious exception is the major axis profile ( $z = 0$  arcsec) of NGC 3390 where the dust lane extends out to  $R \approx \pm 55$  arcsec and indeed causes an apparent break feature at this point (see Fig. 5), hence it is excluded from the fit. In addition, we still find profiles showing a broken exponential structure at more than two times the stellar scaleheight above the disc, where the effect of the dust lane is minimal. Also NGC 4179 does show a break, but has no dust lane at all.

Furthermore, if one assumes that the outer slope is the intrinsic, dust-free slope, the extrapolation into the centre will in almost all cases come up too high (much brighter than the bulge peak). This would require far too much dust in these galaxies. Therefore, the dust quantitatively affects the measured inner scalelengths, however, is not responsible for the break in the profiles (cf. Fig. 4).


#### 3.7.2 Intrinsic small-scale asymmetries

For real galaxies the residuals from masked objects will certainly affect the recovered profile. The problem is ostensibly aggravated by the averaging process of the four quadrants (cf. Section 2.1), which will mirror any residual on one, to all other sides. However, one has to keep in mind that our deprojection method cannot deal with ‘holes’, so we have to somehow fill these masked regions. For example, in the case of NGC 5981 the extended companion superimposed on the upper left-hand side of the bulge could be well traced in the contour lines of the averaged galaxy image (see Fig. 1). Under these worst-case circumstances we made sure that this area is excluded from our fitting region. In the latter case we did not fit beyond  $z > 20$  arcsec.

In addition, the final position of the break radius is strongly affected by the small-scale asymmetries (star-forming regions, dust concentrations, residuals from masked objects) which vary in each of the vertical profiles. Whereas this only slightly affects the measured scalelength, the meeting point of the two exponential fits (break point) is more susceptible to such variations.

#### 3.7.3 Large-scale asymmetries

We do not expect significant large-scale asymmetries in our galaxies, since the input sample is morphologically selected to be symmetric. NGC 5981, which is asymmetric relative to the rest of the sample, is a good test case to estimate the effect of averaging quadrants from both sides. The asymmetry is clearly seen in the radial profiles (cf.



**Figure 4.** Influence of asymmetries: averaging quadrants. The deprojected surface brightness profiles of NGC 5981 using four different ways to combine the four quadrants. The upper left-hand panel and upper right-hand panel show the result using only the two quadrants of the SE and NW side, respectively. The lower left-hand panel shows the profile using only a mean of the two dust-free quadrants. The lower right-hand panel finally shows the resulting profiles after averaging all four quadrants. In each panel, the dashed line (the upper profile) is the face-on-equivalent, radial profile of each galaxy. They are obtained by summing the light of the edge-on galaxy along the columns and deproject the result. The following dotted-dashed lines are the deprojected profiles of individual cuts from top to bottom starting from the major axis towards cuts at larger distance from the plane. The proper distance  $z$  from the plane is given in the caption (together with name and filter) of each panel in the upper right-hand corner. The dotted line (the lower profile) is the deprojected profile of a noise (or source free) cut outside of the galaxy. Overplotted are on each profile two squares marking the fitting region for the disc model. The result of the broken exponential fits are overplotted on each profile as solid lines where the cross marks the break radius.

Fig. B1). The outer slope of the broken exponential fit is significantly shallower on the north-west (NW)(right) side, where also the contour map is more extended. Fitting only a mean quadrant obtained from the NW(right) or south-east (SE)(left) side of the galaxy separately (cf. Fig. 4), we find differences of 5–10 per cent for the scalelengths of the inner region. In the outer disc (beyond the break) the differences are clearly larger, e.g. 40 per cent on the major axis, due to the stronger asymmetry in the outer disc. However, both sides are equally well fit with the broken exponential function, so we do not introduce a bias on the validity of the applied fitting function by averaging the two sides.

## 4 RESULTS

Now we move on to present the results obtained by applying the deprojection method on our observed edge-on galaxies. Again here we first apply the method on the vertically integrated image. This result in a profile that is equivalent to the one obtained from azimuthal averaging, for example by fits with concentric ellipses, of face-on galaxies. A direct comparison between the two views (see Section 3.1) is carried out in Section 4.1.

Furthermore, we deproject a number of vertical slices to explore the existence and properties of, for example, a thick disc component. Here the following questions are addressed: does the inner slope flattens with increasing  $z$ , as expected for some thick disc models? Does the break radius, if present, vary significantly with  $z$ ? Finally, how does the vertical distribution of the outer disc, beyond the break point, changes with  $z$ ? These questions are addressed in Section 4.2.

The radial profiles obtained from deprojecting all the galaxies in the sample together with the best-fitting broken exponential models are shown in Fig. 5. The results for the face-on-equivalent profile of each galaxy in the sample are tabulated in Table 3.


### 4.1 The face-on-equivalent view

#### 4.1.1 Face-on versus edge-on: profile types

As mentioned earlier there are three basic classes of surface brightness profiles that have been found for face-on galaxies, Type I, Type II and Type III (Erwin et al. 2005; Pohlen & Trujillo 2006; Erwin et al., in preparation). Here, in the deprojected face-on-equivalent view, we also find at least one example for each of the three types of surface brightness profiles.

All eight Sbc and later-type galaxies (ESO 380–019, ESO 404–018, FGC 2339, IC 4871, NGC 522, NGC 5290, NGC 5981 and UGC 10459) show clear Type II, downbending profiles. In comparison with Pohlen & Trujillo (2006) and Erwin et al. (in preparation), we cannot increase the level of classification (i.e. assigning subclasses) in order to explore the possible physical origin of the break, for example, star formation threshold or bar related break. This is mainly because inferring accurate bar sizes in edge-on galaxies is almost impossible.

It is interesting to note that the NIR  $K'$ -band profile of NGC 5290, typically assumed to be dust free, shows exactly the same behaviour as the  $R$  band. Therefore, the dust and recent star formation, clearly



**Figure 5.** Deprojected radial profiles of the sample galaxies. The surface brightness profiles of each galaxy are shown in separate panels. The different line types are explained in Fig. 4.

more prominent in the  $R$  band, do not play a role in the basic classification scheme applied here.


The lenticular galaxy, NGC 4179, also exhibits a Type II profile with a ratio of inner to outer scalelength of  $h_{\text{in}}/h_{\text{out}} = 1.4$ . This ratio is somewhat low in comparison to the mean of  $\langle h_{\text{in}}/h_{\text{out}} \rangle = 2.1$  reported by Pohlen & Trujillo (2006) for late-type galaxies, or the median of 2.4 reported by Erwin et al. (in preparation) for early-type galaxies.

One galaxy, NGC 3390, although classified as an Sb galaxy, exhibits a dominant outer spheroidal component closely resembling an S0 galaxy (see Fig. B1). Its face-on-equivalent profile is consistent with being a classical unbroken profile, Type I. Nevertheless, we can also fit to the recovered profile of this galaxy a broken exponential, a Type II. The scalelength ratio in this case is only  $h_{\text{in}}/h_{\text{out}} \lesssim 1.2$ , and so close to unity that in case of a face-on galaxy this would not qualify as a Type II.

The profile of the second S0 galaxy, NGC 1596, is very similar to the antitruncations presented by Erwin et al. (2005). The outer profile is distinctly shallower in slope than the main inner disc profile. Although it should be classified as Type III in this sense, the profile is not well described by our simple broken exponential fitting function. The whole disc profile (beyond  $R \gtrsim \pm 25$  arcsec) appears to be almost continuously upbending and one may argue for an extended, exponential break region between breaks at  $R \approx 70$  and  $\approx 130$  arcsec. Nevertheless, Erwin et al. (2005) and Erwin et al. (in preparation) show some early-type, face-on counterparts with very similar profile shapes.

#### 4.1.2 Face-on versus edge-on: quantitative comparison

As discussed in Section 2 our sample is not unbiased. However, the unusually high frequency of galaxies with Type II profiles suggests

Figure 5 – *continued*

that we indeed have selected against the two other types. For the nine Type II galaxies in our sample, we find that the break radius is in the range between  $1.2$  and  $3.3h_{\text{in}}$ . To compare this with the results of the face-on sample, we have to account for the effect of dust on the inner scalelength. In Section 3.6 we have shown that in the case of dusty edge-on galaxies the resulting scalelengths are typically 20 per cent overestimated (see Fig. 3). Factoring this correction<sup>2</sup> one obtains a break radius in the range of  $1.4$ – $4.0h_{\text{in}}$ , which matches the range given by Pohlen & Trujillo (2006).

The surface brightness at the break radius ranges from  $21.9$  to  $26.3 R\text{-mag/arcsec}^2$ , which also matches the full range for face-on galaxies given by Pohlen & Trujillo (2006). Here, the surface brightness for each galaxy is corrected for galactic extinction and, in the case of the  $V$ -band images, transformed to the  $R$  band assuming  $(V - R) = 0.7$ .

<sup>2</sup> Without the small correction for the different wavelengths used ( $R$  and  $V$  band here compared to  $r'$  band for the face-on sample).

## 4.2 Vertical distribution

### 4.2.1 The inner vertical structure


First, we concentrate on the inner disc region, i.e. the region in between the bulge/bar component and the location of the break. For almost all of the late-type galaxies (Sbc and later, for details see Appendix A) we find an increase in the measured scalelength with the distance from the plane. The increase, at  $3$ – $5z_0$ , is typically of the order of about 20 per cent, relative to the major axis scalelength (see upper panel of Fig. 6). Since we know that the omnipresence of dust extinction will mostly affect the major axis profiles, as evident from the systematic flattening one sees for our artificial galaxies (Fig. 3), our measurements of the increase in scalelength in the optical, starting at the major axis, are only lower limits. Consequently, the major axis slope for four galaxies is indeed observed to be flatter than that obtained from adjacent  $z$  cuts (cf. Fig. 6).

It is interesting to note that this type of flattening towards the mid-plane, which is for  $R$ -band images well explained by the dust

**Table 3.** Exponential disc parameters for the face-on-equivalent profiles.

Galaxy	Filter	Profile type	$b_1$ (arcsec)	$b_2$ (arcsec)	$R_{br}$ (arcsec)	$h_{in}$ (arcsec)	$h_{out}$ (arcsec)	$R_{br}$ ( $h_{in}$ )	$h_{in}$ ( $h_{out}$ )	$\mu_{br}$ (mag/arcsec <sup>2</sup> )	$A_{R,V,K'}$ (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
NGC 522	<i>R</i>	II	22	87	63.1	52.7	8.7	1.2	6.1	22.8	0.23
NGC 1596	<i>R</i>	III	30	210	100.2	27.1	53.8	3.7	0.5	24.4	0.03
NGC 3390	<i>R</i>	I	35	140	–	26.9	–	–	–	–	0.21
NGC 4179	<i>V</i>	II	45	155	97.7	30.9	22.6	3.2	1.4	23.3	0.11
ESO 380–019	<i>V</i>	II	12	117	87.6	27.9	11.5	3.1	2.4	22.9	0.26
NGC 5290	<i>K'</i>	II	30	110	82.5	31.2	13.6	2.7	2.3	21.0	0.00
NGC 5290	<i>R</i>	II	30	125	79.9	33.1	16.3	2.4	2.0	23.7	0.02
NGC 5981	<i>R</i>	II	15	100	66.8	27.9	13.8	2.4	2.0	24.0	0.07
UGC 10459	<i>R</i>	II	10	59	44.2	15.6	5.7	2.8	2.7	25.3	0.03
IC 4871	<i>V</i>	II	20	136	116.5	40.4	9.7	2.9	4.2	24.4	0.27
FGC 2339	<i>R</i>	II	37	66	57.3	17.6	5.9	3.3	3.0	26.4	0.10
ESO 404–018	<i>V</i>	II	0	92	75.6	34.0	9.6	2.2	3.6	24.6	0.13

Notes. (1) Principal name, (2) filter, (3) profile type (I: no break, II: downbending break, III: upbending break), (4,5) inner and outer fitting boundaries, (6,7,8) break radius, inner and outer scalelength in units of arcsec, (9) break radius in relation to the inner scalelength, (10) inner scalelength in relation to the outer scalelength, (11) the surface brightness at the break radius (estimated at the crossing point of the two exponential fits), (12) galactic extinction for the appropriate band from NED according to Schlegel, Finkbeiner & Davis (1998).



**Figure 6.** Measured scalelength with increasing distance from the plane. Upper panel: the open circles are the measured inner scalelengths ( $h_{in}$ ) normalized to the inner scalelengths measured on the major axis  $h_{in,ma}$  for each profile versus the distance ( $z$ ) from the plane normalized to the scaleheight ( $h_z$ ) given in Appendix A. The filled squares mark the values for the outermost profiles. All points for an individual galaxy are connected (solid lines for Type II, dotted–dashed line for Type I and dashed line for Type III) and the name of the galaxy is added. Lower panel: the open circles are the measured outer scalelengths ( $h_{out}$ ) normalized to the outer scalelengths measured on the major axis  $h_{out,ma}$  for each profile versus the distance ( $z$ ) from the plane normalized to the scaleheight ( $h_z$ ).

effect, is also present at the level of approximately 15 per cent in our  $K'$ -band image (NGC 5290). If this is not an intrinsic feature of NGC 5290, one could argue that the dust may still influence the scalelength determination on the major axis even in the NIR band. On the other hand, this seems unlikely since the amount of dust inferred from the  $K'$ -band flattening should have a more severe effect on the  $R$ -band profile (see Appendix A).

In addition, tests with NGC 5981 (see Appendix A) show that using the mean quadrant instead of the dust-free side also results in a systematic underestimation for the measured increase in scalelength with distance from the plane.


#### 4.2.2 The outer vertical structure

Here we focus on the vertical structure of the outer disc, i.e. the region beyond the break radius. For all of the late-type galaxies, namely, Sbc and later, a significant increase of the measured scalelength with distance from the plane is observed (see lower panel of Fig. 6). Whereas for the inner disc the increase in the slope is by a factor of 1.1–1.4, for the outer disc we find a factor of 1.4–2.4. This indicates an extreme flattening of the profile towards larger distances from the plane. Again, the dust, well known for flattening surface brightness profiles, cannot be responsible here since it is concentrated in the plane. This could be regarded as a weakening of the truncation with increasing distance from the plane (see Fig. 7).

#### 4.2.3 The break radius as a function of $z$

The position of the break radius as a function of vertical distance from the plane ( $z$ ) is another constraint that any theory that might explain the truncation phenomenon should satisfy. An inspection of the location of the break radius of the deprojected profiles (Fig. 5) shows that this location does not change significantly as a function of  $z$  in the first two to three  $z$  slices. For the more distant  $z$  slices the slopes of the inner and outer regions become closer, due to the more substantial flattening in the outer disc, which makes it harder to confidently determine the location of the break radius. A clear demonstration of this is when the inner and outer slopes





**Figure 7.** Weakening of truncation with increasing distance from the plane. The open circles are the measured ratios of inner to outer scalelength ( $h_{in}/h_{out}$ ) for each Type II profile with increasing distance ( $z$ ) from the plane normalized to the scaleheight ( $h_z$ ). The filled squares mark the values for the outermost profiles. All points for an individual galaxy are connected (solid lines) and the name of the galaxy is added.

are very close, in such a case the presence of any genuine small-scale structure in the image renders the location of the crossing point very uncertain. This is even more severe when the broken exponential model does not provide a good description of the data, e.g. when the break region is clearly extended (ESO 380–019 or ESO 404–018).

Resorting to the NIR image or concentrating on the smooth S0 galaxies in our sample, in which the substructure (star-forming region or dust patches) are less prominent, does not improve the matter because for these galaxies, as with the other, the flattening of the outer disc dominates the uncertainty in the location of the crossing point.

To summarize, based on the first two to three vertical slices, the current data suggest that the location of the break radius for all our galaxies is consistent with being constant.

## 5 DISCUSSION

### 5.1 Profile classification

In our pilot sample of edge-on galaxies we do indeed find the same three main types of surface brightness profiles as classified by Erwin et al. (2005), Pohlen & Trujillo (2006) and Erwin et al. (in preparation) from their analysis of face-on galaxy images. The broken exponential parameters for the face-on-equivalent view obtained from our edge-on galaxies are well consistent with those found by the analysis of, the more complete, genuine face-on galaxy samples. Only for FGC 2339 the break seems to be at a slightly lower than usual surface brightness, albeit at a typical distance from the centre. However, this is also the galaxy with the value of the lowest central surface brightness. Hence, similar to Pohlen & Trujillo (2006), we still cannot disentangle which of the two statements is correct: the surface brightness at the break radius is constant, or it is coupled to the central surface brightness.

We do not find a difference in the shape of the truncation comparing the two profiles for NGC 5920 obtained from the optical and the NIR image. Both show clearly a broken exponential structure in the region between  $30 \lesssim R \lesssim 110$  arcsec. This is consistent with

the findings of Hunter, Elmegreen & Martin (2006) for the profile shapes of several irregular galaxies using 3.6- and 4.5- $\mu$ m images from the *Spitzer Space Telescope*. Irregular galaxies with double-exponential optical light profiles have the same in the NIR. So barring an exception, NIR observation of truncations do not seem to differ significantly from observation in the optical as proposed by Florido et al. (2006).


Pohlen & Trujillo (2006) have shown that Type I galaxies are rare in late-type galaxies and, according to Erwin et al. (in preparation), more frequent in S0 galaxies. Accordingly, our only Type I candidate (NGC 3390) exhibits an S0-like outer disc structure, despite being classified as Sb. It is interesting to note that in this case the profiles of the individual cuts, close to the major axis but uninfluenced by the dust lane, show some indication of being better described by a broken exponential structure. Therefore, the vertical integration in this case may obscure the existence of a break close to the plane. This is clearly caused by the lack of an apparent break in the vertical slices at larger vertical height above the mid-plane. This might explain the high frequency of not-truncated discs seen in the early-type face-on galaxy sample of Erwin et al. (in preparation).

Despite the discussion above about the general agreement between the edge-on and face-on classifications, an edge-on counterpart of the not truncated, Type I, late-type galaxies (e.g. the face-on NGC 2776 by Pohlen & Trujillo 2006), is yet to be found. Given the expected frequency of such galaxies ( $\approx 10$  per cent) and the size of the parent sample from which we select, we are not able to explain the absence of such galaxies.

If late-type Type III galaxies might be associated with interacting or merging systems, as proposed by Pohlen & Trujillo (2006), the detection of only one Type III galaxy is in full agreement, as we excluded those systems from the original, late-type dominated, sample. In addition, the fact that our Type III galaxy is an early-type one is consistent with the observed increase in frequency of Type III profiles for early-type galaxies (Erwin et al., in preparation). To our knowledge, all edge-on Type III galaxies reported in the literature so far are early types. These galaxies are NGC 4762 studied by several groups (Tsikoudi 1980; Wakamatsu & Hamabe 1984; Wozniak 1994), NGC 4452 shown by Hamabe & Wakamatsu (1989) and NGC 4696C<sup>3</sup> from Pohlen (2001).

For face-on galaxies, Erwin et al. (2005) differentiate between two different types of Type III profiles. One where the excess light is probably part of the disc which shows a relatively sharp transition and both parts are well described with exponentials (Type III-d); and another one where the excess light is associated with the halo/spheroid or bulge (Type III-s). The latter is characterized by a small inner region fitted with an exponential and a continuously upbending outer profile and a changing ellipticity with radius. In the case of NGC 1596 we would argue for a Type III-s classification where the outer structure is associated more with an envelope-like structure. This is supported by the difficulty one faces in fitting NGC 1596 with a consistent thin/thick disc combination (cf. Pohlen et al. 2004b). Any bulge component would have to have an unlikely, extremely high, axis ratio (cf. Appendix A). In this case, the observed outer envelope is therefore much better explained by interaction as recently supported by Chung et al. (2006).

<sup>3</sup> It is worth noticing that NGC 4696C is classified as an Sb galaxy (de Vaucouleurs et al. 1991). However, deep imaging, obtained by Pohlen (2001), clearly shows an outer S0-like envelope.



**Figure 8.** Toy models to explain observed profile structure. Deprojected radial surface brightness profiles of six model galaxies: single disk (panel 1), truncated thin plus not-truncated thick disk (panel 2) and two models with flaring (panels 3–4). The figure has the same set-up as Fig. 5.

## 5.2 Inner disc

Almost all of the analysed profiles exhibit an increase of the radial scalelength with increasing distance from the plane. Notwithstanding the lack of completeness in our sample, this finding could support the notion of *ubiquitous thick discs* as proposed by Dalcanton & Bernstein (2002), where these thick discs have to have a systematically larger scalelength, compared to the thin disc. However, in the absence of a consistent two disc fit to the data at all radii the thick disc model is not the only option. Instead one can interpret the increase in the radial scalelength merely as a result either of a radially flattened component above the inner disc or a flattening of the inner disc itself. For example, in the case of the late-type (Scd) galaxy FGC 2339, the whole inner disc region is dominated by a ring- or bar-like structure in the mid-plane which also alters the shape of the disc above/below the plane over the entire measured vertical extent.

## 5.3 Outer disc

In order to examine what could lead to the observed flattening of the outer disc profile, we invoke several 3D toy models. These 3D models are generated by combining separate components, e.g. a truncated thin disc plus a truncated thick disc. The resultant structures are qualitatively compared to those reconstructed from the data. It is important to emphasize here that no attempt is made to fit the data to these models, rather the intention is to explore whether these toy models could generically reproduce the increased flattening seen in the data. The 3D profiles of the models are shown in Fig. 8.

To highlight the flattening in these models we fit broken exponential profiles with the same fitting routine used for the real galaxies (cf. Section 3.5). For example the upper left-hand panel (1) of Fig. 8

shows a single broken exponential disc model. This type of models does not show any flattening in the reconstructed scalelengths and, therefore, is not favoured by our data.

### 5.3.1 Could we be seeing de Vaucouleurs $R^{1/4}$ bulges?

Bahcall & Kylafis (1985) have shown that a thin disc plus a  $R^{1/4}$  bulge can mimic the shape of the isophotes of a thin disc plus thick disc model (without bulge) down to a surface brightness level of about  $\mu_R \sim 25.5 \text{ mag/arcsec}^2$ . Furthermore, this combination can even produce some flattening of the radial profiles with increasing vertical height beyond the break radius as seen in our models.

Having said that, however, there are strong arguments against explaining the observed flattening with a spherical power-law bulge component. First, in a sample of 19 early-type galaxies (S0–Sbc) Balcells et al. (2003) find that none of the bulges have a  $R^{1/4}$  behaviour. Most of their bulges are better fit with exponentials which are not expected to dominate the radial profiles at large distances. Therefore, it is statistically unlikely to attribute the flattening in the outer disc in most of our late-type edge-on galaxies to a spherical bulge component. Second, the observed shape of the outer contours in all the cases studied here appears too much disc like to be caused by a spheroid (see Figs B1 and 9). Third, it is almost impossible to create a  $R^{1/4}$  bulge which will leave the high S/N major axis profile purely exponential down to the last measured point, but will dominate the radial profile at larger vertical heights above the plane, and doing this all without being clearly visible in a contour map.

In conclusion, since there is no clear sign of a de Vaucouleurs  $R^{1/4}$  in any of the observed images whereas the outer disc flattening



is found in the majority of galaxies in the sample, the answer to the question posed here is, most probably, no.

### 5.3.2 Does a thin plus thick disc combination produce the increased flattening beyond the break?

If one assumes that the thick disc scalelength is in general larger than the thin disc scalelength, the increase in scalelength with  $z$  height in the inner disc will be always present. However, there are various possibilities for producing the extended flattening in the outer disc beyond the break with thin plus thick disc models. The exact shape of the flattening critically depends on the, unknown, outer disc properties of the thick disc.

If, for example, we assume the thick disc to be not truncated, we are still able to produce some flattening beyond the break (see panel 2 of Fig. 8). However, depending on the surface brightness limit the not-truncated thick disc will become visible at large radii. This could be an explanation for some of the very few Types II + III mixed classifications found for late-type galaxies by Pohlen & Trujillo (2006). These galaxies show two breaks in their face-on surface brightness profile. The first is downbending in the inner region, while the other is upbending in the very outer region. However, none of our edge-on sample galaxies shows this behaviour.

By adding a truncation to the thick disc one can produce a wide variety of broken exponential structures depending, for example, on the ratio of inner to outer disc scalelength ( $h_{\text{in}}/h_{\text{out}}$ ). Models with unequal thin and thick disc truncation radii provide even more choice to fit the data.

In conclusion, the answer to the question posed here is yes. However, the degree of flattening will depend on the details of the 3D model.


### 5.3.3 Could the flattening be explained by intrinsic flaring<sup>4</sup> of the stellar disc?

To produce an increased flattening in the outer disc one can invoke models with intrinsic flaring – similar to those used for the Milky Way (see e.g. Gyuk, Flynn & Evans 1999) – that starts beyond the break radius. In the edge-on view, the contour map of the observed galaxy provides a strong constraint on the degree of flaring. The lower panel of Fig. 9 shows three different radially broken exponential single disc models with flaring factors<sup>5</sup> of  $F = 2.5, 5$  and  $10$ . If the flaring factor is chosen to be too large the resulting contour map will blow up in the outer parts, a behaviour that is observed neither in any of our sample galaxies nor in the much larger sample of Pohlen (2001). In general, it seems that a strong flaring of the stellar disc is rarely observed. In addition, in the few reported cases it appears to be rather associated with Type III profiles, e.g. NGC 4762 (Wakamatsu & Hamabe 1984; Freeman & Bland-Hawthorn 2002), and not with Type II as discussed here.

As shown in the panel (3) of Fig. 8, flaring of a single disc beyond the break radius does not always produce the right amount of flattening if compared to our observations. A more promising attempt

<sup>4</sup> With *intrinsic flaring* we do not mean the sometimes observed fact of an increase in scaleheight with radius (e.g. de Grijs & Peletier 1997), which is most probably related to a combination of thin and thick disc with different scalelengths, but the intrinsic change of the scaleheight for a single stellar disc component.

<sup>5</sup> For the Milky Way, Gyuk et al. (1999) have models with  $F = 5$  or  $10$  between the solar radius and 25 kpc.



**Figure 9.** Upper panel: contour map of a radially broken exponential, vertically exponential thin disc superimposed with a standard de Vaucouleurs  $R^{1/4}$  bulge. Lower panel: three combination of a the same thin disc and a thick disc, where the thick disc scaleheight grows (flares) between the break radius ( $R_{\text{br}}$ ) and three times  $R_{\text{br}}$  with an increasing factor (solid, dashed, dotted lines).

to apply flaring is obtained by allowing only the thick disc component to flare beyond the break radius (see panel 4 of Fig. 8). This approach could help in cases where the inner disc indicates only a very small value for the ratio of thin to thick disc scalelength. The flaring is then needed to reproduce the observed significant difference of the outer scalelength on the major axis compared to high above the plane. A flaring of the thick disc is also proposed by several authors (e.g. Siegel et al. 2002; Du et al. 2006) for our Milky Way from star-count studies. This flaring would account for the observed variable scaleheight with different direction of the analysed fields.

It is clear that the data can in principle discriminate between possible 3D models, as each will produce different features in the reconstructed profiles. However, we make no such attempt in this paper except to point out the different possibilities.

## 5.4 Break radius

As we have shown earlier, the location of the break radius in the current sample is found to be consistent with being height independent. In light of the enhanced flattening of the outer disc relative to the inner disc, the determination of the break radius at large vertical cuts is uncertain and more accurate modelling could reveal significant variation in its location towards the larger distance from the major axis.

Nevertheless, we would like to point out that if such a finding is supported by more data, it could have far reaching implications on theories that attempt to understand how the thin and thick disc are

assembled. As such a theory should explain how potentially distinct components of the disc ‘coordinate’ the location of their break radii.

## 6 CONCLUSIONS

This paper has presented a new algorithm for reconstructing the intrinsic 3D structure of galactic discs seen edge-on. The algorithm assumes axial symmetry and utilizes the so-called Fourier slice theorem which in the case of edge-on galaxies is equivalent to an inverse Abel transform. The algorithm is applied to a pilot set of 12 edge-on galaxy images, drawn from a parent sample of 72 galaxies, to obtain face-on-equivalent radial surface brightness profiles, and reconstruct the radial profiles along several vertical cuts, including the major axis.

Comparing the results obtained from the face-on-equivalent profiles with those measured from ‘real’ face-on samples we reach several conclusion. The first, and probably most important, conclusion is that we find in the current edge-on sample at least one example for each of the three basic profile types found in face-on samples: no break, upbending break and downbending break (Erwin et al. 2005; Pohlen & Trujillo 2006; Erwin et al., in preparation). Despite this agreement, our morphologically selected parent sample (Pohlen 2001), choosing undisturbed and prototypical disc galaxies, clearly favours late-type galaxies with class Type II profiles, i.e. downbending breaks. This might be due to a connection between certain profile types and distinct vertical structures. For example, if late-type Type III galaxies are connected to interaction and/or merging processes, they would probably be excluded from edge-on samples. In other words, they would exhibit distortions, lopsidedness or warping, which are all prominent in the edge-on view.

Although our parent sample is not unbiased in this sense, our results do show that the parameters for the downbending breaks of the face-on-equivalent profiles obtained from our edge-on galaxies are consistent with those seen in face-on samples. This implies that truncated profiles of galaxies are independent of geometry and there is no distinction in this regard between edge-on and face-on studies.

In this paper we have been able to study, for the first time, the deprojected radial profiles as a function of vertical height above the plane, which is only possible to explore with edge-on galaxies. Hence, this approach provides a unique opportunity to give additional constraints on theories explaining the origin of the various types of observed profiles in the context of disc formation and evolution.

In this context, we find a generic flattening of the observed profiles in the inner disc region with increasing distance from the plane, which cannot be explained by dust extinction. This implies a more complex vertical structure in the inner region than a single disc. It is possible to explain this flattening by introducing a thick disc with systematically larger scalelength compared to that of its host thin disc.

In the outer disc, beyond the break radius, we find an enhanced flattening of the deprojected profiles, which again sets strong constraints on theories that attempt to explain the observed broken exponential structures. Using some toy models we conclude that thin plus thick disc combinations are, in principle, able to explain the observed structures. However, the details of this combination depend on the exact shape of the outer region of the thick disc. For example, a thick disc without truncation will exhibit less flattening in the outer disc than a thick disc with truncation. So far our data do not allow us to decide if thick discs are in general truncated or not. It remains unclear whether one can fine-tune a simple model

with broken exponential thin plus thick discs in order to explain the observed brightness distribution.

In addition, we find that the location of the break radius does not change significantly as a function of  $z$  in the first two to three  $z$  slices. Unfortunately, the flattening of the radial profile beyond the first few vertical cuts makes it hard to determine the location of the break radius. Although we do not know yet whether thick discs are indeed truncated, we have shown that the vertical structure harbours clues to understand the origin of the different radial profiles and should be pursued. For example, luminous, not-truncated thick disc components could be responsible for the high frequency of not-truncated lenticular galaxies in face-on surveys, while having a broken exponential structure close to the mid-plane of the galaxy.

In conclusion, observations of the vertical structure set strong constraints on modelling/simulating disc galaxies and the interpretation of their various components. In the future, a larger, well-selected sample of edge-on galaxies will be analysed in order to shed light on some of the issues raised here.

## ACKNOWLEDGMENTS

The authors wish to thank Simone Bianchi, who kindly provided us with *dusty galaxy images* produced with his radiative transfer code. We thank Olof van den Berg for using the reduced  $K'$ -band image of NGC 5290. The authors thank P. van der Kruit, R. Sancisi, M. Verheijen and E. Valentijn for helpful discussions, and the anonymous referee for his detailed and very helpful comments. Part of this work was supported by a Marie Curie Intra-European Fellowship within the Sixth European Community Framework Programme. This research has made use the LEDA (<http://leda.univ-lyon1.fr>) and the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research has made use of NASA’s Astrophysics Data System Bibliographic Services.

## REFERENCES

- Bahcall J. N., Kyllafis N. D., 1985, *ApJ*, 288, 252
- Balcells M., Graham A. W., Domínguez-Palmero L., Peletier R. F., 2003, *ApJ*, 582, L79
- Barteldrees A., Dettmar R.-J., 1994, *A&AS*, 103, 475
- Bertin E., Arnouts S., 1996, *A&AS*, 117, 393
- Bianchi S., Ferrara A., Giovanardi C., 1996, *ApJ*, 465, 127
- Binney J. J., Tremaine S., 1987, *Galactic Dynamics*. Princeton Univ. Press, Princeton, NJ
- Binney J. J., Davies R. L., Illingworth G. D., 1990, *ApJ*, 361, 78
- Bland-Hawthorn J., Vlajić M., Freeman K. C., Draine B. T., 2005, *ApJ*, 629, 239
- Byun Y.-I., 1998, *Chin. J. Phys.*, 36, 677
- Chung A., Koribalski B., Bureau M., van Gorkom J. H., 2006, *MNRAS*, 370, 1565
- Dalcanton J. J., Bernstein R. A., 2002, *AJ*, 124, 1328
- Daubechies I., 1988, *Commun. Pure Appl. Math.*, 41, 909
- Debbatista V. P., Mayer L., Carollo C. M., Moore B., Wadsley J., Quinn T., 2006, *ApJ*, 645, 209
- de Grijs R., 1998, *MNRAS*, 299, 595
- de Grijs R., Peletier R. F., 1997, *A&A*, 320, L21
- de Grijs R., Kregel M., Wesson K. H., 2001, *MNRAS*, 324, 1074
- Dehnen W., Gerhard O. E., 1993, *MNRAS*, 261, 311
- Dehnen W., Gerhard O. E., 1994, *MNRAS*, 268, 1019
- de Vaucouleurs G., 1959, *Handb. Phys.*, 53, 311

- de Vaucouleurs G., de Vaucouleurs A., Corwin H. G., Buta R. J., Paturel G., Fouque J. B., 1991, *Third Reference Catalogue of Bright Galaxies*. Springer-Verlag, New York
- de Zeeuw T., 2007, in de Jong R. S., ed., *Astrophys. Space Sci. Proc. Vol. 3, Island Universes: Structure and Evolution of Disc Galaxies*. Springer-Verlag, Dordrecht, p. 571
- Donoho D. L., 1995, *IEEE Transactions on Information Theory*, 41, 613
- Donoho D. L., Johnstone I. M., 1994, *Biometrika*, 81, 425
- Du C., Ma J., Wu Z., Zhou X., 2006, *MNRAS*, 372, 1304
- Elmegreen B. G., Hunter D. A., 2006, *ApJ*, 636, 712
- Erwin P., Beckman J. E., Pohlen M., 2005, *ApJ*, 626, L81
- Fabricant D., Rybicki G., Gorenstein P., 1984, *ApJ*, 286, 186
- Ferguson A., Irwin M., Chapman S., Ibata R., Lewis G., Tanvir N., 2007, in de Jong R., ed., *Ap&SS Proc. Vol. 3, Island Universes: Structure and Evolution of Disk Galaxies*. Springer, Berlin, p. 239
- Florido E., Battaner E., Guíjarro A., Garzón F., Jiménez-Vicente J., 2001, *A&A*, 378, 82
- Florido E., Battaner E., Guíjarro A., Garzón F., Castillo-Morales A., 2006, *A&A*, 455, 467
- Freedman W. L. et al., 2001, *ApJ*, 553, 47
- Freeman K. C., 1970, *ApJ*, 160, 811
- Freeman K., Bland-Hawthorn J., 2002, *ARA&A*, 40, 487
- Gerhard O., Binney J., 1996, *MNRAS*, 279, 993
- Governato F., Willman B., Mayer L., Brooks A., Stinson G., Valenzuela O., Wadsley J., Quinn T., 2007, *MNRAS*, 374, 1479
- Gyuk G., Flynn C., Evans N. W., 1999, *ApJ*, 521, 190
- Hamabe M., Wakamatsu K.-I., 1989, *ApJ*, 339, 783
- Holley-Bockelmann J. K., Mihos J. C., 2001, *BAAS*, 33, 798
- Hunter D. A., Elmegreen B. G., 2006, *ApJS*, 162, 49
- Hunter D. A., Elmegreen B. G., Martin E., 2006, *AJ*, 132, 801
- Jensen E. B., Thuan T. X., 1982, *ApJS*, 50, 421
- Kregel M., 2003, PhD thesis, Univ. Groningen
- Kregel M., van der Kruit P. C., de Grijs R., 2002, *MNRAS*, 334, 646
- Lauberts A., 1982, *ESO/Uppsala Survey of the ESO(B) Atlas*. European Southern Observatory (ESO), Garching
- Li Y., Mac Low M.-M., Klessen R. S., 2006, *ApJ*, 639, 879
- Lucy L., 1974, *AJ*, 79, 745
- Matthews L. D., 2000, *AJ*, 120, 1764
- Möllenhoff C., Popescu C. C., Tuff's R. J., 2006, *A&A*, 456, 941
- Näslund M., Jörsäter S., 1997, *A&A*, 325, 915
- Nelder J. A., Mead R., 1965, *Comput. J.*, 7, 308
- Patterson F. S., 1940, *Harv. Coll. Obser. Bull.*, 914, 9
- Pérez I., 2004, *A&A*, 427, L17
- Pohlen M., 2001, PhD thesis, Ruhr-University Bochum, Germany
- Pohlen M., Trujillo I., 2006, *A&A*, 454, 759
- Pohlen M., Dettmar R.-J., Lütticke R., 2000a, *A&A*, 357, L1
- Pohlen M., Dettmar R.-J., Lütticke R., Schwarzkopf U., 2000b, *A&AS*, 144, 405
- Pohlen M., Dettmar R.-J., Lütticke R., Aronica G., 2002a, *A&A*, 392, 807
- Pohlen M., Dettmar R. J., Lütticke R., Aronica G., 2002b, in Athanassoula E., Bosma A., Mujica R., eds, *ASP Conf. Ser. Vol. 275, Discs of Galaxies: Kinematics, Dynamics and Perturbations*. Astron. Soc. Pac., San Francisco, p. 15
- Pohlen M., Balcells M., Lütticke R., Dettmar R.-J., 2003, *A&A*, 409, 485
- Pohlen M., Beckman J. E., Hüttemeister S., Knapen J. H., Erwin P., Dettmar R.-J., 2004a, in Block D. L., Puerari I., Freeman K. C., Groess R., Block E. K., eds, *Ap&SS Lib. Vol. 319, Penetrating Bars Through Masks of Cosmic Dust*. Kluwer, Dordrecht, p. 713
- Pohlen M., Balcells M., Lütticke R., Dettmar R.-J., 2004b, *A&A*, 422, 465
- Prugniel P., Heraudeau P., 1998, *A&AS*, 128, 299
- Rybicki G. B., 1987, in de Zeeuw P. T., ed., *IAU Symp. 127, Structure and Dynamics of Elliptical Galaxies*. Kluwer, Dordrecht, p. 397
- Sandage A., Tammann G. A., 1980, *A Revised Shapley-Ames Catalog of Bright Galaxies*. Carnegie Institution, Washington
- Sasaki T., 1987, *PASJ*, 39, 849
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
- Siegel M. H., Majewski S. R., Reid I. N., Thompson I. B., 2002, *ApJ*, 578, 151
- Tamm A., Tenjes P., 2006, *A&A*, 449, 67
- Trujillo I., Pohlen M., 2005, *ApJ*, 630, L17
- Tsikoudi V., 1980, *ApJS*, 43, 365
- van der Kruit P. C., 1979, *A&AS*, 38, 15
- van der Kruit P. C., Searle L., 1981, *A&A*, 95, 105
- Wakamatsu K.-I., Hamabe M., 1984, *ApJS*, 56, 283
- Weinberg M. D., 2001, in Woodward C. E., Bica M. D., Shull J. M., eds, *ASP Conf. Ser. Vol. 231, Tetons 4: Galactic Structure, Stars and the Interstellar Medium*. Astron. Soc. Pac., San Francisco, p. 53
- Wozniak H., 1994, *A&A*, 286, L5
- Xilouris E. M., Byun Y. I., Kylafis N. D., Paleologou E. V., Papamastorakis P., 1999, *A&A*, 344, 868
- Yoachim P., Dalcanton J. J., 2006, *AJ*, 131, 226
- Zaroubi S., Goelman G., 2000, *Magn. Reson. Imaging*, 18, 59
- Zaroubi S., Squires G., Hoffman Y., Silk J., 1998, *ApJ*, 500, L87
- Zaroubi S., Squires G., de Gasperis G., Evrard A. E., Hoffman Y., Silk J., 2001, *ApJ*, 561, 600

## APPENDIX A: NOTES ON INDIVIDUAL GALAXIES

Below we describe the observed profiles and give some extended notes on each individual galaxy (sorted alphabetically). A summary of the exponential disc parameters for all fitted profiles of each galaxy can be found in Table A1.

### ESO 380—019

Owing to a not well-defined thick patchy dust lane the inclination is uncertain (clearly not  $90^\circ$ , but closer to  $85^\circ$ ). The extent and size of the bulge component is also uncertain. The major axis radial profile shows a broken exponential structure with a break at  $R \approx \pm(95-100)$  arcsec ( $\mu_V \approx 23.8$  mag/arcsec<sup>2</sup>). The galaxy is moderately asymmetric. In the contour map the SE(left) side is obviously shorter. The single disc fit by Pohlen (2001) yields a vertical scaleheight of  $h_z = 3.5$  arcsec, a radial scalelength of  $h = 22.8$  arcsec and an inclination of  $84.0^\circ$ .

The face-on-equivalent profile of ESO 380—019 is well described as a Type II with an inner shallow exponential (of  $h_{in} = 27.9$  arcsec  $\equiv 2.4h_{out}$ ) and an outer steeper exponential ( $h_{out} = 11.5$  arcsec) with a break at a radial distance of  $R_{br} \approx 88$  arcsec  $\equiv 3.1h_{in}$  and a surface brightness of  $\mu_{br} = 22.9$  V-mag/arcsec<sup>2</sup>.

We find an increase in scalelength of  $\approx 25$  per cent between the major axis and about five times the vertical scaleheight (at 18 arcsec) in the inner disc region. Since the galaxy is rather far from edge-on, the extended dust lane (with maximum extinction at  $\approx 7$  arcsec) could still play a role (cf. Fig. 3). The scalelength beyond the break in the outer disc increases by a factor of 2.5 between 0 and  $5.1h_z$ .

### ESO 404—018

This galaxy shows hardly any bulge component. The disc structure is quite knotty and appears bended like a wave (Weinberg 2001) in addition to being slightly u-shaped at the edges. The radial surface brightness profile shows quite clearly the deviation from a simple not-truncated exponential model on both sides with a sharp break on the NE(left) side, and a smoother one on the SW(right) side starting at  $R \approx \pm 70$  arcsec and is well fitted by a broken exponential. The single disc fit by Pohlen (2001) yields a vertical scaleheight of  $h_z = 2.8$  arcsec, a radial scalelength of  $h = 31.3$  arcsec and an inclination of  $90.0^\circ$ .

The face-on-equivalent profile of ESO 404—018 is well described as a Type II with an inner shallow exponential (of

$h_{\text{in}} = 34.0 \text{ arcsec} \equiv 3.6h_{\text{out}}$ ) and an outer steeper exponential ( $h_{\text{out}} = 9.6 \text{ arcsec}$ ) with a break at a radial distance of  $R_{\text{br}} \approx 76 \text{ arcsec} \equiv 2.2h_{\text{in}}$  and a surface brightness of  $\mu_{\text{br}} = 24.6 \text{ V-mag/arcsec}^2$ .

We find an increase in scalelength of  $\approx 12$  per cent between the major axis and about four times the mean vertical scaleheight (at  $11 \text{ arcsec}$ ) in the inner disc region. Although this galaxy is almost perfectly edge-on we still find the steepest profile on the major axis, so the dust is not playing a major role here. The shape of the profile in the outer disc is not well described with a single exponential. The break regions (starting at  $\approx 75 \text{ arcsec}$ ) seems to be extended and almost exponential itself. This deviation from our simple broken exponential model strongly affects the measurement of the scalelengths beyond the break.

### FGC 2339

This galaxy shows on both sides slightly off-centred ring-, or bar-like knots at  $(R, z) = (+24 \text{ arcsec}, +1 \text{ arcsec})$  and  $(R, z) = (-26 \text{ arcsec}, +1 \text{ arcsec})$ . The underlying disc is quite asymmetric on both sides. The major axis profile on the N(left) side is satisfactorily fitted with a broken exponential showing a break at  $R \approx \pm(55\text{--}60) \text{ arcsec}$  and  $\mu_R \approx 24.0 \text{ mag/arcsec}^2$ . On the S(right) side the profile looks more like a sharply truncated profile. The single disc fit by Pohlen (2001) yields a vertical scaleheight of  $h_z = 1.8 \text{ arcsec}$ , a radial scalelength of  $h = 19.9 \text{ arcsec}$  and an inclination of  $88.5^\circ$ .

The face-on-equivalent profile of FGC 2339, beyond  $35 \text{ arcsec}$  can be described as a Type II with an inner shallow exponential (of  $h_{\text{in}} = 17.6 \text{ arcsec} \equiv 3.0h_{\text{out}}$ ) and an outer steeper exponential ( $h_{\text{out}} = 5.9 \text{ arcsec}$ ) with a break at a radial distance of  $R_{\text{br}} \approx 57 \text{ arcsec} \equiv 3.3h_{\text{in}}$  and a surface brightness of  $\mu_{\text{br}} = 26.4 \text{ R-mag/arcsec}^2$ .

For this galaxy the whole inner region  $R \lesssim 35 \text{ arcsec}$  is dominated by a ring or bar-like structure which introduces an extreme flattening of the profiles at 2 vertical scaleheights and above (cf. van den Berg et al., in preparation). The small inner disc region in the range of  $35 \lesssim R \lesssim 55 \text{ arcsec}$  does not show a clear systematic flattening towards higher  $z$  cuts. The major axis profile is overall the flattest, so probably heavily affected by the mid-plane dust. Nevertheless, there is a 19 per cent increase in scalelength between  $2.3$  and  $3.4h_z$ . The scalelength beyond the break in the outer disc increases by a factor of  $2.3$  between  $0$  and  $2.3 h_z$ .

### IC 4871 (alternative name: IC 4872)

This galaxy is overall very patchy with many distinct H II regions. On the S(right) side, along the major axis at  $R = +100 \text{ arcsec}$  there is an unusually warped patch of star formation visible. The major axis radial profile can be well described with a broken exponential with a very steep outer decline (dropping by  $\approx 2 \text{ mag}$  within  $\approx 10 \text{ arcsec}$ ) on both sides (with slightly different outer slopes), starting at  $R \approx \pm 115 \text{ arcsec}$  ( $\mu_V \approx 23.5 \text{ mag/arcsec}^2$ ). The single disc fit by Pohlen (2001) yields a vertical scaleheight of  $h_z = 3.8 \text{ arcsec}$ , a radial scalelength of  $h = 42.3 \text{ arcsec}$  and an inclination of  $87.5^\circ$ .

The face-on-equivalent profile of IC 4871 is well described as a Type II with an inner shallow exponential (of  $h_{\text{in}} = 40.4 \text{ arcsec} \equiv 4.2h_{\text{out}}$ ) and a very steep outer exponential ( $h_{\text{out}} = 9.7 \text{ arcsec}$ ) with a break at a radial distance of  $R_{\text{br}} \approx 117 \text{ arcsec} \equiv 2.9h_{\text{in}}$  and a surface brightness of  $\mu_{\text{br}} = 24.4 \text{ V-mag/arcsec}^2$ .

We find an increase in scalelength of  $\approx 13$  per cent between the major axis and about 2.5 times the mean vertical scaleheight (at  $8 \text{ arcsec}$ ) in the inner region. The scalelength beyond the break in the outer disc increases by a factor of  $1.7$  between  $0$  and  $2.5 h_z$ .

### NGC 522

Our image has some residual structure in the flat-field which reaches the disc on the SW(right) side at  $(R, z) = (+10 \text{ arcsec}, -15 \text{ arcsec})$  and at  $(R, z) = (+60 \text{ arcsec}, -30 \text{ arcsec})$  causing the unusually shaped contours. There is a galaxy cluster in the background with an edge-on spiral galaxy superimposed at  $(R, z) = (-13 \text{ arcsec}, -17 \text{ arcsec})$  and another elliptical at  $(R, z) = (-60 \text{ arcsec}, -20 \text{ arcsec})$ . The major axis radial profile shows a clear broken exponential behaviour with a very steep outer decline (dropping by  $\approx 5 \text{ mag}$  within  $\approx 25 \text{ arcsec}$ ) on both sides, starting at  $R \approx \pm 60 \text{ arcsec}$  ( $\mu_R \approx 21.5 \text{ mag/arcsec}^2$ ). Even the edge-on profiles show clearly a decrease in the outer slope with increasing  $z$  (see below). The galaxy is slightly asymmetric with the NE(left) side shorter than the SW(right) side. The single disc fit by Pohlen (2001) yields a vertical scaleheight of  $h_z = 4.3 \text{ arcsec}$ , a radial scalelength of  $h = 32.0 \text{ arcsec}$  and an inclination of  $87.5^\circ$ .

The face-on-equivalent profile of NGC 522 is well described as a Type II with an inner shallow exponential (of  $h_{\text{in}} = 52.7 \text{ arcsec} \equiv 6.1h_{\text{out}}$ ) and an outer very steep exponential ( $h_{\text{out}} = 8.7 \text{ arcsec}$ ) with an ‘early’ break at a radial distance of  $R_{\text{br}} \approx 63 \text{ arcsec} \equiv 1.2 h_{\text{in}}$  and a surface brightness of  $\mu_{\text{br}} = 22.8 \text{ R-mag/arcsec}^2$ .

The major axis profile is overall the flattest, so probably heavily affected by the mid-plane dust. We find only a marginal increase in scalelength of  $\approx 5$  per cent between  $0.9$  and  $3.1$  times the mean vertical scaleheight (between  $4$  and  $13 \text{ arcsec}$ ) in the inner region. The scalelength beyond the break in the outer disc increases by a factor of  $1.8$  between  $0$  and  $3.1h_z$ .

### NGC 1596

See Pohlen et al. (2004b) for additional notes. The face-on-equivalent profile of NGC 1596 is classified as Type III, i.e. a break with an upbending profile beyond. However, the profile is not well described by a simple broken exponential. The whole disc profile (beyond  $R \gtrsim \pm 25 \text{ arcsec}$ ) appears to be almost continuously upbending. One may argue for an extended, exponential break region between breaks at  $R \approx 70$  and  $\approx 130 \text{ arcsec}$ . Therefore, the exact position of the break is unclear. Judging from the individually deprojected cuts this structure is driven by the clear upbending breaks on the major axis and at  $z = 8 \text{ arcsec}$  at a radial distance of  $R \approx 110 \text{ arcsec}$ . All cuts at larger distance from the plane ( $z \gtrsim 17 \text{ arcsec}$ ) are consistent with having no break.

We find an increase in scalelength by a factor of  $3.6$  from the major axis (steepest) to a vertical scaleheight of  $33 \text{ arcsec}$  (flatter) in the inner region.

### NGC 3390

There is a nearby, low surface brightness companion, of unknown distance, visible at  $(R, z) = (+140 \text{ arcsec}, -125 \text{ arcsec})$  and another unusually small and round object at  $R = +85 \text{ arcsec}$  on the major axis. This galaxy exhibits a quite unusual appearance. The inner high surface brightness part looks like a normal edge-on Sb galaxy with a well-defined dust-lane. However, the outer low surface brightness part shows that the galaxy is embedded in a huge spheroidal envelope much more like an S0. The galaxy was already classified as an S0-a by Lauberts (1982) and S0? by Sandage & Tammann (1980), whereas de Vaucouleurs et al. (1991) and Prugniel & Heraudeau (1998) quote Sb or Sbc. The radial profiles close to the major axis, similar to other S0 galaxies, decline almost straight into the noise with only a mild break at  $R \approx \pm 90 \text{ arcsec}$  in cuts higher above the plane. This structure makes it impossible to fit a sharply truncated or

broken exponential model. Even the edge-on profiles show clearly a decrease in the outer slope with increasing  $z$  (see below).

The face-on-equivalent profile of IC 3390 is reasonably well described as a Type I (see discussion in main text, Section 4.1). For NGC 3390 we find an increase in scalelength by a factor of 3.1 between the major axis (steepest) and a vertical distance of 26 arcsec (flatter) in the inner disc region as expected from the S0 envelope. A scaleheight of 5.3 arcsec, measured in the  $I$  band by de Grijs (1998), is used in Figs 6 and 7.

#### NGC 4179

See Pohlen et al. (2004b) for additional notes. We find that the face-on-equivalent profile of NGC 4179 is well described as a Type II with an inner shallow exponential (of  $h_{\text{in}} = 30.9$  arcsec  $\equiv 1.4h_{\text{out}}$ ) and an outer slightly steeper exponential ( $h_{\text{out}} = 22.6$  arcsec) with a break at a radial distance of  $R_{\text{br}} \approx 98$  arcsec  $\equiv 3.2 h_{\text{in}}$  and at a surface brightness of  $\mu_{\text{br}} = 23.3$   $V$ -mag arcsec $^{-2}$ .

In the inner disc we find an increase in scalelength of 36 per cent between the major axis and 4.5 times the thin disc scaleheight (6.1 arcsec) reported by Pohlen et al. (2004b). Overall the scalelength increases by a factor of 1.8 to the highest measured cut at  $z = 45$  arcsec. This is very similar to the factor of 1.9 reported by Pohlen et al. (2004b) for the ratio of thick to thick disc scalelength, obtained with a completely different method (3D model fitting). Their thin disc scalelength of 26.2 arcsec is almost identical to our scalelength of 26.7 arcsec at the major axis. Our scalelength of 46.8 arcsec for the vertical cut with the largest distance from the plane (at  $z = 45$  arcsec) is  $\approx 9$  per cent smaller compared to their thick disc scalelength of 51.0 arcsec, which we may reach for cuts even higher above.

#### NGC 5290

The inner disc is embedded in an outer, clearly thicker disc component or in a low surface brightness extension of the bulge. The galaxy exhibits a thick and patchy dust lane, and the two sides are rather asymmetric. In the contour map the E(left) side seems to be thicker and shorter, whereas the W(right) side is longer and tapers off. This asymmetry is reflected in the major axis radial profile. When fitted with a broken exponential structure, the slope of the outer region on the E(left) side is much steeper compared to the W(right) side. The latter seems to be almost  $\approx 15$  arcsec ‘longer’. The single disc fit to the  $R$ -band image by Pohlen (2001) yields a vertical scaleheight

of  $h_z = 7.5$  arcsec, a radial scalelength of  $h = 25.4$  arcsec and an inclination of  $88.5^\circ$ .

The face-on-equivalent profile of NGC 5290 is in both bands ( $R$  and  $K'$ ) well described as a Type II with an inner shallow exponential (of  $h_{\text{in}} = 33.1$  arcsec  $\equiv 2.0 h_{\text{out}}$  in  $R$  and  $h_{\text{in}} = 31.2$  arcsec  $\equiv 2.3 h_{\text{out}}$  in  $K'$ ) and an outer steeper exponential ( $h_{\text{out}} = 16.3$  arcsec in  $R$  and  $h_{\text{out}} = 13.6$  arcsec in  $K'$ ). The break radius is at a similar radial distance ( $R_{\text{br}} \approx 80$  arcsec  $\equiv 2.4 h_{\text{in}}$  in  $R$ , and  $R_{\text{br}} \approx 83$  arcsec  $\equiv 2.7 h_{\text{in}}$  in  $K'$ ) and at a surface brightness of  $\mu_{\text{br}} = 23.7$   $R$ -mag arcsec $^{-2}$  and  $\mu_{\text{br}} = 21.0$   $K'$ -mag/arcsec $^2$ , respectively.

In the  $K'$  band, the scalelength of the major axis in the inner disc region is the flattest compared to all other vertical cuts above (see Fig. 5). For the  $R$  band it is almost as flat as the scalelength at a distance of four times the vertical scaleheight. We observe a systematic decrease in scalelength in both bands up to a vertical distance of  $z \approx 15$  arcsec, after which the scalelength starts to rise again for the deeper  $R$ -band image out to the highest vertical cut at 30 arcsec. If this is not an intrinsic feature of NGC 5290 one could argue that the dust may still influence the scalelength determination even in the NIR band. However, quantitatively this seems very unlikely since the amount of dust inferred from the  $K'$ -band flattening should have a more severe effect on the  $R$ -band profile. In the deeper optical image the scalelength increases by 26 per cent between 2 and 4 vertical scaleheight. In the  $R$  band the scalelength beyond the break in the outer disc increases by a factor of 1.5 between 0 and  $2.6h_z$ .

#### NGC 5981

There is a possible dwarf elliptical companion visible at  $(R, z) = (-10$  arcsec,  $+30$  arcsec), and another companion superimposed on the disc at  $(R, z) = (+77$  arcsec,  $-5$  arcsec). For both we have no velocity information, but there is another large elliptical nearby (NGC 5982, 6.2 arcmin away with  $\Delta v \approx 1300$  km s $^{-1}$ ). NGC 5981 is slightly disturbed and asymmetric. The inner disc is  $\approx 1.5^\circ$  tilted against the outer disc and the bulge component is furthermore slightly tilted against the inner disc. This asymmetry is reflected in the radial profiles. The outer slope of a broken exponential fit is significantly shallower on the NW(right) side, where also the contour map is more extended. The single disc fit by Pohlen (2001) yields a vertical scaleheight of  $h_z = 4.2$  arcsec, a radial scalelength of  $h = 24.1$  arcsec and an inclination of  $86.5^\circ$ .

The face-on-equivalent profile of NGC 5981 is well described as a Type II with an inner shallow exponential (of  $h_{\text{in}} = 27.9$  arcsec  $\equiv$

**Table A1.** Exponential disc parameters for all fitted profiles.


Galaxy	Filter	Profile type	$z$ cut (arcsec)	$b_1$ (arcsec)	$b_2$ (arcsec)	$R_{\text{br}}$ (arcsec)	$h_{\text{in}}$ (arcsec)	$h_{\text{out}}$ (arcsec)	$R_{\text{br}}$ ( $h_{\text{in}}$ )	$h_{\text{in}}$ ( $h_{\text{out}}$ )	$\mu_{\text{br}}$ (mag arcsec $^{-2}$ )
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
ESO 380–019	$V$	II	foe	12	117	87.6	27.9	11.5	3.1	2.4	22.9
ESO 380–019	$V$	II	0	12	117	93.7	26.1	8.6	3.6	3.0	27.4
ESO 380–019	$V$	II	6	12	117	86.9	28.8	10.5	3.0	2.7	27.4
ESO 380–019	$V$	II	12	0	117	77.4	30.4	14.2	2.6	2.1	28.1
ESO 380–019	$V$	II	18	10	117	65.4	32.7	21.5	2.0	1.5	29.1
ESO 380–019	$V$	I	23	10	90	–	34.9	–	–	–	–
ESO 404–018	$V$	II	foe	0	92	75.6	34.0	9.6	2.2	3.6	24.6
ESO 404–018	$V$	II	0	0	92	72.4	31.1	9.0	2.3	3.5	28.0
...											

Notes. (1) Principal name, (2) filter, (3) profile type (I: no break, II: downbending break), (4) vertical position  $z$  of the deprojected slices (‘foe’ means the face-on-equivalent profile), (5,6) inner and outer fitting boundaries, (7,8,9) break radius, inner and outer scalelength in units of arcsec, (10) break radius in relation to the inner scalelength, (11) inner scalelength in relation to the outer scalelength, (12) the surface brightness at the break radius (estimated at the crossing point of the two exponential fits). This is a sample: the full version of this table is available in the online version of this article in the Supplementary Materials section.

$2.0 h_{\text{out}}$ ) and an outer steeper exponential ( $h_{\text{out}} = 13.8$  arcsec) with a break at a radial distance of  $R_{\text{br}} \approx 67$  arcsec  $\equiv 2.4 h_{\text{in}}$  and at a surface brightness of  $\mu_{\text{br}} = 24.0 R\text{-mag arcsec}^{-2}$ .

We find an increase in scalelength of  $\approx 10$  per cent between the major axis and about 3.6 times the mean vertical scaleheight (at 15 arcsec) in the inner region. Using only the two quadrants from the dust-free side for averaging (in opposite to all four) this increase

in scalelength for the inner disc grows to  $\approx 25$  per cent. In this case – as expected from the model fits (cf. Fig. 3) – the scalelength on the major axis is less affected by the dust, so smaller, whereas the scalelength at 3.6 vertical scaleheights is almost identical (29.7 arcsec for the mean quadrant versus 29.2 arcsec for the dust-free part). The scalelength beyond the break in the outer disc increases by a factor of 1.4 between 0 and 3.6  $h_z$  in our mean



**Figure B1.** (a) Radial surface brightness profiles and isophote maps for ESO 380–019 V band, ESO 404–018 V band, FGC 2339 R band and IC 4871 V band (from top left to bottom right). See text for further explanations. (b) Radial surface brightness profiles and isophote maps for NGC 522 R band, NGC 1596 R band, NGC 3390 R band and NGC 4179 V band (from top left to bottom right). (c) Radial surface brightness profiles and isophote maps for NGC 5290 R band, NGC 5290  $K'$  band, NGC 5981 R band and UGC 10459 R band (from top left to bottom right).



quadrant. Owing to the described asymmetries in the disc on both sides this galaxy is well suited to study their effects on the de-projected profiles. Fitting only a mean quadrant obtained from the NW(right) or SE(left) side of the galaxy separately we find differences of 5–10 per cent for the scalelengths of the inner region. In the outer disc (beyond the break) the differences are clearly

larger, e.g. 40 per cent on the major axis, due to the obvious asymmetry.

### UGC 10459

This galaxy has only a very small bulge component. There is a possible background galaxy, with unknown distance, visible at




Figure B1 – continued



$(R, z) = (+33 \text{ arcsec}, -9 \text{ arcsec})$ . The major axis profile shows an obvious broken exponential structure with a break radius at  $R \approx \pm 45 \text{ arcsec}$  ( $\mu_R \approx 23.0 \text{ mag arcsec}^{-2}$ ) separating a shallow inner and a steeper outer slope. The single disc fit by Pohlen (2001) yields a vertical scaleheight of  $h_z = 2.0 \text{ arcsec}$ , a radial scalelength of  $h = 13.4 \text{ arcsec}$  and an inclination of  $87.0^\circ$ .

The face-on-equivalent profile of UGC 10459 is well described as a Type II with an inner shallow exponential (of  $h_{\text{in}} =$

$15.6 \text{ arcsec} \equiv 2.7 h_{\text{out}}$ ) and an outer rather steep exponential ( $h_{\text{out}} = 5.7 \text{ arcsec}$ ) with a break at a radial distance of  $R_{\text{br}} \approx 44 \text{ arcsec} \equiv 2.8 h_{\text{in}}$  and at a surface brightness of  $\mu_{\text{br}} = 25.3 R\text{-mag arcsec}^{-2}$ .

We find an increase in scalelength of  $\approx 40$  per cent between the major axis and about 5.6 times the vertical scaleheight (at  $11 \text{ arcsec}$ ) in the inner disc region. The scalelength on the major axis is slightly larger compared to scalelength of the cut at  $2.1 h_z$ , which is well




Figure B1 – continued

explained by the influence of the dust. The scalelength beyond the break in the outer disc increases by a factor of 2.2 between 0 and  $3.6h_z$ .

## APPENDIX B: FIGURES

The following figures [Figs B1(a)–(c)] show radial surface brightness profiles and isophote maps for four galaxies per page (in alphabetical order). For the isophote map (lower panels) each galaxy is rotated to the major axis. The magnitude of the outer contour ( $\mu_{\text{lim}}$ , defined by a  $3\sigma$  criterion of the background) is indicated in each plot. The consecutive contours are equidistant spaced by 0.5 mag. In the plot, the contour lines are drawn with increasing smoothing towards the outer parts. Note, however, that for the actual fitting a constant scale smoothing is used. For the inner contours out to where the noise begins to increase no smoothing was applied. The following two–three contours are smoothed by replacing each pixel by the mean of  $3^2$ -pixel ‘around’ the central pixel. For the outer two contours this smoothing is increased to  $5^2$ -pixel. The upper panels display the major-axis surface brightness profile (top, solid line) and two parallel radial profiles (lower, solid lines) each above and below the major axis. The exact vertical positions ( $z$ ) for the plotted pro-

files are indicated in the upper right-hand corner of the plot. In some cases the best-fitting, sharply truncated, single disc model obtained by Pohlen (2001) is overplotted (dashed line).

## SUPPLEMENTARY MATERIAL

The following supplementary material is available for this article:

**Table A1.** Exponential disc parameters for all fitted profiles.

This material is available as part of the online article from <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1365-2966.2007.11790.x> (This link will take you to the article abstract.)

Please note: Blackwell Publishing is not responsible for the content or functionality of any supplementary materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a  $\text{\TeX/L\AA\TeX}$  file prepared by the author.