

University of Groningen

Reply to comment on "A local realist model for correlations of the singlet state" by M.P. Seevinck and J.-A. Larsson

de Raedt, Hans; Michielsen, K.; Miyashita, S.; Keimpema, K.

Published in:
European Physical Journal B

DOI:
[10.1140/epjb/e2007-00195-2](https://doi.org/10.1140/epjb/e2007-00195-2)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

De Raedt, H., Michielsen, K., Miyashita, S., & Keimpema, K. (2007). Reply to comment on "A local realist model for correlations of the singlet state" by M.P. Seevinck and J.-A. Larsson. *European Physical Journal B*, 58(1), 55-59. DOI: 10.1140/epjb/e2007-00195-2

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Reply to comment on “A local realist model for correlations of the singlet state” by M.P. Seevinck and J.-Å. Larsson

H. De Raedt^{1,a}, K. Michielsen², S. Miyashita³, and K. Keimpema¹

¹ Department of Applied Physics, Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

² EMBD, Vlasakker 21, 2160 Wommelgem, Belgium

³ Department of Physics, Graduate School of Science, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

Received 15 May 2007

Published online 28 July 2007 – © EDP Sciences, Società Italiana di Fisica, Springer-Verlag 2007

Abstract. The general conclusion of Seevinck and Larsson is that our model exploits the so-called coincidence-time loophole and produces sinusoidal (quantum-like) correlations but does not model the singlet state because it does not violate the relevant Bell inequality derived by Larsson and Gill, since in order to obtain the sinusoidal correlations the probability of coincidences in our model goes to zero. In this reply, we refute their arguments that lead to this conclusion and demonstrate that our model can reproduce results of photon and ion-trap experiments with frequencies of coincidences that are not in conflict with the observations.

PACS. 03.65.-w Quantum mechanics – 02.70.-c Computational techniques; simulations

In order to come to their conclusions, Seevinck and Larsson made the following statements [1]:

- De Raedt et al. claim that their model violates the CHSH inequality, a claim that cannot be found in reference [2];
- the CHSH inequality is inappropriate for models that exploit the so-called coincidence-time loophole [3] and the appropriately modified inequality [3] is not violated by the model of De Raedt et al.;
- the model of De Raedt et al. cannot reproduce all experimental realizations of the EPRB experiment;
- De Raedt et al. claim that their model can reproduce the coincidences of recent experimental results, another claim that cannot be found in reference [2];

and put our model in the context of hidden variable models to obtain an expression for the probability of coincidences.

In this reply, we point out once more that in our work [2] we did not rely on Bell's or CHSH's inequality nor on any generalization thereof to come to our conclusion that it is possible to construct an event-based simulation model that satisfies Einstein's criteria of local causality and realism and can reproduce the expectation values of a system of two $S = 1/2$ particles in the singlet state. In our work [2] we did not make any claims about these

inequalities, neither did we make any statement about coincidences in real experiments. We furthermore demonstrate that the calculation by Seevinck and Larsson [1] of the probability of coincidences for our model is simply wrong and we present results for the frequency of coincidences which compare rather well to the values observed in recent experiments.

Before replying in detail to the comments of Seevinck and Larsson [1] on our model [2], we first want to sincerely apologize that we did not make a reference to the model presented in reference [3] which, like our model, uses coincidence in time as a criterion to decide which pairs of detection events are to be considered as stemming from a single two-particle system. Furthermore, to our knowledge, it is the first work to point out how the original Bell inequality changes when using this post-selection procedure [3]. Although in our work [2], we did not rely on Bell's original inequality or on any of its generalizations [4] to come to our conclusions, we should have made reference to reference [3] only because of the fact that the model presented in reference [3] uses the same pair selection criterion as we use in our model [2].

As stated in paragraph 2 of our paper [2], we consider the original EPRB problem, that is the construction of a model that satisfies Einstein's criterion of local causality for each pair of events and reproduces the expectation values of a system of two $S = 1/2$ particles in the singlet

^a e-mail: h.a.de.raedt@rug.nl

state. Whether or not this model leads to a violation of some inequality is of secondary interest.

We do not share the point of view of Seevinck and Larsson [1] that a system is in the “singlet state” if and only if some correlation violates a certain bound and if the probability of coincidences does not go to zero. This viewpoint does not make much sense if we apply it to the ground state of a hydrogen molecule (a spin singlet), for instance. The singlet state is a concept of quantum theory. Unlike Seevinck and Larsson suggest in their conclusion, the singlet is not defined with any reference to coincidence counts or (generalized) Bell inequalities.

According to quantum theory, the singlet state is completely characterized by the single-particle and two-particle expectation values $E_1(\mathbf{a}_1) = E_2(\mathbf{a}_2) = 0$ and $E(\mathbf{a}_1, \mathbf{a}_2) = -\mathbf{a}_1 \cdot \mathbf{a}_2$, respectively. Because quantum theory has nothing to say about single events [5], it does not give us a recipe to compute $E_1(\mathbf{a}_1)$, $E_2(\mathbf{a}_2)$ and $E(\mathbf{a}_1, \mathbf{a}_2)$ from the record of single events in laboratory experiments or theoretical models.

In the case of EPRB laboratory experiments with photons [6], coincidence in time seems to be a good criterion to identify particle pairs based on the time-tag data of single particle events, since by using this criterion results comparable to those expected from quantum theory can be obtained. Therefore we use the same criterion in our model [2]. However, other criteria to decide which single particles belong to a single two-particle system are not excluded. The criterion depends on the experimental setup but quantum theory does not give any guidance to define a criterion. Having made a definite choice for this criterion, we can compute the single-particle and two-particle expectation values from the record of single events in laboratory or computer experiments and compare the outcome with the results from quantum theory. If and only if we find $E_1(\mathbf{a}_1) = E_2(\mathbf{a}_2) = 0$ and $E(\mathbf{a}_1, \mathbf{a}_2) = -\mathbf{a}_1 \cdot \mathbf{a}_2$, we may say that we found expectation values that correspond to those of a singlet state. No other criteria, like violating an inequality or computing a probability of coincidences for example, are required to come to this conclusion. Note that we cannot say anything more than that the expectation values correspond to those of a singlet state. For example, we cannot make statements such as the source produces singlets, since the results for the expectation values do not only depend on the characteristics of the source and the detection elements but also on the measurement (post-processing) process.

It is self-evident that our model is too simple to describe, in every detail, all conceivable experimental realizations of the EPRB thought experiment but it is the first model that satisfies Einstein’s conditions of local causality and realism and that exactly reproduces the single-particle and two-particle expectation values of the singlet state. In this respect, it may be viewed as the first realization of the EPRB thought experiment (as defined by EPRB), since none of the laboratory experiments of the EPRB experiment have shown results for the single- and two-particle expectation values that compare so well with those of quantum theory. In those experiments, conclu-

sions are usually drawn based on the value for S_{max} only. Moreover, to draw conclusions about local realist modelling of expectation values that agree with those of a singlet state, finding one such model is sufficient. Whether this model then fails to describe all possible laboratory realizations of the EPRB thought experiment becomes irrelevant and it remains to be seen if these laboratory experiments produce data that completely characterize a singlet state, a requirement of the EPRB thought experiment.

Seevinck and Larsson state “we will put the model used by De Raedt et al. in its proper context” [1]. However, they failed to do so in any respect. In spite of the fact that in our paper, we repeatedly stress that in formulating our model we do not rely on concepts of probability theory, they seem to ignore our statements. This is most evident by their statement that “the local hidden variable ... is denoted by $S_{n,i}$ ”. Since, according to Larsson a hidden-variable model is really a probabilistic model [7] and since our model is purely ontological, the concept of a hidden variable cannot be applied to our model as such.

In contrast to the (repeated) statement made in reference [1], we did not claim that the CHSH inequality (see Eq. (2) in Ref. [1]) is valid for our model. There is no such statement in our paper. In our paper, we studied the values of S_{max} as a function of the time window W relative to the time-tag resolution τ and this for several values of the model parameter d . We compared the results with $S_{max} = 2\sqrt{2}$, the quantum theoretical result for the singlet state and also the maximum value for S_{max} that can be obtained for any choice of the quantum state. Although we find that for some model parameters $2 < S_{max} < 4$ we did not claim that our model violates the CHSH inequality, as stated by Seevinck and Larsson [1]. In fact, using elementary algebra it follows immediately from equations (3) and (5) of reference [2] that $|E(\mathbf{a}_1, \mathbf{a}_2)| \leq 1$ and that

$$|E(\mathbf{a}, \mathbf{c}) - E(\mathbf{a}, \mathbf{d}) + E(\mathbf{b}, \mathbf{c}) + E(\mathbf{b}, \mathbf{d})| \leq 4, \quad (1)$$

for the data generated by our computer model. Without any further constraints on the algorithm that generates the data $\{\mathcal{Y}_1, \mathcal{Y}_2\}$ (see Eq. (1) in Ref. [2]), the upperbound (4) in equation (1) cannot be improved. In our paper [2], we use expression equation (5) (see Ref. [2]) to discuss the nature of the quantum state, but attach no meaning to the violation of some bound by our simulation data.

Not surprisingly, also the statement “they furthermore claim that the maximal quantum violation is ...”, [1] is wrong. We did not make any reference to the CHSH inequality in our paper. Looking at Figure 3 of our paper [2], Seevinck and Larsson should have noted that for $d > 3$, our model can produce correlations that are (much) stronger than those of quantum theory (which in view of equation (1) is not a surprise). In fact, the correct statement (see Ref. [2]) is that our model can exhibit correlations that are **stronger** than those of quantum theory of two $S = 1/2$ particles.

As we mentioned before, we agree with Seevinck and Larsson that when time-coincidence is used to decide which pairs of detection events are to be considered as

stemming from a single two-particle system and if one would like to consider a generalized Bell inequality, the relevant inequality to consider would be equation (4) in reference [1] and not the CHSH inequality. In this modified inequality γ is the infimum of the probability of coincidence [3]. Seevinck and Larsson compute γ for our model. They conclude that our model does not violate equation (4) in reference [1]. Moreover, although in the paper on the photon experiment [6] or in the paper on the ion-trap experiment [8], there is no information about the minimum frequency of coincidences (the minimum is required for the application of equation (4) in Ref. [1]), Seevinck and Larsson refer to these papers when they cite the values of $\gamma = 0.05$ and $\gamma = 1$ and then state that our model cannot reproduce the frequencies of coincidences that agree with those found in these two experiments. First of all, the statement that “De Raedt et al. claim that their model can reproduce the coincidences of recent experimental results” [1] is simply wrong: There is no such claim in our paper [2]. Second, it is logically inconsistent to draw conclusions based on the comparison with the ion-trap experiment for which $\gamma = 1$ [1] and third, the calculation of γ [1] for our model is wrong, two statements which we will prove in what follows.

Let us formalize the statements in reference [1] as propositions (denoted by A, B, ...):

- A. The probabilistic, hidden variable model using the time window to define coincidences [3] yields the inequality given in equation (4) of reference [1] (see also Ref. [3]). The upperbound of this inequality is given by $6/\gamma - 4$, where γ denotes the infimum of the probability of coincidences over all possible settings $\mathbf{a}_1, \mathbf{a}_2$ of the detectors.
- B. In the EPRB experiment with ions [8], $\gamma = 1$ [1].
- C. The probabilistic, hidden variable model of reference [3] applies to the ion-trap experiment [8] and hence the experimental data should satisfy the inequality given in equation (4) of reference [1]. Note that the second part of this statement implicitly follows from proposition B.
- D. The ion-trap experiment yields $S_{max} \approx 2.25$ [8]. Strictly speaking, this statement is not made in reference [1], but it is an experimental fact and as such cannot be denied.

Let us now apply the rules of elementary logic.

If $\gamma = 1$ [1], the ion-trap experiment not only violates the original Bell inequality but as $6/\gamma - 4 = 2$, it also violates the inequality given in equation (4) of reference [1]. Thus, we have

$$A \wedge B \wedge C \wedge D \Rightarrow \overline{C}, \quad (2)$$

where \wedge , \Rightarrow and $\overline{}$ denote the logical “and” operation, logical implication, and logical negation, respectively. Clearly, equation (2) expresses a logic contradiction. If we assume that propositions A and D are true (as we do), then we must conclude that B or C or both B and C are false. In any case, the argument used by Seevinck and Larsson leads to a logical contradiction, independent of what we wrote in our paper [2].

We should not exclude the possibility (that is, we might accept proposition \overline{C}) that the model of reference [3] or ours [2] is too simple to describe the ion-trap experiment [8]. This experiment uses a detection pulse during which the bright state of an ion scatters many photons (64 on average) [8]. This process may not be sampling “single-events” but is more likely to probe the ensemble average that is given by quantum theory (although the number of samples, ≈ 64 , is not large).

By trying to put our work in the context of “hidden variable theories”, Seevinck and Larsson also made mistakes in elementary algebra. Seevinck and Larsson assume that the probability of coincidences is given by the denominator of equation (6) in reference [2] (see Appendix A of Ref. [1]), from which they derive an expression for the probability of coincidences γ (see Eq. (8) in Ref. [1]). However, Seevinck and Larsson apparently overlooked the fact that in going from equation (3) to equation (6) (see Ref. [2]), we take the limit $W/T_0 = \tau/T_0 \rightarrow 0$ and let the number of events N in both the numerator and denominator go to infinity. Although the ratio remains finite, which is obvious in the case $\mathbf{a}_1 = \mathbf{a}_2$ ($x_1 x_2 = -1$) where it is equal to minus one, the limit of the denominator may not exist and in fact, it diverges if $\mathbf{a}_1 = \mathbf{a}_2$. This is not a problem of our model: This divergence merely signals that one has to be careful in taking the limits. The mathematical derivation in Appendix A of reference [1] is simply incorrect.

Nevertheless, Seevinck and Larsson raise an interesting question about the role of the frequency (not probability) of coincidences in our model. For nonzero time-tag resolution τ and time window $W \geq \tau$, the frequency of coincidences in our simulation model is given by

$$\Gamma = \frac{1}{N} \sum_{n=1}^N \Theta(W - |t_{n,1} - t_{n,2}|), \quad (3)$$

a well-defined quantity in our simulation model that is easy to compute numerically. Notice that Γ is a function of \mathbf{a}_1 and \mathbf{a}_2 and that $0 \leq \Gamma \leq 1$. Assuming that the results we obtain by using pseudo-random numbers can be described by a probabilistic model, we expect that $\gamma = \min_{\mathbf{a}_1, \mathbf{a}_2} \Gamma$ with probability one if N is sufficiently large. With these additional assumptions, not only the inequality equation (1) holds but also the inequality given by equation (4) of reference [1] holds.

In our model, there are four free parameters, namely the time window W/τ , the maximum time delay T_0/τ , the time-tag exponent d and the number of events N . For $d = 3$, $N \rightarrow \infty$ and in the limit $W/T_0 = \tau/T_0 \rightarrow 0$, our model reproduces exactly, the expression for the two-particle expectation value of a quantum system in the singlet state [2].

For $d = 3$, $W = \tau$, $T_0/\tau = 1000$ and $N = 10^6$ (the results reported in this paper do not change if $N > 5 \times 10^5$), we find that $\min_{\mathbf{a}_1, \mathbf{a}_2} \Gamma \approx 1.27W/T_0$, in concert with the rigorous result (for $d = 3$ and $W = \tau$)

$$\min_{\mathbf{a}_1, \mathbf{a}_2} \left\{ \lim_{W/T_0 \rightarrow 0} \Gamma \right\} = \frac{4}{\pi} \frac{W}{T_0} \approx 1.27 \frac{W}{T_0}. \quad (4)$$




Fig. 1. (Color online) The frequency of coincidences Γ as a function of $\theta = \arccos(\mathbf{a}_1 \cdot \mathbf{a}_2)$ for parameters τ/T_0 , W/T_0 and d chosen such (see text) that the simulation model reproduces the result for a singlet state, $S_{max} = 2.83$ (solid line, red), and the values of $S_{max} = 2.25$ (dashed line, black) and $S_{max} = 2.73$ (dotted line, blue), as obtained from experiments with ions [8] and with photons [6], respectively.

Thus, in the regime $W/T_0 = \tau/T_0 \rightarrow 0$, we find that the minimum frequency of coincidences is proportional to the width of the time bins, as it should be.

Next, we consider the possibility of fitting the results of our model to the experimental data of an EPRB experiment with photons [6] and ions [8]. According to Seevinck and Larsson [1], our model cannot reproduce these experimental data. For simplicity, we set $T_0/\tau = 1000$, $d = 3$ and take $N = 10^6$. Then, there is one free parameter left, namely the (dimensionless) time-window $W/\tau \geq 1$. The fitting procedure consists of changing W/τ such that the value of $S_{max} = \max_{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}} S(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d})$ agrees with values cited in references [6, 8].

In Figure 1, we present our results for the frequency of coincidences for the values of $W/\tau = 285$, $W/\tau = 16$, and $W/\tau = 1$, for which our model yields $S_{max} = 2.25$ [8], $S_{max} = 2.73$ [6], and $S_{max} = 2.83$ (singlet state), respectively.

From Figure 1, we see that in order to reproduce the ion-trap result [8], the frequency of coincidences $\Gamma \geq 0.52$ is quite large. It is important to recognize that with four free parameters at our disposal, it is easy to reproduce almost any number for S_{max} , as long as it is between zero and four. For instance, we find the same value of $S_{max} = 2.25$ for $W = \tau$ and $T_0/\tau = 1.025$ but then $\Gamma \geq 0.87$. In any case, these results refute the statement in the Comment that our model cannot reproduce the experimental result $S_{max} = 2.25$ of the ion-trap experiment [8] with a nonzero value of Γ . Fitting our model (for $d = 3$ and $T_0/\tau = 1000$) to $S_{max} = 2.73$ and $S_{max} = 2.83$ yields $\Gamma > 0.0377$ and $\Gamma > 0.00127$, respectively.

An analysis of experimental data for an EPRB experiment with photons [9] yields $\Gamma \approx 0.01$ (the value of $\gamma \approx 0.05$ cited in reference [1] is the total frequency of coincidences, that is the sum over four experiments, and not the infimum over all possible experiments, as required




Fig. 2. (Color online) Simulation results of the two-particle correlation $E(\mathbf{a}_1, \mathbf{a}_2)$ as a function of $\theta = \arccos(\mathbf{a}_1 \cdot \mathbf{a}_2)$ for the model parameters that yield $S_{max} = 2.25$ (squares, black), $S_{max} = 2.73$ (stars, blue), and $S_{max} = 2.83$ (bullets, red), respectively. The solid line (black) is the result ($E(\mathbf{a}_1, \mathbf{a}_2) = -\mathbf{a}_1 \cdot \mathbf{a}_2$) for the singlet state.

for the application of the inequality given in Eq. (4) of Ref. [1]). Thus, for the same value of S_{max} , our model yields a value of Γ that is larger ($\Gamma = 0.0377$) than the value that can be extracted from experimental data for an EPRB experiment with photons [9].

As our model is flexible enough to yield for S_{max} any number between zero and four with reasonable values of the model parameters, it is of interest to study how the correlation $E(\mathbf{a}_1, \mathbf{a}_2)$ deviates from the result $E(\mathbf{a}_1, \mathbf{a}_2) = -\mathbf{a}_1 \cdot \mathbf{a}_2$ of a system in the singlet state as we fit the values of S_{max} to the experimental results.

In Figure 2, we show the simulation results for the same three cases $S_{max} = 2.25, 2.73, 2.83$. From Figure 2, it is clear that the simulation data that yields $S_{max} = 2.25, 2.73$ cannot be described by a single sinusoidal function, but for $S_{max} \geq 2.73$ the deviations from a single sinusoidal are small and it remains to be seen if experiments can resolve such small differences.

As is evident from Figure 3 in reference [2], for $d > 3$ our model yields the value for the singlet state $S_{max} = 2\sqrt{2}$ without having to consider the limit $W/T_0 = \tau/T_0 \rightarrow 0$. Thus, in order for an experiment and a model of the type considered in our paper to reproduce **all** the features of a quantum system of two $S = 1/2$ particles in the singlet state, it is not sufficient to show that it can yield $S_{max} = 2\sqrt{2}$ for some choice of the parameters. As mentioned before, the singlet state is completely characterized by the single and two-particle expectation values. Hence, in order to make a comparison with the singlet state, it is necessary to measure or compute these two quantities.

Finally, the statement in our paper [2] that “our work suggests that it is possible to construct event-based simulation models that satisfies Einstein’s criteria of local causality and realism and can reproduce the expectation values calculated by quantum theory [10–14]” should not be taken out of the context as Seevinck and Larsson did by omitting the references. In fact, what we have shown in

the work that we refer to is that it is possible to perform an event-based simulation, satisfying Einstein’s criteria of local causality, of a universal quantum computer [13], which according to the theory of quantum computation should suffice to simulate any quantum system [15].

In conclusion, the purpose of our work is to construct an event-based simulation model, satisfying Einstein’s criteria of local causality and realism, that produces the quantum correlations of the singlet state [2]. As we have shown in reference [2] we succeeded, to our knowledge for the first time, in constructing such a model. Our conclusion that we find results that are indistinguishable from those of a singlet state is based on the fact that the calculated single-particle averages and two-particle correlation function agree with the well-known results $E_1(\mathbf{a}_1) = E_2(\mathbf{a}_2) = 0$ and $E(\mathbf{a}_1, \mathbf{a}_2) = -\mathbf{a}_1 \cdot \mathbf{a}_2$ for a system of two $S = 1/2$ particles in the singlet state.

We have also demonstrated in reference [2] and in this reply that knowing S_{max} , a quantity derived from the two-particle correlation function, does not suffice to draw any conclusion about the observation of a singlet(-like) state. We also demonstrate in this reply, that our model can not only produce the results from quantum theory for a system of two $S = 1/2$ particles in the singlet state but that it can also be applied to EPRB laboratory experiments with photons and ions and give results for S_{max} and the frequency of coincidences that are comparable to the values extracted from these experiments.

References

1. M.P. Seevinck, J.Å. Larsson, Eur. Phys. J. B **58**, 51 (2007)
2. K. De Raedt, K. Keimpema, H. De Raedt, K. Michielsen, S. Miyashita, Eur. Phys. J. B **53**, 139 (2006)
3. J.A. Larsson, R.D. Gill, Europhys. Lett. **67**, 707 (2004)
4. J.S. Bell, *Speakable and unspeakable in quantum mechanics* (Cambridge University Press, Cambridge, 1993)
5. D. Home, *Conceptual Foundations of Quantum Physics* (Plenum Press, New York, 1997)
6. G. Weihs, T. Jennewein, C. Simon, H. Weinfurter, A. Zeilinger, Phys. Rev. Lett. **81**, 5039 (1998)
7. J.A. Larsson, Phys. Rev. A **57**, 3304 (1998)
8. M.A. Rowe, D. Kielpinski, V. Meyer, C.A. Sackett, W.M. Itano, C. Monroe, D.J. Wineland, Nature **401**, 791 (2001)
9. <http://www.quantum.at/research/photonentanglement/bellexp/data.html>
10. K. De Raedt, H. De Raedt, K. Michielsen, Comp. Phys. Comm. **171**, 19 (2005)
11. H. De Raedt, K. De Raedt, K. Michielsen, J. Phys. Soc. Jpn Suppl. **76**, 16 (2005)
12. H. De Raedt, K. De Raedt, K. Michielsen, Europhys. Lett. **69**, 861 (2005)
13. K. Michielsen, K. De Raedt, H. De Raedt, J. Comput. Theor. Nanosci. **2**, 227 (2005)
14. H. De Raedt, K. De Raedt, K. Michielsen, S. Miyashita, Comp. Phys. Comm. **174**, 803 (2006)
15. M. Nielsen, I. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000)