



## **University of Groningen**

## Pex19p Contributes to Peroxisome Inheritance in the Association of Peroxisomes to Myo2p

Otzen, Marleen; Rucktaeschel, Robert; Thoms, Sven; Ernmrich, Kerstin; Krikken, Arjen; Erdmann, Ralf; van der Klei, Ida; Emmrich, Kerstin; Krikken, A.M.

Published in: Traffic

DOI:

10.1111/j.1600-0854.2012.01364.x

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version Publisher's PDF, also known as Version of record

Publication date:

Link to publication in University of Groningen/UMCG research database

Citation for published version (APA):

Otzen, M., Rucktaeschel, R., Thoms, S., Ernmrich, K., Krikken, A. M., Erdmann, R., ... Krikken, A. M. (2012). Pex19p Contributes to Peroxisome Inheritance in the Association of Peroxisomes to Myo2p. Traffic, 13(7), 947-959. DOI: 10.1111/j.1600-0854.2012.01364.x

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): http://www.rug.nl/research/portal. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Download date: 11-02-2018




Figure S3: Protein levels in cells used for yeast two-hybrid studies

Western blot analyses of cells producing different mutant forms of the cargo binding domain of Myo2p. The Myo2p cargo binding domain fused to the binding domain of Gal4p revealed that levels of the mutant proteins were similar to the WT control. Pyruvate carboxylase (Pyc1p) was used as loading control.