University of Groningen

Controlling the Dimensionality and Structure of Supramolecular Porphyrin Assemblies by their Functional Substituents

Iacovita, Cristian; Fesser, Petra; Vijayaraghavan, Saranyan; Enache, Mihaela; Stöhr, Meike; Diederich, Francois; Jung, Thomas A.

Link to publication in University of Groningen/UMCG research database

CHEMISTRY
A EUROPEAN JOURNAL

Supporting Information

Controlling the Dimensionality and Structure of Supramolecular Porphyrin Assemblies by their Functional Substituents: Dimers, Chains, and Close-Packed 2D Assemblies

Cristian Iacovita,*[a] Petra Fesser,[b] Saranyan Vijayaraghavan,[a] Mihaela Enache,[c] Meike Stöhr,*[c] François Diederich,*[b] and Thomas A. Jung*[a, d]

chem_201201037_sm_miscellaneous_information.pdf

Fig. 1S1. a) Histogram of the inter-chains separations of the zinc(II)porphyrin at 0.7 ML indicating preferred distances of 4, 6, 8, 10, 12, 14, 16 and 18nm ±0.15nm. b) STM topographic image of two parallel chains. The double arrows show the inter-chains separations defined as the distance between the partially positively charged DMA units.