

University of Groningen

## On the Mechanism of Peptidoglycan Binding and Cleavage by the endo-Specific Lytic Transglycosylase MltE from *Escherichia coli*

Fibriansah, Guntur; Gliubich, Francesca I.; Thunnissen, Andy

*Published in:*  
Biochemistry

*DOI:*  
[10.1021/bi300900t](https://doi.org/10.1021/bi300900t)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2012

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Fibriansah, G., Gliubich, F. I., & Thunnissen, A-M. W. H. (2012). On the Mechanism of Peptidoglycan Binding and Cleavage by the endo-Specific Lytic Transglycosylase MltE from *Escherichia coli*. *Biochemistry*, 51(45), 9164-9177. DOI: 10.1021/bi300900t

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*


# Supplementary Info

Fibriansah *et al.* (2012) On the mechanism of PG binding and cleavage by the endo-specific lytic transglycosylase MltE from *Escherichia coli*

**Table S1.** Se-MAD Data Collection and Phasing Statistics

<b>Data collection</b>	<b>SeMet-sMltE</b>		
	inflection	peak	remote
Beamline		BM14 (ESRF)	
Wavelength (Å)	0.9788	0.9790	0.8855
Space group		$P2_12_12_1$	
Unit cell dimensions		a = 77.6 Å b = 94.6 Å c = 160.8 Å	
Max. resolution (Å)	2.75	2.75	2.75
No. of measured reflections	96998	95894	96424
No. of unique reflections	28972	28898	28853
Completeness (%)	92.8	92.3	92.2
R <sub>sym</sub>	0.055	0.061	0.065
Mean I/s	9.4	8.3	8.0
<b>MAD phasing</b>	figures of merit (centric/acentric): 0.53/0.57		


**Figure S1**




**Figure S1.** Peptidoglycanolytic and chitinolytic assays. (A). Turbidimetric assay of PG solubilization by sMltE. The decrease in turbidity (measured at 600 nm) of insoluble *M. luteus* cell wall (in 50 mM MES pH 6.0 and 200 mM NaCl) was monitored with time after addition of (♦) sMltE-WT, (■) sMltE-E64Q and (▲) buffer only (control). The turbidity decrease data is presented as the percentage decrease from the initial value (using an interpolated OD<sub>600</sub> at t = 0 min), and calculated with the averaged data of triplicate experiments. (B) The chitinolytic activity of MltE was assayed by analysing the products resulting from a reaction with chitohexaose as substrate using TLC. Hen egg-white lysozyme was used as positive control. M: sMltE; L: lysozyme; G1: GlcNAc; G2: (GlcNAc)<sub>2</sub>; G3: (GlcNAc)<sub>3</sub>; G4: (GlcNAc)<sub>4</sub>; G6: (GlcNAc)<sub>6</sub>.

Supplementary Figure S2

A




B




**Figure S2.** Comparison of sMltE with the LT-domains of Slt70 and sMltB. (A) Overall comparison of the C $\alpha$  backbone structures (ribbon representation) of the three proteins. Unique segments in sMltE, Slt70-Cdom (LT domain of Slt70) and sMltB-core (LT domain of MltB) with no equivalents in the other proteins are colored green, yellow and cyan, respectively. The unique  $\beta$ -domain in MltB is colored pink. Segments in sMltE that have an equivalent in either one or both of the other LTs are colored dark-grey and light-grey, respectively. Segments in Slt70-Cdom and sMltB-core that have an equivalent in sMltE are colored dark-grey. Residues that are identical in all three proteins are shown in stick representation. (B) Structure-based sequence alignment of the three *E. coli* proteins. The amino acid residue numbering and secondary structure labeling is based on MltE. Common core residues are enclosed in boxes. Residues that are identical in two or three proteins are shown as white letters on an orange or a red background, respectively. The catalytic acid/base is indicated with an asterisk. Spheres below the sMltB sequence indicate the residues in sMltE with a role in saccharide binding.

### Supplementary Figure S3




**Figure S3** Structural basis for the exo- or endo-specific cleavage specificity of Slit70, MltB and MltE. The overall structures of the three proteins are shown in ribbon representation with the LT domains colored in green and the non-LT domains in pink. A bound glycan chain is shown schematically as a solid cylinder (colored with a red to blue gradient), with a terminal 1,6-anhydroMurNAc residue represented by a blue ball. The U and linker domains in Slit70 and the exo-loop in sMltB present a steric obstruction that limit the accessibility of the PG-binding groove, explaining their exo-lytic activity. The absence of such an obstruction in sMltE explains its endo-lytic activity.

### Supplementary Figure S4



**Figure S4.** The 2F<sub>o</sub>-F<sub>c</sub> electron density maps for the bound ligands are shown, contoured at 1σ. (A) Bulgecin A and the murodipeptide GlcNAc-MurNAc-L-Ala-D-Glu in protein molecule A of the asymmetric unit of sMltE/bulgA•GM-Ala-Glu. (B-C) Chitopentaose bound to protein molecule A in the 1.9 Å and 2.4 Å sMltE-E64Q/G5 structure, respectively.

Supplementary Figure S5



**Figure S5.** Stereo diagram showing an overlay of the apo structure of sMltE (cyan) with the structures of sMltE-E64Q/G5 (purple) and sMltE/bulgA•GM-Ala-Glu (yellow).