

University of Groningen

E0 Conversion in ^{238}U in Heavy Ion Collisions at Coulomb Barrier

DEBOWSKI, M; BALANDA, A; DEBOER, FWN; Bokemeyer, H.; ELZE, TW; Gerl, J; Hoogduin, H.; Van Klinken, J; SALABURA, P; Wollersheim, H. J.

Published in:
Acta Physica Polonica B

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1993

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

DEBOWSKI, M., BALANDA, A., DEBOER, FWN., Bokemeyer, H., ELZE, TW., Gerl, J., ... Xie, H. (1993). E0 Conversion in ^{238}U in Heavy Ion Collisions at Coulomb Barrier. Acta Physica Polonica B, 24(2), 425-427.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

E0 Conversion in ^{238}U

in Heavy ion Collisions at Coulomb Barrier

M. Dębowski¹, A. Balanda¹, F.W.N. de Boer², H. Bokemeyer³,
Th. W. Elze², J. Gerl³, H. Hoogduin⁴, J. van Klinken⁴, P. Salabura¹,
H.J. Wollersheim³, H. Xie³

¹ Jagellonian University, Kraków

² Institut für Kernphysik, Universität Frankfurt

³ Kernfysisch Versneller Instituut, Groningen

⁴ Gesellschaft für Schwerionenforschung, Darmstadt

We propose a Coulomb excitation study for E0 conversion in actinide nuclei. We expect the large amount of conversion electrons in transitions from $K = 0$ states with $\Delta I = 0$. By measurement of those electrons we can extend our knowledge about ^{238}U nuclear structure. Using standard γ -spectroscopy methods we can observe the ground-state rotational band up to spin 30^+ and the octupole band to spin 23^- , other collective bands are hardly seen^{1,2}. In 1985 Venema et al.³ measured excited levels in ^{232}U , ^{234}U and ^{232}Th in $^{232}\text{Th}(\alpha, xn)$ reactions. They used two mini-orange detectors. The strong presence of E0 conversions in $\Delta I = 0$ transitions has been observed (fig.1).




Fig.1 Prompt conversion electrons from $^{232}\text{Th}(\alpha, xn)$ reactions.

In nuclei, transitions between states with equal spin, parity and K -quantum numbers proceed predominantly through E2, M1 and E0 radiation. The contribution of E0 transition increases with Z . For instance, for excitation energy $0.5 - 1.0\text{MeV}$, in nucleus with $Z \sim 80$, E0 transition probability increases as $\sim Z^{12}$, in contrast with E2 $\sim Z^{1.6}$ or M1 independent on Z . Also conversion coefficients for excitation energy $\sim 1.0\text{MeV}$ for pure E2 transition is about three orders of magnitude smaller, than those coefficients for E2 and admixed E0 transitions. For example, $\alpha_{exp}(E0+E2)$ val-

ues for $\Delta I = 0$ transitions depopulating excited levels of two low lying $K = 0$ bands in ^{238}U are $6.7 \pm 3.0^{7,0}$ and 4.4 ± 1.2 respectively⁴. Same transition in ^{232}Th has α_{exp} value of 17.0 ± 6.0^5 . These coefficients are incredibly high, one order of magnitude larger, than expected from nuclear structure of actinide nuclei ($\beta \sim 0.25$). If the conversion coefficients are spin independent or increase with spin, the main decay mode of excited levels of $K = 0$ bands is expected to be interband-conversion electrons more, than inband or interband γ -rays. Also $\Delta I = 0$ transitions from γ - to ground-state band have the E0 conversion contribution due to mixing with β -bands.

Up to now, we know two low lying $K = 0$ rotational bands with band heads 927keV and 995keV , up to spin 4^+ . The third $K = 0$ band has been found with band head $1482\text{keV}^{4,6}$ (see fig.2). Decay of that level will occur mainly through conversion electrons, but it is also possible to decay through Internal Pair Conversion. Although the branching ratio for this process is hundred times smaller, than for electron conversion, it is very interesting to measure the contribution of IPC to background in puzzling e^+e^- peaks^{7,8}.




Fig.2 Decay scheme of ^{238}U .

We want to measure conversion electrons in very heavy systems under Coulomb barrier (e.g. $^{238}\text{U} + ^{181}\text{Ta}$). In such experiments there is very high background of γ -rays and much higher, than in lighter systems δ -radiation ($\sigma_{\delta\text{-electrons}} \sim Z^2$). To suppress the background, we apply mini-orange energy filters. Doppler-broadening reduction, we can obtain moving our electron detectors with mini-oranges away from the target to reduce opening angle. For the opening angle of 1% of 4π , for 1MeV electrons Doppler-broadening is $\Delta E_D = 12\text{keV}$ to compare with $\Delta E_D = 140\text{keV}$ for 4π opening angle. The experimental setup looks as on fig.3.




Fig.3 Experimental setup.

References

- [1] E. Grosse et al., *Physica Scripta* **24** (1981) 337.
- [2] H. Ower et al., *Nucl. Phys.* **A388** (1982) 421.
- [3] W.Z. Venema et al., *Phys. Lett.* **156B** (1985) 163.
- [4] E.N. Shurshikov, *Nucl. Data Sheets* **53** (1988) 601.
- [5] M.R. Schmorak, *Nucl. Data Sheets* **36** (1982) 367.
- [6] Th.W. Elze and J.R. Huizinga, *Nucl. Phys.* **A187** (1972) 545.
- [7] P. Salabura et al., *Phys. Lett.* **B245** (1990) 160.
- [8] W. König et al., *Phys. Lett.* **B218** (1989) 12.