

University of Groningen

The role of neutral hydrogen in the life of galaxies and AGN

Gereb, Katinka

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gereb, K. (2014). The role of neutral hydrogen in the life of galaxies and AGN: A spectral stacking analysis [S.l.]: [S.n.]

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



rijksuniversiteit
groningen

The Role of Neutral Hydrogen in the Life of Galaxies and AGN

A Spectral Stacking Analysis

Proefschrift

ter verkrijging van de graad van doctor aan de
Rijksuniversiteit Groningen
op gezag van de
rector magnificus prof. dr. E. Sterken
en volgens besluit van het College voor Promoties.

De openbare verdediging zal plaatsvinden op
vrijdag 26 september 2014 om 14:30 uur

door

Katinka Geréb

geboren op 4 juli 1986
Sangeorgiu de Padure, Romania

Promotor

Prof. dr. R. Morganti

Copromotor

Prof. dr. T. Oosterloo

Beoordelingscommissie

Prof. dr. J.M. van der Hulst

Prof. dr. J. van Gorkom

Prof. dr. E. Sadler

ISBN 978-90-367-7268-6

ISBN 978-90-367-7267-9 (electronic version)

I am prepared to go anywhere, provided it be forward.
- *David Livingstone*

Cover:

Top left: Andromeda galaxy (M31). Credit: Galaxy Evolution Explorer, NASA/JPL-Caltech.

Top right: Hercules A radio galaxy. Credit: NASA, ESA, S. Baum and C. O'Dea (RIT), R. Perley and W. Cotton (NRAO/AUI/NSF), and the Hubble Heritage Team (STScI/AURA).

The background image displays the Westerbork Synthesis Radio Telescope. Credit: AS-TRON Nico Vermaas - Harm Jan Stiepel.

Contents

1	Introduction	1
1.1	A short history of HI observations	1
1.2	HI gas properties of galaxies	3
1.3	Stacking of HI spectra	4
1.4	Gas and the evolution of radio AGN	6
1.5	Questions addressed in this thesis	10
2	Gas and galaxy properties from a stacking experiment	17
2.1	Introduction	18
2.2	The data: piggyback from the continuum observations	19
2.3	Characteristics of the galaxies in the redshift range $0.06 < z < 0.09$	20
2.4	HI stacking	23
2.5	Results	25
2.5.1	HI content and color separation	25
2.5.2	The connection between cold and ionized gas in red galaxies	26
2.5.3	HI and weak radio sources	27
2.5.4	HI and 24 μm emission in LINERs	30
2.6	Summary and Conclusions	30
2.7	Acknowledgements	31
3	The global cold gas content up to $z = 0.12$	35
3.1	Introduction	36
3.2	Observations and sample selection	37
3.3	Data reduction and HI stacking	38
3.4	Characteristics of the galaxies in the selected sample	38
3.5	Results	42
3.5.1	Stacking in color	42
3.5.2	The HI properties of LINERs and optical AGN	44
3.5.3	AGN and SF properties of the radio population	46
3.5.4	The global SFR and SFE up to $z = 0.12$	47
3.6	Discussion and summary	50
3.7	Acknowledgements	51

4	HI absorption stacking of radio galaxies	57
4.1	Introduction	58
4.2	Sample selection and observations	59
4.2.1	Characterization of the AGN sample	60
4.3	Results	62
4.3.1	HI detection	62
4.3.2	HI stacking: the HI distribution of AGN at low optical depth	63
4.3.3	HI and radio morphology	67
4.3.4	The nature of the HI absorbing systems	69
4.4	Conclusions and implications for the future	70
4.5	Acknowledgements	72
5	The HI absorption ‘Zoo’	77
5.1	Introduction	78
5.2	Description of the sample and observations	79
5.3	Results	80
5.3.1	Fitting complex HI absorption profiles with the busy function	83
5.3.2	Characterization of the profiles with BF parameters	87
5.4	The nature of HI absorption in flux-selected radio galaxies	91
5.4.1	Are powerful AGN interacting with their ambient gaseous medium?	92
5.4.2	Fraction and time-scale of candidate HI outflows	94
5.4.3	Gas rich mergers	96
5.5	The HI properties of compact and extended sources	97
5.6	Summary	99
5.7	Acknowledgements	100
5.8	Appendix	101
5.8.1	Notes on the individual detections	101
5.8.2	Summary table of non-detections	106
6	HI, radio continuum, and optical properties of AGN	117
6.1	Abstract	117
6.2	Introduction	117
6.3	Sample selection	119
6.4	Observations and data reduction	122
6.5	Results	122
6.5.1	SDSS6: 4C 29.30	126
6.5.2	SDSS8	126
6.5.3	SDSS4	127
6.5.4	Galaxies detected in HI emission	127
6.6	Rejuvenated radio AGN	127
6.7	HI and optical properties	128
6.8	Results from the literature	131
6.8.1	HI and radio properties	131
6.8.2	Optical properties	131
6.9	Conclusions, and future perspectives	134
6.10	Appendix	135
6.10.1	SDSS images and spectra	135

Summary	143
7.1 Chapter 2: The Lockman Hole project	143
7.2 Chapter 3: From star forming to inactive galaxies	144
7.3 Chapter 4: Probing the gas content of AGN with HI absorption stacking	144
7.4 Chapter 5: The HI absorption ‘Zoo’	145
7.5 Chapter 6: HI, radio continuum, and optical properties of radio AGN	146
7.6 Future Prospects	146
Nederlandse Samenvatting	151
8.1 HI gas eigenschappen van sterrenstelsels	152
8.2 Stapelen van HI spectra	153
8.3 Gas en de evolutie van radio AGN	153
8.4 Deze thesis	155
Bibliography	151
Acknowledgements	159

