

University of Groningen

Een gekleurd reagens op SH groepen

Witter, Albert

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1960

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Witter, A. (1960). Een gekleurd reagens op SH groepen: toepassing bij de bepaling van de aminozuurvolgorde rondom de cysteinerest in serumalbuminen s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUMMARY

The yellow sulfhydryl reagent, N-(4-dimethylamino-3,5-dinitrophenyl)maleimide (DDPM) has been synthesized and its reaction with cysteine investigated. The addition product of cysteine and DDPM, S-DDPP-cysteine, was found to hydrolyse slowly to the corresponding maleamic acid, even at $p_H = 6.7$. Another reaction, which occurred at slightly acidic p_H , was an intramolecular transamidation reaction, involving the attack of the α -NH₂ group of the cysteine residue at the imide CO group, thus leading to a 1,4-thiazine derivative (Chapter II).

The reaction of DDPM with the SH group of serum albumin (bovine and human) leads to the yellow S-DDPP-serum albumins. By subjecting these to hydrolysis with pepsin, yellow S-DDPP-cysteinylopeptides were formed. These were isolated from the digest by adsorption on talc, paperelectrophoresis and paperchromatography. The amino acid sequences were determined by classical methods, using 1-fluoro-2,4-dinitrobenzene for N-terminal amino acid determination, carboxypeptidase for the C-terminal amino acid determination and subtilisin hydrolysis and partial acid hydrolysis to obtain peptidefragments. The amino acid sequence: leu-glu(NH₂)-asp-glu-glu(NH₂)-glu(NH₂)-cysH-pro-phe was found to occur in bovine as well as in human serum albumins (Chapter III).