

University of Groningen

## About the Chiral Stability of Germacrene B and the Biomimetic Synthesis of Guaiane Sesquiterpenes

Minnaard, Adriaan; Wijnberg, Joannes B.P.A.; Groot, Aede de

*Published in:*  
The Journal of Organic Chemistry

*DOI:*  
[10.1021/jo970902r](https://doi.org/10.1021/jo970902r)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1997

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Minnaard, A. J., Wijnberg, J. B. P. A., & Groot, A. D. (1997). About the Chiral Stability of Germacrene B and the Biomimetic Synthesis of Guaiane Sesquiterpenes. *The Journal of Organic Chemistry*, 62(21). DOI: 10.1021/jo970902r

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## About the Chiral Stability of Germacrene B and the Biomimetic Synthesis of Guaiane Sesquiterpenes

Adriaan J. Minnaard, Joannes B. P. A. Wijnberg,\* and Aede de Groot\*

Laboratory of Organic Chemistry, Agricultural University, Dreijenplein 8,  
6703 HB Wageningen, The Netherlands

Received May 20, 1997<sup>o</sup>

The planar chirality of 15-hydroxygermacrene B (**2**) has been examined by means of the asymmetric Sharpless epoxidation, performed as a kinetic resolution. <sup>1</sup>H NMR experiments with Eu(hfc)<sub>3</sub> demonstrate that the recovered **2** is racemic. Consequently, **2** and most likely also germacrene B (**1**) are not enantiomerically stable at room temperature. The formation of the optically active cis-fused guaiane **4** of high ee with limited Sharpless reagent shows that the asymmetric epoxidation of **2** proceeds highly enantioselectively. The Sharpless epoxidation methodology applied on the (*E,Z*)-germacrene **3** results in the formation of the stable optically active epoxide **5**. Acid-induced cyclization of **5**, leading to a mixture of guaianes, probably proceeds via the trans-fused carbocationic intermediate **A**.

### Introduction

Transannular cyclization reactions of (*E,E*)-germacranes and their monoepoxides have attracted much attention in connection with sesquiterpene biosynthesis. It has been shown<sup>1</sup> that a concerted mechanism in combination with a fixed conformation of the cyclodeca-1(10),4-diene ring system<sup>2</sup> of (*E,E*)-germacranes, at least in the transition state, accounts for the regio- and stereospecificity of these reactions. The 10-membered ring system of (*E,E*)-germacranes can adopt four principal conformations in which the Me(14) and Me(15) group<sup>3</sup> are located at both sides of the ring. These conformers have either a crossed or parallel relation of the double bonds and are interconvertible by rotation of each of the double bonds.<sup>4</sup> Independently, the C(7) substituent of (*E,E*)-germacranes may have two possible orientations, which in turn are related via C(7)–C(8) inversion.<sup>5</sup> In addition, (*E,E*)-germacranes display planar chirality<sup>6</sup> because the two endocyclic double bonds are approximately perpendicular to the plane of the 10-membered ring.

For germacrene B (**1**),<sup>7</sup> a widespread naturally occurring (*E,E*)-germacrane with *sp*<sup>2</sup> hybridization at C(7), the interconversion operations described above count for eight low-energy conformers: four diastereomers and their respective enantiomers. According to MNDO<sup>8</sup> and MM2<sup>9</sup> calculations on **1**, the crown conformation in which

the endocyclic double bonds possess the crossed orientation with both Me(14) and Me(15) at the same side of the molecule is energetically most favorable. The preference of **1** for the crown conformation is supported by X-ray analysis of the 1:1 adduct of **1** with AgNO<sub>3</sub>.<sup>10</sup> This X-ray analysis also revealed the presence of both the enantiomers (*S,S*)-**1** and (*R,R*)-**1** (1:1 ratio) in the adduct (Scheme 1).<sup>11</sup> In another calculation study on the conformational aspects of **1**, the suggestion has been made that resolution of (*S,S*)-**1** and (*R,R*)-**1** should be possible but that the racemization process should be quite facile.<sup>12</sup> It was therefore challenging to study the chiral stability of **1** and related compounds<sup>13</sup> in more detail, all the more so because the *in vivo* formation of **1** is assumably enantioselective.<sup>14</sup>

The successful synthesis of 15-hydroxygermacrene B (**2**)<sup>15</sup> permitted the use of the asymmetric Sharpless epoxidation, performed as a kinetic resolution,<sup>16</sup> as a direct method for studying the chiral stability of this type of (*E,E*)-germacranes<sup>17</sup> through determination of the ee of recovered **2**<sup>18</sup> by means of the enantiopure shift reagent Eu(hfc)<sub>3</sub>.<sup>19</sup> Additionally, it was expected that transannular cyclization of the asymmetric epoxidation product

<sup>o</sup> Abstract published in *Advance ACS Abstracts*, September 15, 1997.

(1) For representative examples, see: (a) Allen, F. H.; Brown, E. D.; Rogers, D.; Sutherland, J. K. *J. Chem. Soc., Chem. Commun.* **1967**, 1116. (b) Sutherland, J. K. *Tetrahedron* **1974**, *30*, 1651. (c) Brown, E. D.; Sam, T. W.; Sutherland, J. K.; Torre, A. *J. Chem. Soc., Perkin Trans. 1* **1975**, 2326. (d) Brown, E. D.; Sutherland, J. K.; Sam, T. W. *J. Chem. Soc., Perkin Trans. 1* **1975**, 2332. (e) Sam, T. W.; Sutherland, J. K. *J. Chem. Soc., Perkin Trans. 1* **1975**, 2336.

(2) The numbering system as given in structure **1** will be followed throughout the text of this paper.

(3) In most naturally occurring (*E,E*)-germacranes, C(14) and C(15) are Me groups. (*E,E*)-Germacranes in which C(14) or C(15) is oxidized are also frequently found in nature, see: Connolly, J. D.; Hill, R. A. *Dictionary of Terpenoids*; Chapman & Hall: London, 1991; Vol. 1, Mono- and Sesquiterpenoids.

(4) During the rotation of a double bond, the vinylic hydrogen passes through the ring.

(5) Wharton, P. S.; Poon, Y. C.; Kluender, H. C. *J. Org. Chem.* **1973**, *38*, 735.

(6) Eliel, E. L.; Wilen, S. H.; Mander, L. N. *Stereochemistry of Organic Compounds*; Wiley Interscience: New York, 1994; pp 1166–1176.

(7) Nishimura, K. *Tetrahedron Lett.* **1969**, 3097.

(8) Fransen, H. R.; Dormans, G. J. M.; Buck, H. M. *Tetrahedron* **1983**, *39*, 2981.

(9) Watson, W. H.; Kashyap, R. P. *J. Org. Chem.* **1986**, *51*, 2521.

(10) (a) Allen, F. H.; Rogers, D. *J. Chem. Soc., Chem. Comm.* **1967**, 588. (b) Allen, F. H.; Rogers, D. *J. Chem. Soc. B* **1971**, 257.

(11) Prefixes like *S,S* denote the chirality associated with the C(1)–C(10) and the C(4)–C(5) double bond, respectively, see: Cahn, R. S.; Ingold, C.; Prelog, V. *Angew. Chem.* **1966**, *78*, 413. For simplicity, C(7)–C(8) inversion is not considered.

(12) Shirahama, H.; Osawa, E.; Matsumoto, T. *Tetrahedron Lett.* **1979**, 2245.

(13) Germacrene has been partly resolved by asymmetric reduction and back oxidation: Hill, R. K.; Fracheboud, M. G.; Sawada, S.; Carlson, R. M.; Yan, S.-J. *Tetrahedron Lett.* **1978**, 945.

(14) Cane, D. E. *Chem. Rev.* **1990**, *90*, 1089.


(15) See preceding paper.

(16) Marshall *et al.* have used the same method in the synthesis of optically active betweenanenes and related (*E*)-cycloalkenes: (a) Marshall, J. A.; Audia, V. H. *J. Org. Chem.* **1987**, *52*, 1106. (b) Marshall, J. A.; Audia, V. H.; Jenson, T. M.; Guida, W. C. *Tetrahedron* **1986**, *42*, 1703 and references cited therein.


(17) It was assumed that replacement of a Me(15) hydrogen atom by a hydroxyl group would have no effect on the conformational behavior.

(18) It is obvious that this ee will depend on the racemization rate of **2**.

(19) (+)-Eu(hfc)<sub>3</sub> stands for europium tris[3-(heptafluoropropyl)-hydroxymethylene]-(+)-camphorate].



of **2** would result in the formation of an optically active cis-fused guaiane.<sup>20</sup>




It is well-known that, with few exceptions, transannular cyclizations of (*E,E*)-germacranes, or more likely their respective 1,10- and 4,5-epoxides, give eudesmane and guaiane sesquiterpenes.<sup>20,21</sup> On the other hand, knowledge of the possible transannular cyclizations of the geometric isomers of (*E,E*)-germacranes, i.e., (*Z,E*)-germacranes (melampolides) and (*E,Z*)-germacranes (heliangolides), appears to be limited to a few studies,<sup>22,23</sup> and the accumulated results do not permit any generalization. With the heliangolide **3** available,<sup>15</sup> it was obvious to investigate the cyclization behavior of the 4,5-epoxide of **3** too. From this cyclization study, it was expected that new information about the possible role of heliangolides in the biosynthesis of sesquiterpenes would be obtained.

## Results and Discussion

The conformational behavior and enantiomer composition of **2** was studied first. Information about its conformational behavior was obtained from <sup>1</sup>H NMR measurements at room and lower temperatures. The signals in the <sup>1</sup>H NMR spectrum of **2** recorded at room temperature were broadened but singular.<sup>24</sup> At low temperature, the number of signals increased, indicating the presence of more than one conformer. Although these observations do not necessarily imply that **2** racemizes at room temperature, they indicate that an averaged <sup>1</sup>H NMR spectrum is obtained at room temperature.

The enantiopure shift reagent (+)-Eu(hfc)<sub>3</sub> was used to determine the enantiomer composition of **2**. In the



normal <sup>1</sup>H NMR spectrum of **2**, the carbinyl protons at C(15) appeared as an AB quartet ( $J = 11.7$  Hz,  $\Delta\nu/J = 5.4$ ). After addition of (+)-Eu(hfc)<sub>3</sub>, the signals of these protons were strongly shifted and doubled, which points to the racemic nature of **2**.<sup>25</sup>

As mentioned in the Introduction, a direct way to establish the chiral stability of **2** involves the asymmetric Sharpless epoxidation, performed as a kinetic resolution. Since **2** probably exists in a single (crown) conformation (vide supra), it is expected that the epoxidation of **2** will only occur from the outer face because the  $\pi$  lobe of the C(4)–C(5) double bond at the interior of the 10-membered ring is effectively shielded and not available for reaction.<sup>26</sup> Consequently, only one enantiomer can undergo reagent preferred epoxidation and none of the diastereomeric epoxy alcohol would be expected to form. The Sharpless epoxidation of **2** was carried out following well-described procedures with (+)-diethyl tartrate (DET) as the chiral ligand.<sup>27</sup> The reaction was designed as a kinetic resolution using 0.55 equiv of *tert*-butyl hydroperoxide (*t*-BuOOH) and stoichiometric amounts of titanium isopropoxide (Ti(OiPr)<sub>4</sub>) at  $-23$  °C. Under these circumstances, the asymmetric epoxidation of **2** produced the optically active guaiane **4** ( $[\alpha]_D -52^\circ$ ) in 17% yield, together with small amounts of unidentified products (Scheme 2). The ee of **4** was found to be 92% through chiral GC analysis.<sup>28</sup> Aqueous AgNO<sub>3</sub> extraction and chromatographical purification were applied to regain unreacted **2** (46%). <sup>1</sup>H NMR experiments with Eu(hfc)<sub>3</sub> showed that this recovered **2** was racemic.<sup>29</sup> This result suggests that the unreacted **2** racemizes either during the epoxidation reaction at  $-23$  °C or as it warms from  $-23$  °C to room temperature during the isolation process. Whatever the case may be, **2** is not enantiomerically stable at room temperature. Assuming that replacement of the hydroxyl group in **2** by a hydrogen atom has no influence on the racemization rate, germacrene B (**1**) will also be enantiomerically unstable at room temperature. Therefore, these compounds cannot be added to the list of enantiomerically stable (*E*)-cycloalkenes.<sup>6</sup> When the asymmetric epoxidation of **2** was performed with 1.1 equiv of *t*-BuOOH, **4** of 56% ee was formed in 36% chemical yield. This finding indicates that during epoxidation at  $-23$  °C with a full equivalent of Sharpless reagent, racemization of **2** occurs slowly relative to the asymmetric epoxidation, so the more reactive enantiomer is not constantly replenished, and, hence, the less reactive enantiomer is epoxidized as well.

(20) The transannular cyclization of (*E,E*)-germacrane 4,5-epoxides constitutes an important step in the biosynthesis of cis-fused guaianes, see: Fischer, N. H. In *Recent Advances in Phytochemistry*, Vol. 24: *Biochemistry of the Mevalonic Acid Pathway to Terpenoids*; Towers, G. H. N., Stafford, H. A., Eds.; Plenum Press: New York, 1990; Chapter 4.

(21) Marco, J. A.; Sanz-Cervera, J. F.; García-Lliso, V.; Domingo, L. R.; Carda, M.; Rodríguez, S.; López-Ortiz, F.; Lex, J. *Liebigs Ann. Chem.* **1995**, 1837 and references cited therein.

(22) Cyclization of geometric isomers of simple (*E,E*)-germacranes have been performed, inter alia, see: (a) Kodama, M.; Yokoo, S.; Matsuki, Y.; Itô, S. *Tetrahedron Lett.* **1979**, 1687 and references cited therein. (b) Williams, J. R.; Callahan, J. F.; Blount, J. F. *J. Org. Chem.* **1981**, *46*, 2665.

(23) (a) Toma, K.; Murae, T.; Takahashi, T. *Chem. Lett.* **1982**, 551. (b) De Pascual Teresa, J.; González, M. S.; Caballero, M. C.; Parra, T.; Bellido, I. S. *Tetrahedron Lett.* **1987**, *28*, 821. (c) González, A. G.; Galindo, A.; Afonso, M. M.; Mansilla, H.; López, M. *Tetrahedron* **1988**, *44*, 4585. (d) Delgado, G.; Guzmán, S. *J. Chem. Soc., Chem. Commun.* **1992**, 606.

(24) In the NMR spectra of **1**, similar effects were observed.

(25) Siever, R. E.; Kime, K. A. *Aldrichimica Acta* **1977**, *10*, 54.

(26) A similar reasoning has been made for the biological epoxidation of (*E,E*)-germacranes, see: Fischer, N. H. *Rev. Latinoamer. Quim.* **1978**, *9*, 41.

(27) (a) Johnson, R. A.; Sharpless, K. B. In *Catalytic Asymmetric Synthesis*; Ojima, I., Ed.; VCH Publishers: New York, 1993; Chapter 4. (b) Katsuki, T.; Martin, V. S. *Org. React.* **1996**, *48*, 1.

(28) A gas chromatograph equipped with a Supelco 30 m BETA-DEX<sup>TM-120</sup> fused silica capillary column, 0.25 mm i.d. and 0.25  $\mu$ m film thickness, was used. H<sub>2</sub> was the carrier gas. Peaks were partly resolved and integrated manually.


(29) In different runs the optical rotation did not exceed 5°.

Initially, the elucidation of the relative stereochemistry of **4** was troublesome because of overlapped signals in its  $^1\text{H}$  NMR spectrum recorded in  $\text{CDCl}_3$  and  $\text{C}_6\text{D}_6$ . The use of a 1:1:1 mixture of  $\text{CDCl}_3$ ,  $\text{C}_6\text{D}_6$ , and  $\text{C}_5\text{D}_5\text{N}$ , however, resulted in unobscured signals for both the bridgehead protons H-1 and H-5. NOE-difference experiments showed a clear NOE between these two protons, thereby establishing the cis fusion of the guaiane skeleton. The NOE observed between H-1 and the 2-propoxy group ascertained the stereochemistry at C(10). Because the C(15) carbinol group and H-5 necessarily possess the trans orientation, the stereochemistry of **4** is as depicted in Scheme 2. Structure **4** also reflects the absolute configuration. The assignment is based on the enantioselectivity principles of the asymmetric Sharpless epoxidation according to which epoxidation of **2** will preferentially take place from the *S,S* conformation.<sup>30</sup>

The formation of cis-fused (–)-**4** must proceed via the resolved epoxide alcohol, initially formed in the Sharpless epoxidation of **2** (see Scheme 2). Under the reaction conditions employed, this epoxide alcohol immediately reacts further by way of epoxide ring opening, probably induced by titanium species acting as Lewis acids, transannular cyclization, and incorporation of 2-propanol present in the reaction medium to afford (–)-**4**. Although other stereochemical courses for the cyclization of (*E,E*)-germacranes cannot be excluded,<sup>31</sup> the formation of (–)-**4** provides strong evidence for the involvement of (*E,E*)-germacrane epoxides in the biosynthesis of cis-fused guaianes.<sup>20</sup>

Beside guaianes possessing a cis-fused ring system, a number of trans-fused guaianes has been found in nature. In most of these trans-fused guaianes, H-5 and C(15) are located at different sides of the molecule.<sup>3</sup> It has been postulated that the biosynthesis of these *trans*-guaianes proceeds via the 4,5-epoxides of melampolides.<sup>32</sup> The isolation of guaianes possessing a syn relationship between H-5 and C(15),<sup>33</sup> on the other hand, suggests the involvement of heliangolides in guaiane biosynthesis. In order to test the latter hypothesis, the heliangolide alcohol **3** was converted into its 4,5-epoxide and subjected to cyclization conditions.<sup>34</sup>

The Sharpless epoxidation of **3** was performed with excess *t*-BuOOH and afforded, after workup and purification, the stable epoxide **5** ( $[\alpha]_D -145^\circ$ )<sup>35</sup> in 52% yield (Scheme 3). The sharp signals and couplings in the  $^1\text{H}$  NMR spectrum suggest that **5** probably exists in one conformation at room temperature. Upon treatment of **5** with acid, a fast reaction was observed which resulted in a mixture of essentially four guaianic products in a 1:1:1:4 ratio as judged by GC. Separation of this product mixture appeared to be troublesome, and only after repeated chromatography on silica gel, one of the minor compounds, **8**, could be obtained in almost pure form. A



more successful separation of the remaining product mixture was achieved with  $\text{AgNO}_3$ -impregnated silica gel. In this way, the main product **6** ( $[\alpha]_D +48^\circ$ )<sup>36</sup> and another minor compound, **7**, contaminated with small amounts of **8**, were obtained. The fourth product, most likely the C(9)–C(10) double bond isomer of **6** and **7**,<sup>37</sup> was lost during the latter purification step. The absence of olefinic signals in the  $^1\text{H}$  NMR spectrum of **6** and the appearance of four olefinic singlets in its  $^{13}\text{C}$  NMR spectrum are consistent with the assigned structure. The presence of an exocyclic double bond in **7** follows from the appearance of a broad two-proton singlet at  $\delta$  4.72 ( $^1\text{H}$ ) and an olefinic triplet at  $\delta$  107.23 ( $^{13}\text{C}$ ) in its NMR spectra. The vinylic one-proton doublet ( $J = 0.9$  Hz) and the three-proton doublet ( $J = 7.0$  Hz) at  $\delta$  5.32 and 1.02, respectively, in the  $^1\text{H}$  NMR spectrum of **8** are typical and, together with the other  $^1\text{H}$  and  $^{13}\text{C}$  NMR data, in full agreement with the assigned structure.

The formation of **6**, **7**, and **8** presumably proceeds via the intermediate **A**. This tertiary carbocation can undergo proton loss to give the isomeric guaianes **6** and **7**, but can also react further by way of two consecutive 1,2-H shifts ( $\text{C}(1) \rightarrow \text{C}(10)$  and  $\text{C}(5) \rightarrow \text{C}(1)$ ) to afford another tertiary carbocationic intermediate (**B**). Proton loss leading to a conjugated diene system accounts for the formation of **8** from **B**. Because tandem 1,2-H shifts are suprafacial and occur most likely via a concerted process,<sup>38</sup> the ring junction in the initially formed intermediate **A** is probably *trans*. Combined with the syn relationship between H-5 and C(15), this has led to the stereochemistry depicted in the structures **6**, **7**, and **8** (Scheme 3).

(30) With (+)-DET as the chiral ligand, (*R,R*)-**2** remains largely unoxidized, see ref 27a, pp 104–108.

(31) Kuroyanagi, M.; Ueno, A.; Koyoma, K.; Natori, S. *Chem. Pharm. Bull.* **1990**, *38*, 55.

(32) Herz, W. In *Chemistry in Botanical Classification. Nobel Symposium 25*; Bendz, G., Santesson, T., Eds.; Academic Press: New York, 1973; pp 153–172.

(33) (a) Bohlmann, F.; Borthakur, N.; Jakupovic, J.; Pickard, J. *Phytochemistry* **1982**, *21*, 1357. (b) Barrero, A. F.; Sánchez, J. F.; Molina, J.; Barrón, A.; Del Mar Salas, M. *Phytochemistry* **1990**, *29*, 3575.

(34) The possible role of melampolides in guaiane biosynthesis is currently studied at our department. To be published.

(35) Chiral GC could not be used to establish the ee of **5** because of its thermal instability.

(36) Because only a single peak was observed, the ee of **6** could not be determined with chiral GC. In this respect, it should be noted that the enantioselectivity of the Sharpless epoxidation of allylic alcohols almost invariably approaches 100%. See reference 27a, p 104.

(37) The presence of this isomer was deduced from the appearance of a broad triplet at  $\delta$  5.52 in the  $^1\text{H}$  NMR spectrum of the crude product mixture.

(38) Carey, F. A.; Sundberg, R. J. *Advanced Organic Chemistry*; Plenum Press: New York, 1990; part A, p 314.

The reaction outcome of the cyclization of **5** reinforces our hypothesis that heliangolides can act as precursors for guaianes with H-5 and C(15) in a syn arrangement. The formation of **8** from **5**, in which probably 1,2-H shifts are involved, further suggests that the biosynthesis of naturally occurring guaianes with a C(5)–C(6) double bond and a syn relationship between H-1 and C(14)<sup>39</sup> probably proceeds in the same way. Similar 1,2-H shifts have also been proposed for the biosynthesis of pseudoguaianes,<sup>40</sup> but in that case a C(4) → C(5) shift of C(15) instead of proton loss follows the 1,2-H shifts. It is obvious that the syn orientation of H-5 and C(15), as in intermediate **A**, prevents a concerted 1,2-C shift.

### Concluding Remarks

The question whether germacrene B (**1**) would be enantiomerically stable could be answered by the asymmetric Sharpless epoxidation of **2**, performed as a kinetic resolution. It appears that the racemization rate of **2** is fast at room temperature and this conclusion can safely be extrapolated to germacrene B itself.

The formation of the optically active guaiane **4** of high ee with limited Sharpless reagent shows that the asymmetric epoxidation of **2** proceeds highly enantioselectively. Therefore, enantioselective *in vivo* epoxidation of germacrene B may be possible as well.<sup>41</sup> In addition, the cis ring junction in **4** indicates that **2** preferably reacts from the most stable crown conformation, which is consistent with previous studies.<sup>1</sup>

The acid-induced transannular cyclization of the heliangolide 4,5-epoxide **5**, in which the guaianes **6** and **7** and the rearranged guaiane **8** are formed, may represent a possible biosynthetic pathway toward guaianes with an aberrant stereochemistry at C(4).<sup>33,39a</sup>

### Experimental Section<sup>42</sup>

**Materials.** All reagents were purchased from Aldrich or Janssen. The compounds **2** and **3** were prepared as described.<sup>15</sup>

**Kinetic Resolution/Asymmetric Epoxidation of 2.** To a stirred solution of 0.134 mL (0.45 mmol) of Ti(OiPr)<sub>4</sub> in 2 mL of dry CH<sub>2</sub>Cl<sub>2</sub> was added 0.090 mL (0.52 mmol) of (+)-DET at –23 °C. The solution was stirred for 5 min, and then a solution of 0.100 g (0.45 mmol) of **2** in 2 mL of CH<sub>2</sub>Cl<sub>2</sub> was added. After stirring at –23 °C for an additional 30 min, 0.045 mL (≈ 0.25 mmol) of *t*-BuOOH (5–6 M in decane) was added. Stirring was continued at –23 °C for 1 h, and then a mixture of 8 mL of acetone and 0.5 mL of water was added. After being stirred at –23 °C for another 30 min and at 0 °C for 3 h, the reaction mixture was filtered, and the filtrate was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were washed with brine, dried, and evaporated at rt. The remaining residue was dissolved in 10 mL of *tert*-butyl methyl ether and added to a

mixture of 5 mL of brine and 0.2 g of NaOH at 0 °C. The two-phase mixture was stirred at 0 °C for 1 h. After separation and drying, the organic layer was evaporated at rt to afford 0.115 g of the crude product mixture. This mixture was dissolved in 10 mL of *tert*-butyl methyl ether and extracted with five portions of 20% aqueous AgNO<sub>3</sub>. The combined aqueous layers were washed with *tert*-butyl methyl ether and cooled to 0 °C. After addition of 15 mL of 25% aqueous NH<sub>3</sub>, the aqueous layer was extracted with *tert*-butyl methyl ether. The combined organic layers were washed with brine, dried, and evaporated at rt to afford 0.037 g (37%) of the starting material **2**. The *tert*-butyl methyl ether solution remaining after aqueous AgNO<sub>3</sub> extraction was washed with brine, dried, and evaporated. Flash chromatography (3:1 to 3:2 petroleum ether (bp 40–60 °C)/*tert*-butyl methyl ether) gave, in order of elution, an additional 0.009 g (9%) of **2** and 0.022 g (17%) of [(1*R*)-(1*α*,3*αα*,4*α*,8*αα*)]-decahydro-1-hydroxymethyl-4-methyl-7-(1-methylethylidene)-4-[(1-methylethyl)-oxy]-1-azulenol (**4**): [α]<sub>D</sub> –52.3° (c 0.38), 92% ee;<sup>29</sup> <sup>1</sup>H NMR<sup>43</sup> δ 1.07 (d, *J* = 6.5 Hz, 3 H), 1.11 (d, *J* = 6.5 Hz, 3 H), 1.11 (s, 3 H), 1.46–1.66 (m, 4 H), 1.73–1.85 (m, 2 H), 1.75 (br s, 3 H), 1.90 (br s, 3 H), 1.92–2.00 (m, 2 H), 2.21 (br dd, *J* = 5.8, 12.4 Hz, 1 H), 2.62 (br d, *J* = 12.8 Hz, 1 H), 2.87 (ddd, *J* = 5.3, 14.1, 17.0 Hz, 1 H), 3.07 (br ddd, *J* ≈ 6, 6, 13 Hz, 1 H), 3.73 (d, AB system, *J* = 10.8 Hz, 1 H), 3.82 (septet, *J* = 6.5 Hz, 1 H), 3.91 (d, AB system, *J* = 10.8 Hz, 1 H), 4.33 (br s, 2 OH); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 20.06 (q), 20.27 (q), 24.64 (t), 25.20 (q), 25.30 (q), 26.85 (t), 27.00 (t), 28.59 (q), 31.54 (t), 32.50 (t), 47.93 (d), 50.22 (d), 62.40 (d), 66.85 (t), 77.05 (s), 84.91 (s), 122.63 (s), 130.86 (s); MS *m/z* (relative intensity) 296 (M<sup>+</sup>, 1), 236 (70), 218 (28), 205 (100), 187 (66), 147 (25), 122 (67), 107 (27), 43 (32); HRMS calcd for C<sub>18</sub>H<sub>32</sub>O<sub>3</sub> (M<sup>+</sup>) 296.2351, found 296.2355.

The same procedure was employed by using 0.062 g (0.28 mmol) of **2** and 0.062 mL (≈ 0.31 mmol) of *t*-BuOOH (5–6 M in decane), except that the AgNO<sub>3</sub> extraction was omitted. Flash chromatography (3:1 to 3:2 petroleum ether (bp 40–60 °C)/*tert*-butyl methyl ether) afforded 0.004 g (6%) of unreacted **2** and 0.029 g (36%) of **4**: [α]<sub>D</sub> –35.5° (c 0.69), 56% ee.

**(1*R*,2*S*)-7-Methyl-4-(1-methylethylidene)-(1,2-*b*)-oxiranyl-7-cyclodecenemethanol (5).** The procedure used for the epoxidation of **2** was employed by using 0.123 g (0.56 mmol) of **3** and 0.28 mL (≈ 1.40 mmol) of *t*-BuOOH (5–6 M in decane). The reaction mixture was stored at –20 °C in the refrigerator overnight and worked up as described. Flash chromatography (2:1 petroleum ether (bp 40–60 °C)/*tert*-butyl methyl ether) afforded 0.067 g (52%) of **5**: [α]<sub>D</sub> –145° (c 0.70); <sup>1</sup>H NMR δ 1.31 (ddd, *J* = 4.8, 13.5, 13.5 Hz, 1 H), 1.65 (s, 6 H), 1.73 (s, 3 H), 1.80–2.30 (m, 8 H), 2.51 (m, 1 H), 2.83 (br d, *J* = 15.0 Hz, 1 H), 3.07 (br d, *J* = 10.0 Hz, 1 H), 3.55 (dd, AB system, *J* = 6.8, 12.2 Hz, 1 H), 3.83 (dd, AB system, *J* = 4.3, 12.2 Hz, 1 H), 5.30 (dd, *J* = 8.0, 8.0 Hz, 1 H); <sup>13</sup>C NMR δ 17.02 (q), 20.63 (q), 21.53 (q), 22.78 (t), 27.99 (t), 33.82 (t), 34.18 (t), 38.63 (t), 60.60 (d), 64.07 (t), 64.73 (s), 122.83 (d), 128.11 (s), 129.82 (s), 135.88 (s); MS *m/z* (relative intensity) 236 (M<sup>+</sup>, 18), 221 (36), 205 (70), 147 (100), 135 (94), 121 (73), 107 (96), 93 (87), 55 (80); HRMS calcd for C<sub>15</sub>H<sub>24</sub>O<sub>2</sub> (M<sup>+</sup>) 236.1776, found 236.1778.

**Acid-Induced Cyclization of 5.** To a stirred solution of 0.040 g (0.17 mmol) of **5** in 5 mL of dry CH<sub>2</sub>Cl<sub>2</sub> at room temperature (rt) was added 0.01 g (0.05 mmol) of TsOH dissolved in 1.5 mL of dry CH<sub>2</sub>Cl<sub>2</sub>. After being stirred for 10 min, the reaction mixture was diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were washed with saturated aqueous NaHCO<sub>3</sub> and brine, dried, and evaporated to afford 0.040 g of a crude product mixture. Repeated flash chromatography (3:1 hexane/EtOAc) gave 0.002 g of almost pure **8**. Flash chromatography of the remaining mixture using silica gel impregnated with 7% of AgNO<sub>3</sub> afforded 0.010 g of **6** and 0.002 g of **7**.<sup>44</sup> The spectroscopic data of **6**, **7**, and **8** are given below.

**[(1*R*)-(1*α*,8*αβ*)]-Octahydro-1-hydroxymethyl-4-methyl-7-(1-methylethylidene)-1-azulenol (6):** [α]<sub>D</sub> +48° (c 0.49);

(43) This spectrum was recorded in a 1:1:1 mixture of CDCl<sub>3</sub>, C<sub>6</sub>D<sub>6</sub>, and C<sub>5</sub>D<sub>5</sub>N.

(44) The <sup>1</sup>H NMR spectrum of **7** revealed the presence of a small amount of **8**.

(39) (a) Bohlmann, F.; Jakupovic, J. *Phytochemistry* **1979**, *18*, 131. (b) Yu, S.; Fang, N.; Mabry, T. J.; Abboud, K. A.; Simonsen, S. H. *Phytochemistry* **1988**, *27*, 2887. (c) Zdero, C.; Bohlmann, F.; King, R. M. *Phytochemistry* **1990**, *29*, 3201. (d) Gao, F.; Wang, H.; Mabry, T. J.; Watson, W. H.; Kashyap, R. P. *Phytochemistry* **1990**, *29*, 551. (e) Gao, F.; Wang, H.; Mabry, T. J. *Phytochemistry* **1990**, *29*, 1601.

(40) (a) Fischer, N. H.; Wiley, R. A.; Perry, D. L. *Rev. Latinoamer. Quim.* **1976**, *7*, 87. (b) Herz, W. *Israel J. Chem.* **1977**, *16*, 32.

(41) The biotransformation of racemic germacrene to enantiomerically enriched germacrene-4,5-epoxide by plant cell cultures has been reported, see: (a) Hikino, H.; Konno, C.; Nagashima, T.; Kohama, T.; Takemoto, T. *Chem. Pharm. Bull.* **1977**, *25*, 6. (b) Sakui, N.; Kuroyanagi, M.; Ishitobi, Y.; Sato, M.; Ueno, A. *Phytochemistry* **1992**, *31*, 143. (c) Sakamoto, S.; Tsuchiya, N.; Kuroyanagi, M.; Ueno, A. *Phytochemistry* **1994**, *35*, 1215.

(42) For a general description of the experimental procedures employed in this research, see: Minnaard, A. J.; Wijnberg, J. B. P. A.; de Groot, A. *Tetrahedron* **1994**, *50*, 4755.

$^1\text{H}$  NMR  $\delta$  1.56 (br s, 3 H), 1.60 (br s, 6 H), 1.55–2.60 (m, 13 H), 3.46 (d, AB system,  $J = 11.0$  Hz, 1 H), 3.65 (d, AB system,  $J = 11.0$  Hz, 1 H);  $^{13}\text{C}$  NMR  $\delta$  20.33 (2 q), 22.21 (q), 28.82 (t), 29.47 (t), 31.94 (t), 35.17 (t), 36.07 (t), 47.35 (d), 68.49 (t), 83.12 (s), 122.71 (s), 128.99 (s), 132.07 (s), 137.40 (s); MS  $m/z$  (relative intensity) 236 ( $\text{M}^+$ , 46), 218 (20), 205 (100), 187 (30), 149 (15), 123 (6), 105 (16), 97 (39); HRMS calcd for  $\text{C}_{15}\text{H}_{24}\text{O}_2$  ( $\text{M}^+$ ) 236.1776, found 236.1778.

**[(1*R*)-(1 $\alpha$ ,3 $\alpha$ ,8 $\alpha$ )]-Decahydro-1-hydroxymethyl-4-methylidene-7-(1-methylethylidene)-1-azulenol (7):**  $^1\text{H}$  NMR  $\delta$  1.20 (ddd,  $J = 3.8, 11.7, 11.7$  Hz, 1 H), 1.55–2.15 (m, 5 H), 1.60 (br s, 2 OH), 1.65 (br s, 3 H), 1.67 (br s, 3 H), 2.25–2.65 (m, 6 H), 3.50 (dd, AB system, 5.3, 10.7 Hz, 1 H), 3.62 (dd, AB system, 3.2, 10.7 Hz, 1 H), 4.72 (br s, 2 H);  $^{13}\text{C}$  NMR  $\delta$  20.25 (2 q), 26.68 (t), 30.96 (t), 31.69 (t), 36.35 (t), 36.62 (t), 49.08 (d), 53.34 (d), 69.58 (t), 82.10 (s), 107.23 (t), 124.87 (s), 130.62 (s), 152.58 (s); MS  $m/z$  (relative intensity) 236 ( $\text{M}^+$ , 15), 218 (17), 206 (15), 205 (100), 187 (37), 145 (11), 131 (12), 105 (11), 91 (14); HRMS calcd for  $\text{C}_{15}\text{H}_{24}\text{O}_2$  ( $\text{M}^+$ ) 236.1776, found 236.1777.

**[(1*R*)-(1 $\alpha$ ,3 $\alpha$ ,4 $\beta$ )]-Octahydro-1-hydroxymethyl-4-methyl-7-(1-methylethylidene)-1-azulenol (8):**  $^1\text{H}$  NMR  $\delta$  1.02

(d,  $J = 7.0$  Hz, 3 H), 1.20 (m, 1 H), 1.60 (br s, 2 OH), 1.68 (br s, 3 H), 1.71 (br s, 3 H), 1.60–2.70 (m, 9 H), 3.61 (dd, AB system,  $J = 4.8, 10.6$  Hz, 1 H), 3.72 (br d, AB system,  $J = 10.6$  Hz, 1 H), 5.32 (br s, 1 H);  $^{13}\text{C}$  NMR  $\delta$  20.19 (q), 20.66 (q), 20.77 (q), 29.93 (t), 30.52 (t), 33.30 (t), 35.47 (d), 42.80 (t), 51.01 (d), 69.30 (t), 83.55 (s), 120.06 (d), 124.44 (s), 132.31 (s), 152.68 (s); MS  $m/z$  (relative intensity) 236 ( $\text{M}^+$ , 57), 218 (54), 205 (100), 187 (86), 149 (30), 145 (32), 131 (30), 121 (32), 91 (32), 55 (33), 41 (30); HRMS calcd for  $\text{C}_{15}\text{H}_{24}\text{O}_2$  ( $\text{M}^+$ ) 236.1776, found 236.1776.

**Acknowledgment.** We thank A. van Veldhuizen for recording  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra and H. Jongejan for mass spectral data.

**Supporting Information Available:**  $^1\text{H}$  NMR spectra for compounds **4–8** (5 pages). This material is contained in libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS; see any current masthead page for ordering information.

JO970902R