

University of Groningen

Correction : Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia between 34 and 37 weeks' gestation (HYPITAT-II)

Broekhuijsen, Kim; Langenveld, Josje; van Baaren, Gert-Jan; van Pampus, Marielle G.; van Kaam, Anton H.; Groen, Henk; Porath, Martina; Franssen, Maureen T. M.; Mol, Ben W.; HYPITAT-II Study Grp

Published in:
BMC Pregnancy and Childbirth

DOI:
[10.1186/1471-2393-13-232](https://doi.org/10.1186/1471-2393-13-232)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Broekhuijsen, K., Langenveld, J., van Baaren, G.-J., van Pampus, M. G., van Kaam, A. H., Groen, H., ... HYPITAT-II Study Grp (2013). Correction : Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia between 34 and 37 weeks' gestation (HYPITAT-II): a multicentre, open-label randomised controlled trial (vol 13, pg 232, 2013). BMC Pregnancy and Childbirth, 13, [232]. <https://doi.org/10.1186/1471-2393-13-232>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CORRECTION

Open Access

Correction: Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia between 34 and 37 weeks' gestation (HYPITAT-II): a multicentre, open-label randomised controlled trial

Kim Broekhuijsen^{1*}, Josje Langenveld², Gert-Jan van Baaren³, Mariëlle G van Pampus⁴, Anton H van Kaam³, Henk Groen⁵, Martina Porath⁶, Maureen TM Franssen¹, Ben W Mol³ and HYPITAT-II study group

The earliest draft versions of the protocol for our study described the composite adverse maternal outcome as one or more of progression to severe disease, pulmonary edema, thrombo-embolic disease, HELLP syndrome, eclampsia, placental abruption or maternal death. However, there is ongoing debate as to whether progression to severe disease should be considered an adverse maternal outcome [1,2]. Therefore, after obtaining funding which enabled us to increase our sample size to the current sample size of 680, we decided to study a composite adverse maternal outcome excluding progression to severe disease. These changes were incorporated in the protocol as submitted to and approved by the institutional review board;* the current protocol is available from our website (http://www.studies-obsgyn.nl/hypitat2/page.asp?page_id=642). Unfortunately, the change to the maternal outcome definition was not incorporated into the published protocol, which incorrectly includes progression to severe disease in the composite adverse maternal outcome [3].

We also discovered minor differences between the published protocol and the IRB approved protocol. The definition for neonatal morbidity should have contained meconium aspiration syndrome, pneumothorax and/or pneumomediastinum, periventricular leucomalacia, convulsions and other neurological abnormalities. Finally, low 5-minute Apgar score should have been defined as below 7 (as opposed to below 3), and low umbilical artery pH as below 7.05 (as opposed to below 7.0).

These discrepancies were discovered and the correction submitted for publication during recruitment.

* Medical Ethics Committee, Academic Medical Centre, Amsterdam, the Netherlands (ref. 2008/244).

Author details

¹Department of Obstetrics and Gynecology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

²Department of Obstetrics and Gynecology, Atrium Medical Center, Heerlen, The Netherlands. ³Department of Obstetrics and Gynecology, Academic Medical Center, Amsterdam, The Netherlands. ⁴Department Obstetrics and Gynecology, Onze Lieve Vrouwe Gasthuis, Amsterdam, The Netherlands.

⁵Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands. ⁶Department of Obstetrics and Gynecology, Maxima Medical Center, Veldhoven, The Netherlands.

Received: 24 January 2013 Accepted: 6 December 2013

Published: 23 December 2013

References

1. Koopmans CM, Bijlenga D, Groen H, Vijgen SM, Aarnoudse JG, Bekedam DJ, HYPITAT study group, et al: Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia after 36 weeks' gestation (HYPITAT): a multicentre, open-label randomised controlled trial. *Lancet* 2009, **374**:979-988.
2. Bewley S, Shennan A: Hypitad and the fallacy of pregnancy interruption. *Lancet* 2010, **375**(9709):119.
3. Langenveld J, Broekhuijsen K, van Baaren G, van Pampus MG, van Kaam AH, Groen H, Porath M, Mol BW, HYPITAT-II study group: Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia between 34 and 37 weeks' gestation (HYPITAT-II): a multicentre, open-label randomised controlled trial. *BMC Pregnancy Childbirth* 2011, **11**:50.

doi:10.1186/1471-2393-13-232

Cite this article as: Broekhuijsen et al.: Correction: Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia between 34 and 37 weeks' gestation (HYPITAT-II): a multicentre, open-label randomised controlled trial. *BMC Pregnancy and Childbirth* 2013 **13**:232.

* Correspondence: kbroekhuijsen@gmail.com

¹Department of Obstetrics and Gynecology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands
Full list of author information is available at the end of the article