



University of Groningen

Modal Analysis and Experimental Study of High-Order Mode Contribution to Standing Waves in Quasi-Optical Systems

Jellema, Willem; Withington, S.; Trappe, Neil; Murphy, J. A.; Wild, Wolfgang

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version Publisher's PDF, also known as Version of record

Publication date: 2004

Link to publication in University of Groningen/UMCG research database

Citation for published version (APA):

Jellema, W., Withington, S., Trappe, N., Murphy, J. A., & Wild, W. (2004). Modal Analysis and Experimental Study of High-Order Mode Contribution to Standing Waves in Quasi-Optical Systems. Abstract from 15th International Symposium on Space Teraherb Technolog, Northampton, MA, United States.

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policyIf you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): http://www.rug.nl/research/portal. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Download date: 22-05-2019

Modal Analysis and Experimental Study of High-Order Mode Contribution to Standing Waves in Quasi-Optical Systems

Willem Jellema^{1,4}, Stafford Withington², Neil Trappe³, J. Anthony Murphy³, Wolfgang Wild^{1,4}

¹Space Research Organization of the Netherlands (SRON), P.O. Box 800, 9700 AV Groningen, the Netherlands

In many submillimeter-wave optical systems performance is in some way affected by the presence of standing waves. Usually these effects are studied empirically or only modeled to first-order by assuming a quasi-sinusoidal variation with distance and wavelength. In this paper we present an analytical technique to calculate the coupling between two corrugated horns including the total transmission and reflection properties. In our model we have combined the waveguide mode matching technique with a Gaussian beam mode description of the free space propagation. We illustrate the approach for the case of two coupled corrugated horns as the distance between them is varied. We furthermore present experimental test results confirming that multiple reflections do not always result in a classical periodic standing wave pattern. In particular when the horns are close together and the fields are not well matched, high Q cavity effects are observed resulting in irregular standing wave patterns. These cavity effects can be attributed to high-order modes which get trapped between transmitting and receiving horn. We finally discuss the consequences of our observations for a few examples of typical quasi-optical arrangements illustrating that the presented effects can have important consequences for submillimeter-wave optical systems.

 ²Cavendish Laboratory, Madingley Road, Cambridge CB3 0HE, United Kingdom
³National University of Ireland Maynooth, Co, Kildare, Maynooth, Ireland
⁴Kapteyn Institute, P.O. Box 800, 9700 AV Groningen, the Netherlands