

University of Groningen

Prevalence, risk factors and molecular epidemiology of highly resistant gram negative rods in hospitalized patients in the Dutch region Kennemerland

Souverein, Dennis; Euser, Sjoerd M.; Herpers, Bjorn L.; Diederens, Bram; Houtman, Patricia; van Seventer, Marina; van Ess, Ingeborg; Kluytmans, Jan; Rossen, Johannes; Den Boer, Jeroen W.

Published in:
Antimicrobial Resistance and Infection Control

DOI:
[10.1186/s13756-016-0107-6](https://doi.org/10.1186/s13756-016-0107-6)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Souverein, D., Euser, S. M., Herpers, B. L., Diederens, B., Houtman, P., van Seventer, M., ... Den Boer, J. W. (2016). Prevalence, risk factors and molecular epidemiology of highly resistant gram negative rods in hospitalized patients in the Dutch region Kennemerland. *Antimicrobial Resistance and Infection Control*, 5, [8]. DOI: 10.1186/s13756-016-0107-6

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

RESEARCH

Open Access



Prevalence, risk factors and molecular epidemiology of highly resistant gram negative rods in hospitalized patients in the Dutch region Kennemerland

Dennis Souverein^{1*}, Sjoerd M. Euser¹, Bjorn L. Herpers¹, Bram Diederer¹, Patricia Houtman², Marina van Seventer², Ingeborg van Ess³, Jan Kluytmans⁴, John W. A. Rossen⁵ and Jeroen W. Den Boer¹

Abstract

Background: This paper describes (1) the Highly Resistant Gram Negative Rod (HR-GNR) prevalence rate, (2) their genotypes, acquired resistance genes and (3) associated risk factors of HR-GNR colonization among the hospitalized population in the Dutch region Kennemerland.

Methods: Between 1 October 2013 and 31 March 2014, cross-sectional prevalence measurements were performed in three regional hospitals as part of each hospital's infection control program. Rectal swabs were analyzed at the Regional Public Health Laboratory Kennemerland by direct culturing. Genotypes and acquired resistance genes of positive isolates were determined using Whole Genome Sequencing with the MiSeq instrument (Illumina). Association between several independent variables and HR-GNR positivity was examined using logistic regression models.

Results: Out of 427 patients, 24 HR-GNR positive isolates were recovered from 22 patients, resulting in a regional HR-GNR colonization prevalence (95 % CI) of 5.2 % (3.6–7.9). Of these 22 positive patients, 15 were Extended Spectrum Beta-Lactamase (ESBL) positive (3.5 % (2.1–5.7)), 7 patients were positive for a Fluoroquinolones and Aminoglycosides (Q&A) resistant Enterobacteriaceae (1.6 % (0.8–3.3)) and from one patient (0.2 % (0–1.3)) a *Stenotrophomonas maltophilia* resistant towards co-trimoxazole was isolated. No carbapenemase producing Enterobacteriaceae (CPE), multi-resistant *Acinetobacter* species or multi-resistant *Pseudomonas aeruginosa* were isolated. The ESBL genes found were *bla*_{CTX-M-1} (*n* = 4, 25.0 %), *bla*_{CTX-M-15} (*n* = 3, 18.8 %), *bla*_{CTX-M-27} (*n* = 2, 12.5 %), *bla*_{CTX-M-14b} (*n* = 2, 12.5 %), *bla*_{CTX-M-9} (*n* = 2, 12.5 %), *bla*_{CTX-M-14} (*n* = 1, 6.3 %), *bla*_{CTX-M-3} (*n* = 1, 6.3 %), *bla*_{SHV-11} (*n* = 1, 6.3 %) and *bla*_{SHV-12} (*n* = 1, 6.3 %). Being known HR-GNR positive in the past was the only significant associated risk factor for HR-GNR positivity, odds ratio (95 % CI): 7.32 (1.82–29.35), *p*-value = 0.005.

Conclusions: Similar ESBL prevalence rates and genotypes (3.5 %) were found in comparison to other Dutch studies. When previously HR-GNR positive patients are readmitted, they should be screened for HR-GNR colonization since colonization with GR-GNRs could be prolonged. We recommend for future studies to include all defined HR-GNRs in addition to ESBLs in prevalence studies, in order to obtain a more comprehensive overview of colonization with HR-GNRs.

Keywords: HR-GNRs, ESBL, Prevalence, Risk factors, Whole genome sequencing, Netherlands

* Correspondence: d.souverein@streeklabhaarlem.nl

¹Department of Epidemiology and Infection Prevention, Regional Public Health Laboratory Kennemerland, Boerhaavelaan 26, 2035 RC Haarlem, The Netherlands

Full list of author information is available at the end of the article

Background

Worldwide there is an alarming increase in the prevalence of Highly Resistant Gram Negative Rods (HR-GNRs) among clinical isolates [1–3]. The emergence and spread of HR-GNRs is a public health threat since infections caused by HR-GNRs are associated with an increased risk of morbidity, mortality, and healthcare costs (estimated mean additional costs per case between € 5449.- and € 27,245.-) compared to susceptible micro-organisms [4, 5]. In the Netherlands, the group of HR-GNRs is defined as (1) Enterobacteriaceae that are Extended Spectrum Beta-Lactamase (ESBL) and/or carbapenemase positive (CPE) and/or resistant towards Fluoroquinolones and Aminoglycosides (Q&A), (2) *Acinetobacter* species that are CPE and/or resistant to Q&A, (3) *Stenotrophomonas maltophilia* resistant towards co-trimoxazole and (4) multi-resistant *Pseudomonas aeruginosa* (Table 1) [6].

Several Dutch studies reported ESBL (colonization) prevalence rates in different human populations and regions ranging from 4.7 to 10.1 % [7–12]. In addition, molecular analyses of ESBL positive isolates found in animals (veal calves, broilers and companion animals) and humans showed several associations that suggest transmission [10–14]. However, exact transmission routes and risk factors are largely unknown since epidemiological links are frequently missing, which limits the interpretation of molecular typing results. Previous studies on livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) have shown that differences in prevalence rates exist between different regions of the Netherlands, stressing the importance of regional prevalence studies which may also apply to the HR-GNR (and ESBL) epidemiology [15].

To our knowledge, the above mentioned prevalence studies are the only (recent) studies carried out in the Netherlands that determined the prevalence of ESBL-producing Enterobacteriaceae among different human populations. No studies were found that examined the prevalence and genotypes of all defined HR-GNRs among hospitalized patients (Table 1). Furthermore, the Regional

Public Health Laboratory Kennemerland (RPHLK) is a microbiological diagnostic and expertise laboratory that performs infectious disease diagnostics for primary, secondary and tertiary care facilities in the Dutch region Kennemerland allowing the possibility to perform standardized prevalence measurements from a regional point of view.

In this cross-sectional prevalence measurement, we aimed to investigate (1) the HR-GNR prevalence rate, (2) their genotypes and acquired resistance genes, and (3) associated risk factors of HR-GNR colonization among the hospitalized population in the Dutch region Kennemerland.

Methods

Ethics statement

According to the Dutch regulation for research with human subjects, neither medical or ethical approval was required to conduct the study since the data were collected as part of each hospitals standard infection control program. Additionally, we received approval to conduct the study from the institutional review board of the Spaarne Gasthuis. The data were anonymized and analyzed under code.

Study design, setting, participants, data collection and variables

Between 1 October 2013 and 31 March 2014, cross-sectional (point)prevalence measurements were performed in the three regional hospitals in the Dutch region Kennemerland as part of each hospitals infection control program. Rectal swabs were obtained from hospitalized patients (independent of the hospitalization time) at all participating wards (internal medicine, cardiology, neurology, surgery, urology, pulmonology, intensive care unit, pediatrics, geriatrics, orthopedics and gynecology). Outpatients as well as patients on day care were excluded.

Additionally, the following data were obtained from the hospital and laboratory information system in order to identify possible risk factors: (1) basic patient characteristics (gender and age), (2) antibiotic usage (during current admission and up-to six months before admission), (3) admission information (during current admission and up to 1 year before admission) and (4) historic HR-GNR isolates (since January 2008).

Laboratory detection HR-GNRs

Rectal swabs (Copan eSwab including 1 mL of modified liquid amies) were analyzed for the presence of HR-GNRs at the RPHLK by direct culturing on both an ESBL screening agar (ChromID ESBL-ID, bioMérieux, enriched with a mixture of antibiotics, including cefpodoxime) and a CLED GM20 agar (with 20 mg/L gentamicin, Oxoid). Gram-negative rods growing on

Table 1 Dutch HR-GNR definition

Organism	ESBL	CAR	QUI	AMG	CFT	PIP	COT
Enterobacteriaceae	A	A	B	B	-	-	-
<i>Acinetobacter</i> species	-	A	B	B	-	-	-
<i>Stenotrophomonas maltophilia</i>	-	-	-	-	-	-	A
<i>Pseudomonas aeruginosa</i>	-	C	C	C	C	C	-

A: This type of resistance mechanism or resistance against this antimicrobial agent indicates a HR-GNR

B: Resistance against minimal two antimicrobial agents indicates a HR-GNR

C: Resistance against minimal three antimicrobial agents indicates a HR-GNR
ESBL extended spectrum beta lactamase, CAR carbapenems, QUI fluoroquinolones, AMG aminoglycosides

CFT ceftazidime, PIP piperacillin, COT co-trimoxazole

these two agars were identified using MALDI-TOF (Bruker Daltonics, Germany). Antibiotic susceptibility testing was performed using the automated system VITEK2 (bioMérieux, France). All isolates suspected for the production of ESBL were confirmed using the combination disk method (ceftazidime and cefotaxime or cefepime with and without clavulanic acid) [16]. Strains suspected for carbapenemase production were confirmed using the modified Hodge test [16]. All positive isolates were stored at -80°C .

Molecular characterization of HR-GNR positive isolates by whole genome sequencing

DNA was extracted using the UltraClean microbial DNA isolation kit (Mo Bio Laboratories, Carlsbad, CA, USA) according to the manufacturer's protocol. A DNA library was prepared using the Nextera XT kit (Illumina, San Diego, CA, USA) according to the manufacturer's instructions. Subsequently, Whole Genome Sequencing (WGS) was performed using the MiSeq instrument (Illumina) for generating paired-end 250-bp reads, aiming at a coverage of at least 60-fold. *De novo* assembly was performed as described previously using CLC Genomics Workbench v7.0.3 (CLC bio A/S, Aarhus, Denmark) after quality trimming ($Q_s \geq 28$) with optimal word sizes based on the maximum N50 value [17, 18]. The sequence type (ST) was identified by uploading the assembled genomes to the multilocus sequence type (MLST) server (version 1.7) and the acquired resistance genes were determined with the CGE Resfinder 1.2 tool [19, 20]. STs previously undescribed were submitted to *Stenotrophomonas maltophilia* MLST database (<http://pubmlst.org/smaltophilia/>), to the *Enterobacter cloacae* MLST database (<http://pubmlst.org/ecloacae/>) or to the Enterobase database (<http://enterobase.warwick.ac.uk/>).

Infection prevention policy in the participating hospitals

The infection prevention policy in the three regional hospitals is based on the Dutch Working party of Infection prevention (WIP), which is considered highly effective and widely accepted by all Dutch hospitals in order to prevent nosocomial spread of Multi Drug Resistant Micro-Organisms (MDROs) [6]. The hospitals infection prevention policy includes the training of hospital staff (such as not wearing hand jewelry and the daily use of clean hospital uniforms). Patient rooms are daily cleaned and disinfected when indicated. As part of the policy, patients with an increased risk of HR-GNR acquisition, such as admission in a hospital abroad are screened before admission and nursed in contact isolation, consisting of: (1) nursing in a single room (2) the use of protective clothing, including gloves and gown and/or mask (when indicated), (3) compliance to hand hygiene protocols and (4) disinfection of medical

devices and equipment after use. In addition, an alert pop-up is entered in the hospital information system as warning when HR-GNR positive patients are readmitted to the hospital. In case of an unexpected HR-GNR positive patient, all contacts (patients that were still hospitalized and shared the same room with the index patient for at least 24 h during the current hospital stay) were screened for HR-GNR colonization and isolated (when positive) as described.

Outcomes and data analysis

The primary outcome was the rectal HR-GNR prevalence rate at patient level. These prevalence rates were calculated by dividing the number of HR-GNR positive patients (and per HR-GNR subgroup) by the total number of sampled patients. Confidence intervals (95 %) of the prevalence rates were calculated using the Wilson score [21]. Proportions (such as prevalence rates and type of micro-organism) were compared between hospitals using the Pearson chi-square test or Fisher's exact test (when appropriate).

The association between several independent variables (sex, age, historic antibiotic use, antibiotic use during admission, historic admission (up to 1 year), known HR-GNR positive in the past (since January 2008) and time from start admission to sampling) and the dependent variable HR-GNR positivity (yes/no) was examined using logistic regression models. These associations were presented as odds ratio's, including 95 % confidence intervals and p -values. The independent continuous variables were first checked for linearity and when not linear associated reported as quartiles. Univariate significant associations were further analyzed using a multivariate logistic regression model. All analyses were performed using PASW SPSS Statistics version 18.0. Results were interpreted as statistically significant when the p -value was < 0.05 .

Results

Study population

In total, out of 566 eligible patients, 427 patients (75.4 %) were sampled during this present regional prevalence measurement. The mean age (SD) of this study population was 65.1 (21.1) years and 217 patients (50.8 %) were male. The median time (range) between admission and sampling was 3 (0–48) days. From all sampled patients, twelve (2.8 %) patients were known to be HR-GNR positive in the past (since January 2008) of whom three (25.0 %) patient(s) were positive and nine (75.0 %) were negative in the present prevalence measurement. The median time (range) between the first HR-GNR positive isolate and the current prevalence measurement for the three positive and nine negative patients was 414 (69–1991) and 648 (28–2273) days, respectively. A

total of 167 (39.3 %) patients used antibiotics during the present hospital stay and 99 (23.3 %) patients used antibiotics up to 6 months before the present hospital stay. A total of 204 patients (48.0 %) were admitted (up to 1 year) before the present hospital stay. Stratified study population characteristics for the separate hospitals are shown in Table 2. From two patients, no demographic and historic data were available and these were excluded from further analyses.

Prevalence of HR-GNRs and micro-organisms in hospitalized patients

A total of 22 patients were culture positive for one or more types of HR-GNRs, resulting in a regional HR-GNR colonization prevalence (95 % CI) of 5.2 % (3.6–7.9). Of these 22 positive patients, 15 were ESBL positive (3.5 % (2.1–5.7)), 7 patients were positive with an isolate resistant towards Q&A (1.6 % (0.8–3.3)) and from one patient (0.2 % (0–1.3)) a *Stenotrophomonas maltophilia* resistant towards co-trimoxazole was isolated. No CPE, multi-resistant *Acinetobacter* species or multi-resistant *Pseudomonas aeruginosa* were isolated. From three patients (0.7 %), more than one distinctive HR-GNR phenotype was isolated of whom one patient (0.2 %) was positive for more than one type of HR-GNR (both an ESBL positive isolate and isolate resistant towards Q&A). Stratified prevalence rates for each hospital and type of HR-GNR are shown in Table 3. No statistically significant differences in prevalence rates were found between hospitals ($p = 0.180$). When patients were divided in two groups based on hospitalization time, a HR-GNR prevalence (95 % CI) of 5.3 % (2.6–10.5) and 4.7 % (2.9–7.9) was found respectively for patients that were 0–1 day hospitalized (7 out of 133 patients) and longer than one day hospitalized (14 out of 292 patients) ($p = 0.836$).

Most of the isolated HR-GNR micro-organisms were *Escherichia coli* (73.1 %) followed by *Enterobacter cloacae*

(11.5 %), *Klebsiella pneumoniae* (11.5 %) and *Stenotrophomonas maltophilia* (3.8 %). There was no significant difference in the total distribution of all micro-organisms between the three hospitals ($p = 0.421$).

Molecular characterization of ESBL positive isolates

Twenty-four HR-GNR positive isolates belonging to 22 HR-GNR positive patients were genotyped using WGS. All 16 phenotypically ESBL positive isolates sampled from 15 patients harbored ESBL genes of which one isolate (*Klebsiella pneumoniae*, ST48) contained two ESBL genes ($bla_{CTX-M-15}$ and bla_{SHV-11}). ESBL genes found were $bla_{CTX-M-1}$ ($n = 4$, 25.0 %), $bla_{CTX-M-15}$ ($n = 3$, 18.8 %), $bla_{CTX-M-27}$ ($n = 2$, 12.5 %), $bla_{CTX-M-14b}$ ($n = 2$, 12.5 %), $bla_{CTX-M-9}$ ($n = 2$, 12.5 %), $bla_{CTX-M-14}$ ($n = 1$, 6.3 %), $bla_{CTX-M-3}$ ($n = 1$, 6.3 %), bla_{SHV-11} ($n = 1$, 6.3 %) and bla_{SHV-12} ($n = 1$, 6.3 %) (Table 4).

Eleven of the 16 ESBL positive isolates were determined as *Escherichia coli* (68.8 %). ST131 was found twice (12.5 %) and both isolates harbored the $bla_{CTX-M-27}$ gene. All other ESBL positive *Escherichia coli* sequence types were found once ($n = 1$, 6.3 %) and included ST10, ST69, ST88, ST93, ST349, ST635, ST685, ST2178 and a new sequence type (ST5929). Three of the 16 ESBL positive isolates were determined as *Klebsiella pneumoniae* (18.8 %) consisting of ST22, ST45 and ST48. Additionally, all three ESBL positive *Klebsiella pneumoniae* isolates harbored the $bla_{CTX-M-15}$ gene. The other three ESBL positive isolates were determined as *Enterobacter cloacae* (18.8 %) of which ST50 was found twice and a new sequence type (ST421) once. Two of the three ESBL positive *Enterobacter cloacae* isolates harbored the $bla_{CTX-M-9}$ gene. In addition, Table 4 also shows phenotypic resistance patterns and identified acquired resistance genes towards Q&A, revealing that the presence of acquired resistance genes in an isolate and phenotypic resistance do not always co-occur.

Table 2 Demographic characteristics of patients

Patient characteristics	Total ^a	Hospital 1	Hospital 2	Hospital 3 ^a
Number of unique patients	427 (100)	126 (29.5)	167 (39.1)	134 (31.4)
Sex				
Male	217 (51.1)	57 (45.2)	90 (53.9)	70 (53.0)
Used antibiotics 6 months before current admission	99 (23.3)	41 (32.5)	34 (20.4)	24 (18.2)
Used antibiotics during current admission	167 (39.3)	33 (26.2)	76 (45.5)	58 (43.9)
Admitted before current admission (up to 1 year)	204 (48.0)	65 (51.6)	77 (46.1)	62 (47.0)
Known HR-GNR positive in the past ^b	12 (2.8)	2 (1.6)	7 (4.2)	3 (2.3)
Mean age in years (SD)	65.1 (21.1)	61.2 (22.9)	65.7 (22.4)	68.1 (16.9)
Median time from admission to sampling in days (Range)	3.0 (0–48)	2 (0–29)	4 (0–48)	4 (0–43)

Data are presented as numbers (%) unless indicated otherwise

Percentages were calculated in reference to the specific hospital

^aFrom two patients no demographic characteristics were known (in total: 425 patients with known characteristics and 132 for hospital 3)

^bKnown HR-GNR positive in the past (since January 2008)

Table 3 Prevalence rates of rectal HR-GNRs colonization on a patient level

	Regional		Hospital 1		Hospital 2		Hospital 3	
	Prevalence	95 % CI	Prevalence	95 % CI	Prevalence	95 % CI	Prevalence	95 % CI
HR-GNR	5.2 % (22/427)	3.6 %–7.9 %	2.4 % (3/126)	0.8 %–6.7 %	7.2 % (12/167)	4.6 %–12.8 %	5.2 % (7/134)	2.6 %–10.4 %
ESBL	3.5 % (15/427)	2.1 %–5.7 %	1.6 % (2/126)	0.4 %–5.6 %	5.4 % (9/167)	2.8 %–9.9 %	3.0 % (4/134)	1.2 %–7.4 %
Q&A	1.6 % (7/427)	0.8 %–3.3 %	0.8 % (1/126)	0.1 %–4.3 %	2.4 % (4/167)	0.9 %–6.0 %	1.5 % (2/134)	0.4 %–5.3 %
Other ^a	0.2 % (1/427)	0.0 %–1.3 %	0.0 % (0/126)	0.0 %–0.0 %	0.0 % (0/167)	0.0 %–0.0 %	0.7 % (1/134)	0.1 %–3.4 %

^aAll other HR-GNR beside ESBL and Q&A, see Table 1

Table 4 Molecular characteristics of HR-GNR positive isolates

Patient	Hospital	Species	MLST	HR-GNR type ^{a, b, c}	ESBL gene(s)	Acquired AMG ^d resistance genes	Acquired QUI ^e resistance genes	AMG ^d resistance ^f	QUI ^e resistance ^f
P1	1	<i>E. coli</i>	ST685	ESBL	CTX-M-1	-	-	S	S
P2	1	<i>K. pneumoniae</i>	ST48	ESBL	CTX-M-15, SHV-11	strA, strB	oqxA, oqxB, QnrB66	S	S
P3	1	<i>E. coli</i>	ST69	Q&A	-	strA, strB, aph(3')-Ia, aadA5, aac(3)-IId	-	R	R
P4	2	<i>E. coli</i>	ST5929 ^g	ESBL	CTX-M-1	-	-	S	S
P5	2	<i>E. coli</i>	ST69	ESBL	CTX-M-1	aadA5	-	S	S
P6	2	<i>E. coli</i>	ST2178	ESBL	CTX-M-3	-	-	S	S
		<i>E. coli</i>	ND	ESBL	ND	ND	ND	S	S
		<i>K. pneumoniae</i>	ST22	ESBL	CTX-M-15	strA, strB, aph(3')-Ic, aac(3)-Iia, aac(6')-Ib-cr	oqxA, oqxB, aac(6')-Ib-cr, QnrB66	R	S
P7	2	<i>E. coli</i>	ST349	ESBL	CTX-M-14b	strA, strB	-	S	S
P8	2	<i>E. coli</i>	ST93	ESBL	CTX-M-14b	-	-	S	R
P9	2	<i>E. coli</i>	ST131	ESBL	CTX-M-27	strA, strB, aadA5	-	S	R
P10	2	<i>E. cloacae</i>	ST50	ESBL	CTX-M-1	aadA2, aadB	oqxA, oqxB, QnrA1	I	R
P11	2	<i>E. cloacae</i>	ST421 ^g	ESBL	CTX-M-9	aadB	oqxA, oqxB, QnrA1	S	S
P12	2	<i>E. cloacae</i>	ST50	ESBL	CTX-M-9	aadA2, aadB	oqxA, oqxB, QnrA1	I	R
		<i>E. coli</i>	ST88	Q&A	-	aph(3')-Ic	-	R	R
P13	2	<i>E. coli</i>	ST69	Q&A	-	strA, strB, aac(3)-IId	-	R	R
P14	2	<i>E. coli</i>	ST131	Q&A	-	strA, aac(3)-IId, aadA5	-	R	R
P15	2	<i>E. coli</i>	ST648	Q&A	-	aac(6')-Ib-cr	aac(6')-Ib-cr	R	R
P16	3	<i>E. coli</i>	ST10	ESBL	CTX-M-14	strA, strB, aac(3)-IId, aadA1, aadA4, aac(6')-Ib-cr	aac(6')-Ib-cr	R	S
		<i>K. pneumoniae</i>	ST45	ESBL	CTX-M-15	strA, strB, aac(3)-Ila, aac(6')-Ib-cr	oqxA, oqxB, aac(6')-Ib-cr, QnrB66	R	I
P18	3	<i>E. coli</i>	ST131	ESBL	CTX-M-27	-	-	S	R
P19	3	<i>E. coli</i>	ST635	ESBL	SHV-12	strA, strB, aac(3)-IIb, aacA4, aac(6')-Ib-cr, aac(6')-IIc	aac(6')-Ib-cr, QnrB4, QnrB66	R	I
		<i>E. coli</i>	ST57	Q&A	-	aph(4)-Ia, aac(3)-IVa, aadA2	-	R	R
P20	3	<i>E. coli</i>	ST57	Q&A	-	aph(4)-Ia, aac(3)-IVa, aadA2	-	R	R
		<i>E. coli</i>	ND	Q&A	-	ND	ND	R	R
P21	3	<i>E. coli</i>	ST131	Q&A	-	strA, strB, aac(3)-IId, aadA5	-	R	R
P22	3	<i>S. maltophilia</i>	ST152 ^g	CTX-R	-	aph(3')-IIc, aacA4, aac(6')-Ib-cr	aac(6')-Ib-cr	ND	ND

^a: ESBL: Extended Spectrum Beta Lactamase; ^b: Q&A: combined resistance towards Fluoroquinolones and Aminoglycosides; ^c: CTX-R: resistance towards Co-trimoxazole; ^d: QUI: Fluoroquinolones; ^e: AMG: Aminoglycosides; ^f: resistance based on VITEK2 results; S sensitive, I intermediate, R resistant, ND not determined, NA not available, ^g new sequence type

Molecular characterization of non-ESBL HR-GNR positive isolates

Eight (33.3 %) of the 24 genotyped HR-GNR positive isolates (belonging to eight patients) were non-ESBL isolates. Seven of these eight isolates showed combined resistance towards Q&A. All seven isolates resistant towards Q&A were determined as *Escherichia coli* of which ST69 and ST131 were both found twice (28.6 %). All other sequence types (ST57, ST88, and ST648) were recovered once (14.3 %). Only one isolate resistant towards Q&A (ST648) harbored the acquired *aac(6')* *Ib-cr* gene encoding for resistance towards both Fluoroquinolones and Aminoglycosides. The other isolates resistant towards Q&A harbored only acquired resistance genes encoding for Aminoglycoside resistance although they also showed phenotypic resistance towards Fluoroquinolones. These acquired Aminoglycoside resistance genes were *strA* ($n = 4$, 57.1 %), *strB* ($n = 3$, 42.9 %), *aac(3) -IId* ($n = 3$, 42.9 %), *aadA5* ($n = 3$, 42.9 %), *aph(3')* *-Ia* ($n = 1$, 14.3 %), *aph(4) -Ia* ($n = 1$, 14.3 %), *aac(3) -IVa* ($n = 1$, 14.3 %), *aadA2* ($n = 1$, 14.3 %) and *aph(3')* *-Ic* ($n = 1$, 14.3 %).

One of the eight non-ESBL HR-GNR positive isolates was determined as *Stenotrophomonas maltophilia*, which showed resistance towards co-trimoxazole (12.5 %) and was characterized as ST152 (new sequence type). This isolate harbored the acquired *sul1* gene encoding for Sulphonamide resistance.

Associated risk factors for being HR-GNR positive in hospitalized patients

Table 5 shows the associations between independent variables and the dependent variable (HR-GNR positive in the prevalence measurement). Both continuous variables (age and time from admission to sampling) were reported as quartiles (in reference to the first category) since they were not linearly associated with the outcome. Being known HR-GNR positive in the past was the only significant associated risk factor, odds ratio (95 % CI): 7.32 (1.82–29.35), p -value = 0.005. The other tested risk factors were associated with an increased risk of HR-GNR positivity, although not significant. The association with being HR-GNR positive in the past did not markedly change after adjustment for possible confounders (antibiotic use (during the current and up to 6 months before the current admission) age, sex, admission before the current admission and time from start admission to sampling) using multivariate logistic regression analysis, odds ratio (95 % CI): 6.54 (1.35–31.61), p -value = 0.020 (Table 6). No significant effect modifiers were identified.

Discussion

The present study shows the results of a cross-sectional HR-GNR (including ESBL) prevalence measurement in hospitalized patients of three hospitals in the Dutch region Kennemerland (with 650,000 inhabitants). In total, 427 rectal swabs derived from unique patients in these

Table 5 Univariate associations between possible risk factors in clinical patients and being HR-GNR positive

Risk factor	HR-GNR positive patients ($n = 21$) ^a	HR-GNR negative patients ($n = 404$) ^a	Odds ratio (95 % CI)	P -value
Sex				
Female	9 (42.9)	199 (49.3)	1 (ref)	-
Male	12 (57.1)	205 (50.7)	1.29 (0.53–3.14)	0.568
Used antibiotics 6 months before current admission	6 (28.6)	93 (23.0)	1.34 (0.51–3.55)	0.559
Used antibiotics during admission	10 (47.6)	157 (38.9)	1.43 (0.59–3.45)	0.425
Admitted before current admission (up to 1 year)	14 (66.7)	190 (47.0)	2.25 (0.89–5.70)	0.086
Known HR-GNR positive in the past	3 (14.3)	9 (2.2)	7.32 (1.82–29.35)	0.005
Age (years)				
Group 1 (0–56 years)	4 (19.0)	103 (25.5)	1 (ref)	-
Group 2 (57–70 years)	5 (23.8)	99 (24.5)	1.30 (0.34–4.98)	0.701
Group 3 (71–79 years)	9 (42.9)	101 (25.0)	2.30 (0.69–7.69)	0.178
Group 4 (80–94 years)	3 (14.3)	101 (25.0)	0.77 (0.17–3.50)	0.730
Time from admission to sampling (days)				
Group 1 (0–1 days)	7 (33.3)	126 (31.2)	1 (ref)	-
Group 2 (2–3 days)	3 (14.3)	90 (22.3)	0.60 (0.15–2.38)	0.468
Group 3 (4–6 days)	3 (14.3)	85 (21.0)	0.64 (0.16–2.53)	0.519
Group 4 (7–48 days)	8 (38.1)	103 (25.5)	1.40 (0.49–3.98)	0.531

^aIn total, data for 425 patients were available for analyses, as for two patients demographic data were unknown

Table 6 Multivariate logistic regression model between possible risk factors in clinical patients

Risk factor	Unadjusted association		Fully adjusted model ^a	
	Odds ratio (95 % CI)	P-value	Odds ratio (95 % CI)	P-value
Sex	1.29 (0.53–3.14)	0.568	1.12 (0.43–2.91)	0.823
Used antibiotics 6 months before current admission	1.34 (0.51–3.55)	0.559	0.84 (0.28–2.51)	0.756
Used antibiotics during admission	1.43 (0.59–3.45)	0.425	1.15 (0.41–3.23)	0.791
Admitted before current admission (up to 1 year)	2.25 (0.89–5.70)	0.086	2.21 (0.80–6.10)	0.126
Known HR-GNR positive in the past	7.32 (1.82–29.35)	0.005	6.54 (1.35–31.61)	0.020
Age (years)				
Group 1 (0–56 years)	1 (ref)	-	1 (ref)	-
Group 2 (57–70 years)	1.30 (0.34–4.98)	0.701	0.84 (0.20–3.45)	0.806
Group 3 (71–79 years)	2.30 (0.69–7.69)	0.178	1.73 (0.49–6.15)	0.395
Group 4 (80–94 years)	0.77 (0.17–3.50)	0.730	0.60 (0.12–2.93)	0.530
Time from admission to sampling (days)				
Group 1 (0–1 days)	1 (ref)	-	1 (ref)	-
Group 2 (2–3 days)	0.60 (0.15–2.38)	0.468	0.47 (0.11–2.01)	0.311
Group 3 (4–6 days)	0.64 (0.16–2.53)	0.519	0.52 (0.12–2.29)	0.385
Group 4 (7–48 days)	1.40 (0.49–3.98)	0.531	1.10 (0.34–3.62)	0.870

^aCorrected for (1) Antibiotic use during current admission, (2) Antibiotic use up 6 months before current admission, (3) sex, (4) age, (5) admitted before current admission, (6) time from start admission to sampling and (7) known HR-GNR positive in the past

hospitals were analyzed and resulted in a total HR-GNR and ESBL prevalence rate of 5.2 and 3.5 %, respectively. Furthermore, 7 patients (1.6 %) were positive with an isolate resistant towards Q&A, and from one patient (0.2 %) a *Stenotrophomonas maltophilia* resistant towards co-trimoxazole was isolated. In line with other Dutch prevalence studies, no CPE positive bacteria were found, indicating a relatively low prevalence in our region.

Several Dutch studies have previously reported about the ESBL prevalence rates among different human populations. First, Overdeest et al. found a rectal ESBL colonization prevalence of 4.9 % within hospitalized patients [7]. Second, a prevalence survey among 125 residents living in five nursing homes and two (hospital) rehabilitation wards in the central region of the Netherlands showed an ESBL prevalence of 6.0 % [8]. Third, a study among 720 patients with gastrointestinal complaints visiting the general practitioner showed an ESBL prevalence of 10.1 % [9]. Fourth, two studies reported on the ESBL prevalence among travelers (before travel) resulting in an ESBL prevalence rate of 8.6 and 9.0 % [22, 23]. Fifth, a cross-sectional ESBL prevalence measurement performed in a representative sample of the Dutch community population showed an ESBL prevalence of 4.7 % [24]. A comparison between the ESBL prevalence rate in our present study (3.5 %) and the other studies showed that the ESBL prevalence rate in our region was relatively low. A possible explanation for this difference could be the different culture methods that were used. Some studies used an (selective

or non-selective) enrichment broth, which is associated with higher sensitivities compared with direct culture methods [25, 26]. Furthermore, differences in population characteristics or a geographical variation in risk factors may also explain the differences in ESBL prevalence rates.

All phenotypic characterized ESBL positive isolates in our study harbored ESBL genes, mostly *bla*_{CTX-M} (88.2 %). WGS showed that the *bla*_{CTX-M-1} (25.0 %), *bla*_{CTX-M-15} (18.8 %), *bla*_{CTX-M-14b} and *bla*_{CTX-M-9} (both 12.5 %) ESBL genes were found most often. In line with our results, another study among hospitalized patients isolated the *bla*_{CTX-M-1} (45.8 %) ESBL gene most often, followed by *bla*_{CTX-M-15} (16.7 %) and *bla*_{TEM-52} (12.5 %) [7]. Surprisingly, a study among a representative sample of the Dutch community population also isolated the ESBL gene *bla*_{CTX-M-1} (35 %), *bla*_{CTX-M-15} (33 %) and *bla*_{CTX-M-14} (18 %) most often, showing that isolates obtained from hospitalized and non-hospitalized individuals share similar ESBL genes [24]. This finding suggests that the positive HR-GNR patients may have already been positive at admission and act as a reservoir for other patients. In addition, studies among cats, dogs and chicken (retail) meat isolated the *bla*_{CTX-M-1} most frequently, indicating that shared reservoirs and/or transmission dynamics exist [11, 12]. Some studies showed other distributions in ESBL genes. A study among symptomatic general practitioner patients with gastrointestinal complaints, found predominantly *bla*_{CTX-M-15} (47 %) ESBL-genes [9]. Furthermore, the two studies that investigated the ESBL prevalence among healthy

travelers (before travel) found predominantly *bla*_{CTX-M-15} (47 %) and *bla*_{CTX-M-9} (90 %) ESBL genes [22, 23]. These differences may be explained by heterogeneity in study populations. Travelers, symptomatic general practitioner patients and hospitalized patients possibly show different risk behavior, which may be reflected in the molecular typing results. More research is needed to further elucidate these population differences. As shown, most Dutch ESBL prevalence studies report predominantly *bla*_{CTX-M} ESBL genes and a great variation in *Escherichia coli* sequence types that is in line with our results.

As shown in our study, 6 out of 7 phenotypically characterized isolates resistant towards Q&A had no acquired resistance genes encoding for quinolone resistance, suggesting that quinolone resistance in these isolates was mainly caused by chromosomal mutations. The same was seen for ESBL positive isolates that showed phenotypical resistance towards fluoroquinolones or aminoglycosides (not both). As described earlier by Guan et al., plasmid-encoded quinolone resistance genes do not confer quinolone resistance by themselves, but augment the effect of other resistance mutations [27]. Probably the same conclusion is applicable to acquired Aminoglycoside resistance genes and phenotypic expression. More research is needed to elucidate the role of these resistance genes and phenotypic expression.

Traveling (predominantly to South Asia) and having a high degree of contact with broilers are today's most important identified risk factors for colonization with ESBL positive bacteria [12, 22, 23, 28, 29]. In our study, we analyzed several potential risk factors for HR-GNR colonization. Antibiotic use (during and/or 6 months before admission) did show a higher odds ratio for HR-GNR positivity (including ESBL), although this association was not significant. The same was found for sex (higher odds for males), 'admission before the current admission (up to one year)' and age (for the first three quartiles). Being known HR-GNR positive in the past was the only significantly associated risk factor, also when corrected for possible confounders. Additionally, studies carried out among long term rehabilitation patients also identified 'being known HR-GNR positive' as independent risk factor for HR-GNR colonization [30, 31]. This finding suggests that colonization with HR-GNRs persists for a longer period and should not be ignored at re-admission at the hospital. Additional longitudinal studies are needed to quantify the influence of the period of HR-GNR colonization since this information is essential to assess the importance of isolating patients at readmission [6]. Another studied risk factor, time to sampling (from start admission to sampling) was not significantly associated with HR-GNR colonization, indicating that the role of nosocomial colonization was minimal. This finding

suggests that basic hygiene in the studied hospitals is good. Even when we excluded the first quartile (0–1 day), as one may hypothesize that HR-GNR positive patients in this group were already positive at admission, no significant association was detected. To elucidate the influence of this possible risk factor, prospective study designs are needed in which all patients are screened at admission and during hospitalization (at specific time intervals).

Furthermore, we suggest that future studies aiming at identifying transmission routes and/or reservoirs for HR-GNR involving pigs, horses or other (companion) animals, should include both humans and animals (and their isolates) that are epidemiologically linked. When such studies are performed in a longitudinal design, transmission dynamics as well as origin and transmission of resistance genes can be studied (from humans to animals or the other way around).

To our knowledge, no studies are available that determined the colonization prevalence of Q&A resistant Enterobacteriaceae. We recommend for future studies to incorporate all defined HR-GNRs, in addition to ESBLs, in prevalence studies, in order to obtain a more comprehensive overview of colonization with HR-GNRs.

The present study has several limitations. First, only a small number of potential risk factors were included in the risk factor analysis lacking the ability to identify more 'potential' risk factors such as contact with HR-GNR positive household members, travel history, food preferences, other medication use, having pets and stay or transfer from a nursing home. Second, as mentioned before, no enrichment broth was used which could have underestimated the HR-GNR prevalence rate. Third, historic data on antibiotic use were retrospectively retrieved from the hospital pharmacy database, which only contains data on clinical prescribed antibiotics, and not on antibiotic use in the primary care setting. Fourth, our study was performed in a single region of the Netherlands with a relatively small number of cases limiting the power to identify all important risk factors. Therefore, our results should be interpreted with caution.

Conclusion

In conclusion, no local differences in HR-GNR prevalence rates and micro-organisms were found between the three regional hospitals. In addition, the Dutch region Kennemerland showed similar ESBL prevalence rates among the hospitalized patients population in comparison to other Dutch regions. When previously HR-GNR positive patients are readmitted they should be screened for HR-GNR colonization since colonization with GR-GNRs could be prolonged. Molecular typing results showed that comparable ESBL genotypes were found as earlier described in both humans and animals supporting the hypothesis of multiple reservoirs

and risk factors. Future studies must focus more on these postulated reservoirs and risk factors with appropriate study designs.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DS, SME, BLH, BD, PH, MS, IE, JK and JWB designed the study, interpreted the results, revised the manuscript and wrote the manuscript. DS and SME performed the statistical analysis. DS, SME and BLH interpreted the results from the statistical analysis. PH, MS and IE collected patient specimens and data. JWR performed the molecular analysis of the bacterial isolates, interpreted the molecular typing results, revised the manuscript and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We are grateful to Sigrd Rosema for her help in analyzing the WGS data. We thank the team of the curators of the University of Freiburg (Germany) for importing novel alleles, profiles and isolates at the *Stenotrophomonas maltophilia* MLST website (<http://pubmlst.org/smaltophilia/>), the team of the curators of the National Center for Global Health and Medicine (Japan) for importing novel alleles, profiles and isolates at the *Enterobacter cloacae* MLST website (<http://pubmlst.org/ecloacae/>) and the team of the curators of the Warwick Medical School (United Kingdom) for importing novel alleles, profiles and isolates at the Enterobase website (<http://enterobase.warwick.ac.uk/>).

Author details

¹Department of Epidemiology and Infection Prevention, Regional Public Health Laboratory Kennemerland, Boerhaavelaan 26, 2035 RC Haarlem, The Netherlands. ²Department of Infection Prevention, Spaarne Gasthuis, Haarlem, The Netherlands. ³Department of Infection Prevention, Rode Kruis Ziekenhuis, Beverwijk, The Netherlands. ⁴Laboratory for Microbiology and Infection Control, Amphia Hospital, Breda, and University Medical Center, Utrecht, The Netherlands. ⁵Department of Medical Microbiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

Received: 4 February 2016 Accepted: 3 March 2016

Published online: 08 March 2016

References

- Willemsen I, Elberts S, Verhulst C, Rijnsburger M, Filius M, Savelkoul P, et al. Highly resistant gram-negative microorganisms: incidence density and occurrence of nosocomial transmission (TRIANGLE study). *Infect Control Hosp Epidemiol*. 2011;32(4):333–41.
- Falagas ME, Karageorgopoulos DE. Extended-spectrum beta-lactamase-producing organisms. *J Hosp Infect*. 2009;73(4):345–54.
- Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. *Clin Microbiol Rev*. 2005;18(4):657–86.
- Rottier WC, Ammerlaan HS, Bonten MJ. Effects of confounders and intermediates on the association of bacteraemia caused by extended-spectrum β -lactamase-producing Enterobacteriaceae and patient outcome: a meta-analysis. *J Antimicrob Chemother*. 2011;67(6):1311–20.
- Cosgrove SE. The relationship between antimicrobial resistance and patient outcomes: mortality, length of hospital stay, and health care costs. *Clin Infect Dis*. 2006;42(2):S82–9.
- Bijzonder resistente micro-organismen (BRMO). Werkgroep Infectie Preventie (WIP). 2012. <http://www.nvmm.nl/system/files/121217%20BRMO-%20final.pdf>. Accessed 15 Jul 2015.
- Overdeest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, et al. Extended-Spectrum Beta-Lactamase genes of *Escherichia coli* in chicken meat and humans, the Netherlands. *Emerg Infect Dis*. 2011;17(7):1216–22.
- Hoogendoorn M, Smalbrugge M, Stobberingh EE, van Rossum SV, Vlamincx BJ, Thijsen SF. Prevalence of antibiotic resistance of the commensal flora in Dutch nursing homes. *J Am Med Dir Assoc*. 2013;14(5):336–9.
- Reuland EA, Overdeest IV, Al Naiemi N, Kalpoe JS, Rijnsburger MC, Raadsen SA, et al. High prevalence of ESBL-producing Enterobacteriaceae carriage in Dutch community patients with gastrointestinal complaints. *Clin Microbiol Infect*. 2012;19(6):542–9.
- Hordijk J, Wagenaar JA, Kant A, van Essen-Zandbergen A, Dierikx C, Veldman K, et al. Cross-sectional study on prevalence and molecular characteristics of plasmid mediated ESBL/AmpC-producing *Escherichia coli* isolated from veal calves at slaughter. *PLoS One*. 2013;8(5):e65681.
- Huijbers PM, Graat EA, Haenen AP, van Santen MG, van Essen-Zandbergen A, Mevius DJ, et al. Extended-spectrum and AmpC β -lactamase-producing *Escherichia coli* in broilers and people living and/or working on broiler farms: prevalence, risk factors and molecular characteristics. *J Antimicrob Chemother*. 2014;69(10):2669–75.
- Hordijk J, Schoormans A, Kwakernaak M, Duim B, Broens E, Dierikx C, et al. High prevalence of fecal carriage of extended spectrum β -lactamase/AmpC-producing Enterobacteriaceae in cats and dogs. *Front Microbiol*. 2013;4:242.
- Wu G, Day MJ, Mafura MT, Nunez-Garcia J, Fenner JJ, Sharma M, et al. Comparative analysis of ESBL-positive *Escherichia coli* isolates from animals and humans from the UK, The Netherlands and Germany. *PLoS One*. 2013;8(9):e75392.
- Ewers C, Bethe A, Semmler T, Guenther S, Wieler LH. Extended-spectrum β -lactamase-producing and AmpC-producing *Escherichia coli* from livestock and companion animals, and their putative impact on public health: a global perspective. *Clin Microbiol Infect*. 2012;18(7):646–55.
- van Rijen MM, Van Keulen PH, Kluytmans JA. Increase in a Dutch hospital of methicillin-resistant *Staphylococcus aureus* related to animal farming. *Clin Infect Dis*. 2008;46(2):261–3.
- Guideline Laboratory detection of highly resistant microorganisms, version 2.0. NVMM. 2013. <http://www.nvmm.nl/system/files/2012.11.15%20richtlijn%20BRMO%20%28version%202.0%29%20-%20RICHTLIJN.pdf>. Accessed 15 Jul 2015.
- Zhou K, Ferdous M, de Boer RF, Kooistra-Smid AM, Grundmann H, Friedrich AW, et al. The mosaic genome structure and phylogeny of Shiga toxin-producing *Escherichia coli* O104:H4 is driven by short-term adaptation. *Clin Microbiol Infect*. 2015;21(5):468. e7-468.e18.
- Ferdous M, Zhou K, Mellmann A, Morabito S, Croughs PD, de Boer RF, et al. Is Shiga Toxin-Negative *Escherichia coli* O157:H7 Enteropathogenic or Enterohemorrhagic *Escherichia coli*? Interim molecular analysis using whole-genome sequencing. *J Clin Microbiol*. 2015;53(11):3530–8.
- Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, et al. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol*. 2012;50(4):1355–61.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*. 2012;67(11):2640–4.
- Wilson EB. Probable inference, the law of succession, and statistical inference. *J Am Stat Assoc*. 1927;22:209–12.
- Paltansing S, Vlot JA, Kraakman ME, Mesman R, Bruijning ML, Bernards AT, et al. Extended-spectrum β -lactamase-producing enterobacteriaceae among travelers from the Netherlands. *Emerg Infect Dis*. 2013;19(8):1206–13.
- von Wintersdorff CJ, Penders J, Stobberingh EE, Oude Lashof AM, Hoebe CJ, Savelkoul PH, et al. High rates of antimicrobial drug resistance gene acquisition after international travel, The Netherlands. *Emerg Infect Dis*. 2014;20(4):649–57.
- van Hoek AH, Schouls L, van Santen MG, Florijn A, de Greeff SC, van Duijkeren E. Molecular characteristics of extended-spectrum cephalosporin-resistant enterobacteriaceae from humans in the community. *PLoS One*. 2015;10(6):e0129085. doi:10.1371/journal.pone.0129085.
- Murk JL, Heddema ER, Hess DL, Bogaards JA, Vandenbroucke-Grauls CM, Debets-Ossenkopp YJ. Enrichment broth improves detection of extended-spectrum-beta-lactamase-producing bacteria in throat and rectal surveillance cultures of samples from patients in intensive care units. *J Clin Microbiol*. 2009;47(6):1885–7.
- Kluytmans-van den Bergh MF, Verhulst C, Willemsen LE, Verkade E, Bonten MJ, Kluytmans JA. Rectal carriage of extended-spectrum beta-lactamase-producing Enterobacteriaceae in hospitalized patients: selective pre-enrichment increases the yield of screening. *J Clin Microbiol*. 2015;53(8):2709–12.
- Guan X, Xue X, Liu Y, Wang J, Wang Y, Wang J, et al. Plasmid-mediated quinolone resistance—current knowledge and future perspectives. *J Int Med Res*. 2013;41(1):20–30.
- Kuenzli E, Jaeger VK, Frei R, Neumayr A, DeCrom S, Haller S, et al. High colonization rates of extended-spectrum β -lactamase (ESBL)-producing *Escherichia coli* in Swiss travellers to South Asia— a prospective observational multicentre cohort study looking at epidemiology, microbiology and risk factors. *BMC Infect Dis*. 2014;14:528.

29. Yaita K, Aoki K, Suzuki T, Nakaharai K, Yoshimura Y, Harada S, et al. Epidemiology of extended-spectrum β -lactamase producing *Escherichia coli* in the stools of returning Japanese travelers, and the risk factors for colonization. *PLoS One*. 2014;9(5):e98000.
30. Bilavsky E, Temkin E, Lerman Y, Rabinovich A, Salomon J, Lawrence C, et al. Risk factors for colonization with extended-spectrum beta-lactamase-producing enterobacteriaceae on admission to rehabilitation centres. *Clin Microbiol Infect*. 2014;20(11):O804–10.
31. Del Rosario-Quintana C, Tosco-Núñez T, Lorenzo L, Martín-Sánchez AM, Molina-Cabrillana J. Prevalence and risk factors of multi-drug resistant organism colonization among long-term care facilities in Gran Canaria (Spain). *Rev Esp Geriatr Gerontol*. 2015;50(5):232–6.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

