

University of Groningen

The dust masses of powerful radio galaxies

Tadhunter, C.; Dicken, D.; Morganti, R.; Konyves, V.; Ysard, N.; Nesvadba, N.; Ramos Almeida, C.

Published in:
Monthly Notices of the Royal Astronomical Society

DOI:
[10.1093/mnrasl/slu135](https://doi.org/10.1093/mnrasl/slu135)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Tadhunter, C., Dicken, D., Morganti, R., Konyves, V., Ysard, N., Nesvadba, N., & Ramos Almeida, C. (2014). The dust masses of powerful radio galaxies: clues to the triggering of their activity. *Monthly Notices of the Royal Astronomical Society*, 445, L51-L55. <https://doi.org/10.1093/mnrasl/slu135>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The dust masses of powerful radio galaxies: clues to the triggering of their activity

C. Tadhunter,¹★ D. Dicken,² R. Morganti,^{3,4} V. Konyves,² N. Ysard,⁵ N. Nesvadba⁵ and C. Ramos Almeida^{6,7}

¹Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, UK

²Laboratoire AIM-Paris-Saclay, CEA/DSM/Irfu, Orme des Merisiers, Bat 709, F-91191 Gif sur Yvette, France

³ASTRON, PO Box 2, NL-7990 AA Dwingeloo, the Netherlands

⁴Kapteyn Astronomical Institute, University of Groningen, PO Box 800, NL-9700 AV Groningen, the Netherlands

⁵Institut d'Astrophysique Spatiale (IAS), Universit Paris-Sud, F-91405 Orsay, France

⁶Instituto de Astrofísica de Canarias, C/ Via Láctea, E-38205 La Laguna, Tenerife, Spain

⁷Departamento de Astrofísica, Universidad de La Laguna, E-38205 La Laguna, Tenerife, Spain

Accepted 2014 August 15. Received 2014 August 15; in original form 2014 May 9

ABSTRACT

We use deep *Herschel Space Observatory* observations of a 90 per cent complete sample of 32 intermediate-redshift 2Jy radio galaxies ($0.05 < z < 0.7$) with strong emission lines to estimate the dust masses of their host galaxies and thereby investigate the triggering mechanisms for their quasar-like AGN. The dust masses derived for the radio galaxies ($7.2 \times 10^5 < M_d < 2.6 \times 10^8 M_\odot$) are intermediate between those of quiescent elliptical galaxies on the one hand, and ultraluminous infrared galaxies (ULIRGs) on the other. Consistent with simple models for the co-evolution of supermassive black holes and their host galaxies, these results suggest that most radio galaxies represent the late time re-triggering of AGN activity via mergers between the host giant elliptical galaxies and companion galaxies with relatively low gas masses. However, a minority of the radio galaxies in our sample (~ 20 per cent) have high, ULIRG-like dust masses, along with evidence for prodigious star formation activity. The latter objects are more likely to have been triggered in major, gas-rich mergers that represent a rapid growth phase for both their host galaxies and their supermassive black holes.

Key words: galaxies:active – galaxies: interactions – galaxies: ISM – galaxies:jets – quasars: general.

1 INTRODUCTION

Radio-loud AGN are particularly important for understanding the link(s) between galaxy evolution and nuclear activity, because, in contrast to samples of radio-quiet AGN in the local Universe (e.g. Seyfert galaxies), they are invariably associated with early-type host galaxies, allowing relatively ‘clean’ searches to be made for the signs of the triggering events. They also drive powerful relativistic jets that are capable of projecting the power of the AGN into the gaseous haloes of the host galaxies and clusters of galaxies, directly affecting the cooling of the hot IGM/ICM gas – one of the most important forms of AGN-induced feedback (Best et al. 2006; McNamara & Nulsen 2007).

Recently, we have undertaken a major programme of optical (Ramos Almeida et al. 2011, 2012) and mid- to far-IR (Dicken et al. 2008, 2009, 2012) observations of the 2Jy sample of south-

ern radio galaxies ($0.05 < z < 0.7$; Tadhunter et al. 1998), aimed at directly investigating the triggering mechanisms for radio-loud AGN. Our deep Gemini optical imaging observations show tidal features (tails, shells, fans, bridges) at relatively high surface brightness levels in a majority of objects (> 85 per cent), consistent with the idea that radio-loud AGN represent a fleeting active phase in the evolution of the subset of giant elliptical galaxies that have recently undergone galaxy interactions and mergers (Ramos Almeida et al. 2012). However, the radio-loud AGN phase does not correspond to a single stage of the merger sequence, nor to a single type of galaxy merger or interaction (Ramos Almeida et al. 2011, 2012). Such apparent diversity is also consistent with the fact that only a minority (≤ 35 per cent) of our sample show evidence for recent star formation activity, based on the direct detection of young stellar populations at optical wavelengths (Tadhunter et al. 2002, 2011), far-IR excesses from *Spitzer* photometry (Dicken et al. 2009), and PAH features in mid-IR *Spitzer* spectroscopy (Dicken et al. 2012).

Our previous work suggests that only a small fraction (≤ 20 per cent) of morphologically disturbed elliptical galaxies are

★E-mail: c.tadhunter@sheffield.ac.uk

capable of hosting a powerful radio-loud AGN (Ramos Almeida et al. 2012). Therefore, simply undergoing a galaxy interaction of *any type* is not by itself sufficient to trigger a powerful radio-loud AGN in a giant elliptical galaxy. One of the most important additional factors is likely to be gas content: it is probable that powerful, FR II-like radio activity is triggered by accretion of the cool/warm phases of the ISM, rather than the hot phase (e.g. Hardcastle, Evans & Croston 2007). Therefore, we expect the ability to host a powerful radio-loud AGN to be related at some level to the total reservoir of cool ISM which, in the case of the elliptical galaxy hosts of radio-loud AGN, is likely to have had an external origin.

Simple arguments allow us to predict the total gas reservoir required to trigger quasar-like activity and sustain it over the quasar lifetime (t_{qso} yr). If we define a quasar to have a bolometric luminosity of $L_{\text{BOL}} > 10^{38}$ W, a mass inflow rate of $\dot{M} > 0.2 M_{\odot} \text{ yr}^{-1}$ on to the central supermassive black hole is required to produce the quasar activity, assuming an efficiency of 10 per cent. Therefore, over the lifetime of the quasar the supermassive black hole will accrete a mass $M_{\text{ac}} > 0.2 t_{\text{qso}} M_{\odot}$, amounting to $M_{\text{ac}} > 2 \times 10^6 M_{\odot}$ for a typical quasar lifetime of $t_{\text{qso}} \sim 10^7$ yr (Martini 2004). However, on the basis of the black hole mass versus host bulge mass correlations, we know that only a small fraction of the gas accreted by the bulge reaches the supermassive black hole: for every solar mass accreted by a supermassive black hole, ~ 500 solar masses of stars must be formed in the bulge of the host galaxy (Marconi & Hunt 2003). In this case, the *total* gas reservoir that is expected to be associated with a particular quasar triggering event (M_{tot}) is much larger: $M_{\text{tot}} \sim 100 t_{\text{qso}} M_{\odot}$ or $M_{\text{tot}} \sim 10^9 M_{\odot}$ for $t_{\text{qso}} \sim 10^7$ yr. Although this argument is oversimplified in the sense that some of gas may already have been used up in star formation or ejected by feedback processes by the time the quasar is observed, it provides an order-of-magnitude estimate of the total gas reservoir required.

In this Letter, we use deep *Herschel* observations of a sample of nearby powerful radio galaxies with quasar-like nuclei to address the issue of whether the total cool ISM reservoirs are as massive as expected based on this simple model for the co-evolution of galaxy bulges and their central supermassive black holes. We also use the results to investigate the nature of the triggering merger events. A cosmology with $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_{\text{m}} = 0.27$, and $\Omega_{\lambda} = 0.73$ is assumed throughout this Letter.

2 SAMPLE, OBSERVATIONS, AND DATA REDUCTION

The full sample for this project, as described in Dicken et al. (2009, 2012), comprises all 46 radio sources with redshifts $0.05 < z < 0.7$, radio fluxes $S_{2.7 \text{ GHz}} > 2.0 \text{ Jy}$, declinations $\delta < +10$ deg, and radio spectral indices $\alpha < -0.7$ (assuming $F_{\nu} \propto \nu^{+\alpha}$). This southern 2Jy sample is now the best observed of all samples of powerful radio galaxies in terms of the depth and completeness of its multiwavelength data.¹

All 46 objects in the southern 2Jy sample were observed at 100 and 160 μm with the *Herschel* PACS instrument as part of a programme of ‘Must Do’ observations in the last days of *Herschel*; and the 16 objects with the brightest *Spitzer* 70 μm fluxes were also observed with the *Herschel* SPIRE instrument at 250, 350, and 500 μm in order to characterize the shapes of their longer wavelength SEDs. The depths of the PACS observations allowed us to




Figure 1. Example radio to infrared SEDs for PKS0806-10 (top) and PKS0349-27 (bottom). The symbols are as follows. Open squares: total radio fluxes; open circles: core radio fluxes; filled triangles: *Herschel*/SPIRE fluxes; filled circles: *Herschel*/PACS fluxes; filled squares: *Spitzer*/MIPS fluxes. The solid lines represent the *Spitzer*/IRS spectra for the sources.

detect 100 and 87 per cent of the sample at 100 and 160 μm , respectively. In addition, for the purpose of quantifying the degree of any non-thermal contamination of the *Herschel* photometry, we made sub-mm 870 μm observations with LABOCA/APEX of the 20 objects previously identified with *Spitzer* as having the possibility of non-thermal contamination (Dicken et al. 2008).

Although our full sample comprises a mixture of weak-line radio galaxies and strong-line radio galaxies (SLRG: $\text{EW}_{[\text{O III}]}) > 10 \text{ \AA}$), in this Letter we concentrate on the 35 objects in our sample with strong emission lines, indicating the presence of quasar-like AGN in addition to the powerful radio jets and lobes. Discussion of the full sample, including details of the observations, data reduction, far-IR photometry and SEDs, will be presented in Dicken et al. (in preparation), while basic information about the SLRG sample used in this Letter is presented separately in the appendix in Table A1 (available as Supporting Information).

In Fig. 1 (top), we show an example SED of one of the brighter sources in our sample for which we have full flux information across the *Herschel* PACS and SPIRE bands, while in Fig. 1 (bottom) we show the SED of a more typical, fainter object that was only detected with PACS.

Given that all the objects in our sample are radio-loud AGN, it is important to be cautious about the potential contamination of the far-IR continuum by non-thermal emission from the synchrotron-emitting radio jets, lobes, and cores. As discussed in Dicken et al. (2008), such contamination is most likely to be important in objects where the flat-spectrum cores and inner jets are unusually strong, or the steep-spectrum radio lobes are contained within the *Herschel* beam for the PACS and SPIRE observations. By using simple

¹ See http://2jy.extragalactic.info/The_2Jy_Sample.html

power-law extrapolations of our high-frequency (20 GHz) radio and LABOCA 870 μm measurements, we find that non-thermal contamination is significant at far-IR wavelengths in 6/35 (17 per cent) of the objects in our SLRG sample (see Dicken et al., in preparation, for details). For three of these objects – all compact steep spectrum sources – it has proved possible to accurately subtract the non-thermal power law, and thereby derive fluxes and luminosities for the thermal dust emission. However, for the remaining three sources (PKS0035-02, PKS1136-13, and PKS1306-09) the non-thermal contamination makes a larger contribution at far-IR wavelengths, and we cannot confidently subtract this component because of uncertainties about the extrapolation of the SEDs. Therefore, we exclude the latter three objects from our analysis,² making our sample of 2Jy SLRG 90 per cent complete overall.

3 DETERMINING THE COOL DUST MASSES

The determination of dust masses from the far-IR data is highly sensitive to the assumptions made in the analysis, for example the form of the long-wavelength dust emissivity law, as characterized by the β index for a modified blackbody fit ($F_\nu \propto \nu^\beta B(\nu, T)$), and the number of cool dust components of different temperature assumed in the model (e.g. Clements, Dunne & Eales 2010).

In order to simplify our analysis, and allow the direct comparisons with the dust masses of ULIRGs and normal non-active elliptical galaxies presented in Section 4 and Fig. 2, we assumed a single temperature (T_d) for the cool dust in each object. Furthermore, we assumed that all objects in our sample have the same β index. In this case, the dust mass is given by

$$M_d = \frac{S_{160}^K D^2}{\kappa_{160}^m B(160, T_d)(1+z)}, \quad (1)$$

where S_{160}^K is the K -corrected flux density at 160 μm , D is the luminosity distance, κ_{160}^m is the dust opacity per unit mass at a wavelength of 160 μm , and $B_\nu(160, T_d)$ is the flux density of a blackbody of temperature T_d at 160 μm . For the purpose of this analysis, we assume $\kappa_{160}^m = 1.06 \text{ m}^2 \text{ kg}^{-1}$ (Draine 2003).

We determined a typical β index for the objects in our sample by comparing the 100/160 and 160/250 colours measured for the brighter objects observed with SPIRE, with the predicted colours for modified black bodies of different temperature and beta index (see Dicken et al., in preparation). From this comparison, we find that $\beta = 1.2$ is most consistent with the observations of the far-IR bright 2Jy objects in our sample.

We then determined the temperatures of the cool dust components for all the objects in our sample with both 100 and 160 μm fluxes by comparing the 100/160 flux ratios with those predicted by modified blackbody curves of different T_d but the same β index ($\beta = 1.2$), with uncertainties in the temperatures estimated from the uncertainties in the 100/160 flux ratios. The cool dust temperatures fall in the range $30 < T_d < 70 \text{ K}$, with a median dust temperature $T_d = 41 \text{ K}$. Finally, the dust masses were calculated from the K -corrected 160 μm fluxes and estimated temperatures using equation (1) above. Note that the uncertainties in the dust masses take into

²Note that the upper limiting dust masses, derived for these objects by assuming that all their 160 μm radiation is dust emission, fall well within the range measured for the other objects in our sample; excluding these objects does not affect any of the comparisons with the other samples in Section 4.




Figure 2. Dust mass distributions for 2Jy radio galaxies (top), ULIRGs (middle), and elliptical galaxies (bottom); see text for details of the sample selection. The vertical dashed and dotted lines show the equivalent dust masses of the Milky Way and LMC, respectively (both calculated from total hydrogen gas masses assuming $M_g/M_d = 140$).

account the uncertainties in both the far-IR fluxes and the dust temperatures, but are inevitably dominated by the uncertainties in the temperatures.

Six of the SLRG in our sample were not detected at 160 μm . In such cases, we derived the dust masses using the K -corrected 100 μm fluxes, assuming the median dust temperature derived from the other 26 objects in the sample, and $\kappa_{100}^m = 2.71 \text{ m}^2 \text{ kg}^{-1}$ (Draine 2003) in equation (1). The distribution dust masses for these six objects is similar to that of the sample as a whole. Therefore, the inclusion/exclusion of these objects does not alter our conclusions.

Clearly, our dust mass results are sensitive to the assumed value of the β index, which affects both the temperatures derived from the 100/160 flux ratios and the K -correction of the far-IR fluxes. However, we find that changing the β index from 1.2 to 2.0 – the value assumed in some studies of dust masses in elliptical galaxies,

for example – only increases the median dust mass by 80 per cent. Moreover, comparisons between the dust masses derived for various overlapping samples of low-redshift ULIRGs (Klaas et al. 2001; Farrah et al. 2003; Clements, Dunne & Eales 2010) and elliptical galaxies (Smith et al. 2012; di Serego Alighieri et al. 2013; Martini, Dicken & Storchi-Bergmann 2013) suggest that the effect of changing the β index, number of dust components, and general modelling approach is at most a factor of ~ 2 – 3 in the dust mass. This will not significantly affect the main results of the comparisons with the samples of the quiescent elliptical galaxies and ULIRGs presented in the next section.

4 RESULTS

The main result of this study is that the SLRG in our complete sample are moderately dust-rich. Fig. 2 shows a histogram of the dust masses derived from our *Herschel* data. The dust masses cover the range $7.2 \times 10^5 < M_d < 2.6 \times 10^8 M_\odot$, with a median dust mass of $8.5 \times 10^6 M_\odot$. Assuming a typical gas-to-dust ratio of 140 (Draine 2007; Parkin et al. 2012), these translate into gas masses in the range $1.0 \times 10^8 < M_g < 3.7 \times 10^{10} M_\odot$, with a median gas mass of $1.2 \times 10^9 M_\odot$. We note that this median gas mass is remarkably close to that predicted on the basis of the simple argument presented in the Introduction. Therefore, it appears that there is indeed sufficient cool ISM in these radio galaxies to sustain the quasar-like activity for the requisite time-scales, and simultaneously grow the stellar mass of the host galaxies, maintaining the black hole/host scaling relationships.

To put these results into context, Fig. 2 also shows the distributions of dust masses derived for nearby ULIRGs and elliptical galaxies. The ULIRG dust masses were taken from the results of the variable β index modified blackbody fits to the far-IR/sub-mm SEDs of the sample of 23 nearby ($z < 0.17$) ULIRGs presented by Clements et al. (2010), while the elliptical galaxy dust masses were taken from the modified blackbody $\beta = 2$ results for the non-active ellipticals in Smith et al. (2012) and di Serego Alighieri et al. (2013), as well as from the template fitting results of Martini et al. (2013); the final sample of elliptical galaxies comprises 40 objects with distances in the range $15 < D < 66$ Mpc.

The median dust masses of the ULIRG and elliptical galaxy samples plotted in Fig. 2 are 8×10^7 and $2 \times 10^5 M_\odot$, respectively. At the high dust mass end of the distributions there is clearly an overlap between the dust masses of the radio galaxies and those of the ULIRGs; the lower end of the radio galaxy dust mass distribution also overlaps with the higher end of the elliptical dust mass distribution. However, on average, the radio galaxies have dust masses that are more than an order of magnitude *higher* than those of typical elliptical galaxies, but an order of magnitude *lower* than those of typical ULIRGs.

We emphasize that our comparison between the 2Jy radio galaxies and the elliptical galaxies is conservative, because (a) we have included upper limiting dust masses for the elliptical galaxies (~ 53 per cent of cases) as if they were detections; (b) for objects in common between the samples we have favoured the Smith et al. (2012) single modified blackbody fits – which give generally larger dust masses – over the Martini et al. (2013) template fitting results; and (c) the elliptical galaxy dust mass estimates are dominated by the results of the Smith et al. (2012) and di Serego Alighieri et al. (2013) $\beta = 2.0$ single modified blackbody fits, which tend to give higher dust masses than the $\beta = 1.2$ fits used for the radio galaxies.

In terms of the comparison with the ULIRGs, the single modified blackbody fits of Clements et al. (2010) with variable beta index (as

shown in Fig. 2) are the most comparable with the $\beta = 1.2$ results presented for the radio galaxies. However, even if we compare the $\beta = 2.0$ results for the 2Jy radio galaxies with the $\beta = 2.0$ results of Klaas et al. (2001) – which give the lowest dust masses of all the low- z ULIRG studies – the radio galaxy dust masses remain a factor ~ 4 lower on average.

We further note that the total cool ISM mass implied by the median radio galaxy dust mass ($M_g = 1.2 \times 10^9 M_\odot$) is significantly lower than that of the Milky Way ($4.8 \times 10^9 M_\odot$; Draine 2011), and about a factor of 2 higher than that of the Large Magellanic Cloud (LMC, $5 \times 10^8 M_\odot$; Westerlund 1990).

The other interesting feature of the radio galaxy dust mass distribution is that a large proportion (78 per cent) of the nine radio galaxies with the highest dust masses ($M_d > 2 \times 10^7 M_\odot$) – comparable with those of ULIRGs – belong to the rare class of radio galaxies that show evidence for prodigious star formation activity (Tadhunter et al. 2011; Dicken et al. 2012). Indeed, the two highest dust mass radio galaxies – PKS2314+05 and PKS2135-20 – are classified as ULIRGs based on their far-IR luminosities.

5 DISCUSSION AND CONCLUSIONS

The dust masses of the radio source host galaxies, and the total cool ISM masses that they imply, provide important clues to the nature of the events that triggered the powerful AGN and jet activity in these systems, complementing existing studies of the 2Jy sample host galaxy morphologies (Ramos Almeida et al. 2011, 2012) and levels of star formation activity (Dicken et al. 2012).

The radio galaxies with the highest dust masses have cool ISM contents, morphologies, and star formation properties that are similar to those of ULIRGs in the local Universe. In such cases, it is likely that the AGN have been triggered in major gas-rich mergers in which a substantial fraction of the stellar mass is being built up via merger-induced star formation and in which the black holes are undergoing a major growth phase. However, such starburst radio galaxies are rare in the local radio galaxy population (Tadhunter et al. 2011; Dicken et al. 2012). More typical radio galaxies have much lower cool ISM masses, and most of them lack evidence for recent star formation activity (Dicken et al. 2012), even if they do show morphological signs of recent galaxy interactions (Ramos Almeida et al. 2011, 2012). In these more typical cases it is likely that we are witnessing the late-time re-triggering of quasar activity via mergers between mature giant elliptical galaxies and companion galaxies that have gas masses between those of the LMC and M33 in the Local Group. As we showed in the Introduction, such mergers are capable of delivering sufficiently massive reservoirs of cool gas to the radio source hosts to fuel quasar-like activity on the requisite time-scales.

Finally, we note that, while most of the nearby elliptical galaxies that lack powerful AGN activity have low dust masses (see Fig. 2), a small but significant fraction of such objects have much larger dust masses. An interesting example in the local Universe is Centaurus A, which is associated with a relatively low-power FR I radio source and currently lacks a powerful quasar AGN, but none the less contains a substantial reservoir of cool ISM in its well-known dust lane ($M_d = 1.6 \times 10^7 M_\odot$, $M_g = 2.7 \times 10^9 M_\odot$; Parkin et al. 2012). This raises the question of why Centaurus A, and gas rich but quiescent elliptical galaxies in general, do not always appear as quasars. Most likely, the presence of a substantial gas reservoir is a necessary but not sufficient condition for triggering luminous quasar activity; other factors, such as the detailed distribution and kinematics of the cool ISM are also likely to be important.

In the particular case of Centaurus A, it is plausible that we are observing the system in a late post-merger phase in which the ISM in the dust lane has settled into a dynamically stable configuration (see also Tadhunter et al. 2011). In this case, the radial gas flows into the nuclear regions are insufficient to sustain luminous quasar activity. In contrast, objects with quasar nuclei may represent an earlier phase in the merger when the gas is still settling into a stable configuration, and the rate of radial flow of the gas into the nucleus is much higher. Clearly, high-resolution CO observations that measure both the distribution and kinematics of the cool ISM in nearby radio galaxies with quasar-like nuclei will be crucial for making further progress in this field. Such observations will also provide an important check on the cool ISM masses that we have estimated from the thermal dust emission.

ACKNOWLEDGEMENTS

We thank everyone involved with the *Herschel* Observatory and the PACS and SPIRE instruments. CT acknowledges support from STFC consolidated grant ST/J001589/1. CRA is supported by a Marie Curie Intra European Fellowship within the 7th European Community Framework Programme (PIEF-GA-2012-327934). RM gratefully acknowledges support from the ERC under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC Advanced Grant RADIOLIFE-320745.

REFERENCES

- Best P., Kaiser C. R., Heckman T. M., Kauffmann G., 2006, MNRAS, 368, L67
- Clements D. L., Dunne L., Eales S., 2010, MNRAS, 403, 274
- di Serego Alighieri S. et al., 2013, A&A, 552, A8
- Dicken D., Tadhunter C., Morganti R., Buchanan C., Oosterloo T., Axon D., 2008, ApJ, 678, 712
- Dicken D. et al., 2009, ApJ, 694, 268
- Dicken D. et al., 2012, ApJ, 745, 172
- Draine B., 2003, ARA&A, 41, 241
- Draine B. T., 2011, Physics of the Interstellar and Intergalactic Medium. Princeton Univ. Press, Princeton, NJ, p. 5
- Draine B. T. et al., 2007, ApJ, 663, 866
- Farrah D., Afonso J., Efstathiou A., Rowan-Robinson M., Fox M., Clements D., 2003, MNRAS, 343, 585
- Hardcastle M., Evans D. A., Croston J. H., 2007, MNRAS, 376, 1849
- Klaas U. et al., 2001, A&A, 379, 823
- McNamara B., Nulsen P., 2007, ARA&A, 45, 117
- Marconi A., Hunt L., 2003, ApJ, 589, L21
- Martini P., 2004, in Ho L. C., ed., Carnegie Observatories Astrophysics Series, Vol. 1: Coevolution of Black Holes and Galaxies. Cambridge Univ. Press, Cambridge, p. 169
- Martini P., Dicken D., Storchi-Bergmann T., 2013, ApJ, 766, 121
- Parkin T. J. et al., 2012, MNRAS, 422, 2291
- Ramos Almeida C., Ramos Almeida C., Tadhunter C. N., Inskip K. J., Morganti R., Holt J., Dicken D., 2011, MNRAS, 410, 1550
- Ramos Almeida C. et al., 2012, MNRAS, 419, 687
- Smith M. W. L. et al., 2012, ApJ, 748, 123
- Tadhunter C., Morganti R., Robinson A., Dickson R., Villar-Martin M., Fosbury R. A. E., 1998, MNRAS, 298, 1035
- Tadhunter C., Dickson R., Morganti R., Robinson T. G., Wills K., Villar-Martin M., Hughes M., 2002, MNRAS, 330, 997
- Tadhunter C., González Delgado R., Rodríguez Zaurín J., Villar-Martín M., Morganti R., Emonts B., Ramos Almeida C., Inskip K., 2011, MNRAS, 412, 960
- Westerlund B. E., 1990, A&AR, 2, 29

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Table A1. General properties of the sample of 2Jy strong line radio galaxies (SLRG) included in the paper. The third and fourth columns give the optical (BLRG: broad-line radio galaxy; NLRG: narrow-line radio galaxy; Q: quasar) and radio (FR I: Fanaroff–Riley type I source; FR II: Fanaroff–Riley type II source; CSS: compact steep spectrum source; GPS: gigahertz-peaked spectrum source) classifications, with any uncertain classifications in brackets. Columns five and six indicate whether the source was detected by *Herschel* PACS at 100 and 160 mm, respectively, while the final column indicates whether the source was observed using *Herschel*/SPIRE. An asterisk in column 1 indicates that the source was excluded from the final sample with dust mass estimates because of non-thermal contamination that could not be corrected. Details of the full 2Jy sample are given in Dicken et al. (2008, 2009, 2012) (<http://mnrasl.oxfordjournals.org/lookup/suppl/doi:10.1093/mnrasl/slu135/-/DC1>).

Please note: Oxford University Press are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.