

University of Groningen

N-Hydroxyimide Ugi Reaction toward α -Hydrazino Amides

Chandgude, Ajay L; Dömling, Alexander

Published in:
Organic letters

DOI:
[10.1021/acs.orglett.7b00205](https://doi.org/10.1021/acs.orglett.7b00205)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Chandgude, A. L., & Dömling, A. (2017). N-Hydroxyimide Ugi Reaction toward α -Hydrazino Amides. *Organic letters*, 19(5), 1228-1231. DOI: 10.1021/acs.orglett.7b00205

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

N-Hydroxyimide Ugi Reaction toward α -Hydrazino AmidesAjay L. Chandgude^{1b} and Alexander Dömling*^{1b}

Department of Drug Design, University of Groningen, Antonius Deusinglaan 1, 9713 AV Groningen, The Netherlands

Supporting Information

ABSTRACT: The Ugi four-component reaction (U-4CR) with *N*-hydroxyimides as a novel carboxylic acid isostere has been reported. This reaction provides straightforward access to α -hydrazino amides. A broad range of aldehydes, amines, isocyanides and *N*-hydroxyimides were employed to give products in moderate to high yields. This reaction displays N–N bond formation by cyclic imide migration in the Ugi reaction. Thus, *N*-hydroxyimide is added as a new acid component in the Ugi reaction and broadens the scaffold diversity.



The Ugi reaction (U-4CR) is a widely used multi-component reaction (MCR) for the synthesis of bis-amides and peptidomimetics.¹ This reaction has emerged as a powerful synthetic method for the organic, pharmaceutical, and polymer industries.² However, it cannot meet the ever-increasing need for molecular complexity and diversity in organic and medicinal chemistry. An increasing demand for novel scaffolds has led to the interest in U-4CR post-modifications and single-reactant replacement (SRR) by isosteres.³ U-4CR postmodifications are useful for the synthesis of various heterocycles and peptidic scaffolds.⁴ Nevertheless, isostere use in the Ugi reaction is rather limited.⁵ An amine component could be replaced by secondary amines, hydroxylamines, and hydrazines. As with amines, use of acid isosteres in the Ugi reaction is also limited.

In the Ugi reaction, the carboxylic acid plays several prominent structural roles, including activation of the intermediate imine, the reversible addition to the nitrilium ion, and participation in the irreversible Mumm rearrangement to form the final bis-amide product. Because of carboxylic acid's substantial role in the reaction, isosteric replacement by other agents is difficult to accomplish. In 1962, Ugi reported the first acid isosteric replacements by inorganic acids, such as hydrazoic acids, cyanates, thiocyanates, etc. (Figure 1).⁶ To date, only a few acid isosteres have been reported. For example, our group reported the thioacetic acid as an isostere.⁷ El Kaïm and co-workers reported the phenol as an acid isostere in the Ugi reaction involving Smiles rearrangement to form an *N*-arylamine.⁸ Other Ugi–Smiles and similar strategies have been described by El Kaïm (thiophenol),⁹ Charton (squaric acid),¹⁰ and Neo (hydroxycoumarins).¹¹ Further, Lewis acids and CO₂ were used as acid isosteres in the U-4CR.⁵

As for the related Passerini reaction,¹² organic acid isostere replacement has remained largely unexplored for the Ugi-4CR as there are only a few examples of isostere use in the Ugi-4CR for the synthesis of peptidomimetics.⁵ Therefore, finding new isosteres in U-4CR for the synthesis of diverse and complex peptidomimetic derivatives is of high interest.

We hypothesized that *N*-hydroxyimides could be used as a novel acid isostere in the U-4CR reaction, which can directly

Previous acid components in U-4CR



Introduction of *N*-hydroxamic acid in U-4CR

Figure 1. Previously reported and new acid isosteres in U-4CR.

provide the α -hydrazino amides as Ugi reaction products. α -Hydrazino amides are aza analogues of β -peptides and are of interest for several reasons.¹³ These foldamers exhibit the special hydrazino turn, and the hydrazidic bond is very resistant to protease.¹⁴ Hydrazino amides are found in many natural products such as linatine, a vitamin B6 antagonist;^{15a} negamycine, an antibiotic;^{15b} and matlystatins, antimicrobial compounds.^{15c} They also have broad applications in medicinal chemistry including use as proteasome inhibitors,^{16a} antimicrobials,^{16b} DNA and RNA interactors,^{16c} (S)-(–)-carbidopa

Received: January 30, 2017

Published: February 21, 2017


for the treatment of Parkinson's disease,¹⁷ or as human leukocyte elastase (HLE) inhibitors.¹⁸

Hydrazino peptides are mainly synthesized by two methods: first, by using hydrazine derivatives¹⁹ and, second, by coupling of an amino group with another amine typically employing oxaziridines.¹⁴ The general use of hydrazine and oxaziridine is limited by unavailability of diverse derivatives and their highly toxic and unstable nature. Another drawback is that their synthesis is laborious. Therefore, the development of new methods for the synthesis of this important foldamer is highly desirable.

Herein, we report the successful use of the *N*-hydroxyimides as an acid isostere in the U-4CR for a direct route to the synthesis of α -hydrazino amides. This is the second example of cyclic imide migration to nitrogen (O \rightarrow N imide transfer) in the Mumm rearrangement to form an N–N bond. This type of N–O bond breaking and N–N bond formation in a Mumm-type rearrangement has been recently reported.²⁰ This reaction illustrates the use of *N*-hydroxamic acid for N–N bond formation without phthalimidation.

We started our optimization by using propionaldehyde, benzylamine, cyclohexyl isocyanide, and *N*-hydroxyphthalimides (NHPI) as model reactants. Reaction in methanol did not form any desired product (Table 1, entry 1). In polar aprotic solvents such as THF and CH₃CN, only traces of product were formed (see the Supporting Information for details of optimization conditions). In the polar protic solvent

Table 1. Optimization Conditions^a



entry	solvent	temp (°C)	catalyst	time (h)	% yield ^b
1	MeOH	rt		12	
2	THF	rt		12	trace
3	CH ₃ CN	rt		12	trace
4	DCE	rt		12	25
5	toluene	rt		12	20
6 ^c	DCE	rt	ZnCl ₂		22
7 ^c	DCE	rt	Sc(OTf) ₃	12	15
8 ^c	DCE	rt	I ₂	12	22
9 ^c	DCE	rt	TMSCl	12	19
10 ^c	DCE	rt	InCl ₃	12	10
11 ^c	THF	50	I ₂	12	trace
12 ^c	DCE	50	ZnCl ₂	12	nd
13 ^c	toluene	rt	ZnCl ₂	12	51
14 ^c	xylene	rt	ZnCl ₂	12	49
15 ^c	chlorobenzene	rt	ZnCl ₂	12	31
16 ^c	TFE	rt	ZnCl ₂	12	38
17 ^d	DCE	rt	ZnCl ₂	12	40
18 ^d	toluene	rt	ZnCl ₂	12	66
19 ^e	toluene	rt	ZnCl ₂	12	47
20 ^f	toluene	rt	ZnCl ₂	2	43
21 ^g	toluene	rt	ZnCl ₂	2	50

^aThe reaction was carried out with propionaldehyde (1.0 mmol), benzylamine (1.0 mmol), cyclohexyl isocyanide (1.0 mmol), and *N*-hydroxyphthalimide (1.5 mmol) in 2 mL of solvent. ^bYield of isolated product **5a**. ^c10 mol % of catalyst used. ^d30 mol % of ZnCl₂ used. ^e50 mol % of ZnCl₂ used. ^fReaction performed in sonication with 10 mol % of ZnCl₂. ^gReaction performed in sonication with 30 mol % of ZnCl₂. nd = not determined.

MeOH, the U-3CR product, α -amino amide, was formed as a major product; in contrast, it formed only in trace amounts in solvents such as THF and toluene. This U-3CR product formation might be due to the low acidity of *N*-hydroxyimide (pK_a \sim 7.5), which functioned only as a catalyst. This observation led us to try a nonpolar solvent and Lewis acid to activate *N*-hydroxyimide for the further optimization. Indeed, nonpolar solvents such as DCE and toluene allowed moderate product formation of 25% and 20%, respectively, at room temperature.

Next, we screened various Lewis acids, (Table S2) such as InCl₃, I₂, Sc(OTf)₃, etc. (10 mol %) in DCE as a solvent. We found that ZnCl₂ was the best of the screened Lewis acids (Table 1, entries 7–11). An increase in the temperature failed to improve the product yield (Table 1, entries 11 and 12). In further solvent screening with ZnCl₂, (Table S3), we observed that toluene and xylene gave similar yields, 51% and 49%, respectively (Table 1, entries 13 and 14). The nature of the solvent played a critical role in the success of the reaction. Next, we performed a catalyst equivalence study in toluene as solvent. An increase in the catalyst quantity to 30 mol % gave the best yield of 66% (Table 1, entry 18). However, a further increase in the quantity of ZnCl₂ to 50 mol % gave a lower yield, 47% (Table 1, entry 19). The use of sonication in this reaction did not have any effect on product yield (Table 1, entries 20 and 21).^{12a}

With these optimized conditions in hand, next we examined the generality of this U-4CR by using various aldehydes, amines, isocyanides, and *N*-hydroxyimides (Table 2). Aliphatic aldehydes offered good yields, up to 78% (Table 2, entries 1–3). Aromatic aldehydes are also useful substrates in this reaction (Table 2, entries 6–9).

Electron-withdrawing and -donating groups in aromatic aldehydes at different positions such as ortho and para provided moderate to good yields. Amines with protected functional groups like acetal and halogens were well-tolerated in this reaction, affording moderate to good yields of the products (Table 2, entries 3, 4, and 6). The acid-protected amino acid β -alanine ester gave only 18% yield (Table 2, entry 5). Various aliphatic and aromatic isocyanides like cyclohexyl, phenylethyl, 2-nitrobenzyl, benzyl, 4-methoxyphenyl, and β -cyanoethyl were well-suited within the developed methodology. Among *N*-hydroxyimides, *N*-hydroxysuccinimides (NHS) also proceeded smoothly similarly to NHPI and gave 48–59% yield (Scheme 1, **5n–p**). However, the reaction with hydroxybenzotriazole (HOBt) resulted in only trace product formation (Scheme 1, **5q**).

A broad functional group tolerance in this reaction could be of interest for the postmodification condensations. Thus, among the vast number of possible postmodification reactions with this modified U-4CR, we attempted several. Hydrazines are important intermediates for the synthesis of many heterocycles and scaffolds.¹⁹ The U-4CR product (**5d**) treatment with hydrazine hydrate deprotects the NHPI and forms the free hydrazine derivatives **6**.^{12a} We obtained free hydrazine in a good yield of 64% after overnight reaction (Scheme 2).


Next, we turned our attention to creating access for pharmaceutically important α -amino amide molecules. We converted the U-4CR product (**5p**) to U-3CR product α -amino amide **7** in good yield (73%). This AlCl₃-catalyzed reaction cleaved the N–N bond (Scheme 3) to form the final product.²¹

Table 2. Substrate Scope^a




entry	1	2	3	4	% yield ^b
1				NHPI	66 (5a)
2				NHPI	74 (5b)
3				NHPI	78 (5c)
4				NHPI	58 (5d)
5 ^c				NHPI	18 (5e)
6				NHPI	57 (5f)
7				NHPI	51 (5g)
8				NHPI	46 (5h)
9				NHPI	21 (5i)
10				NHPI	91 (5j)
11				NHPI	38 (5k)
12				NHPI	71 (5l)
13				NHPI	65 (5m)

^aReaction conditions: **1** (1.0 mmol), **2** (1.0 mmol), **3** (1.0 mmol), and **4** (1.5 mmol), ZnCl₂ (30 mol %) in toluene (2 mL) at rt for overnight.
^bIsolated yield. ^c1.5 equiv of triethylamine used.

We did not carry out detailed mechanistic studies but envision the following mechanism (Scheme 4). ZnCl₂ activates an imine **A** to allow the nucleophilic addition of isocyanide **3** to form the nitrilium intermediate **C**. The hydroxamate

Scheme 1. *N*-Hydroxyimide Scope

Scheme 2. Deprotection toward Hydrazine Formation

Scheme 3. Deprotection toward α -Amino AmideScheme 4. Anticipated Mechanism for the Ugi–*N*-Hydroxyimide Reaction

nucleophilically traps the nitrilium intermediate **C**. Finally, this intermediate **D** undergoes an irreversible Mumm-like $N \rightarrow N$ migration to form the α -hydroxyimide **5**.

In conclusion, we have reported *N*-hydroxyimides as novel acid isosteres in the U-4CR toward the one-step synthesis of α -hydroxyimides via N – N bond formation. This mild and general reaction requires catalytic amounts of ZnCl₂. This protocol uses readily available *N*-hydroxyimides, which replace

the toxic and unstable hydrazines/oxaziridine use for the synthesis of α -hydrazino amides. The method is applicable for a wide range of aldehydes and amines and has the potential for multiple postmodifications. Such scaffolds will be useful to fill the screening decks of the European Lead Factory (ELF).²² Moreover, as this reaction has significant potential in peptidomimetics synthesis, studies on postmodification reactions are now in progress.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.7b00205](https://doi.org/10.1021/acs.orglett.7b00205).

General experimental procedures; compound characterization data; ¹H and ¹³C spectra of all compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: a.s.s.domling@rug.nl

ORCID

Ajay L. Chandgude: [0000-0001-8236-9626](https://orcid.org/0000-0001-8236-9626)

Alexander Dömling: [0000-0002-9923-8873](https://orcid.org/0000-0002-9923-8873)

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the University of Groningen. The Erasmus Mundus Scholarship “Svaagata” is acknowledged for a fellowship to A.C. The work was financially supported by the NIH (2R01GM097082-05) and by the Innovative Medicines Initiative (Grant Agreement No. 115489). Funding has also been received from the European Union’s Horizon 2020 research and innovation programme under MSC ITN “Accelerated Early stage drug dIScovery” (AEGIS) (Grant Agreement No. 675555).

■ REFERENCES

- (1) (a) Ugi, I.; Werner, B.; Domling, A. *Molecules* **2003**, *8*, 53–66. (b) Domling, A.; Ugi, I. *Angew. Chem., Int. Ed.* **2000**, *39*, 3168–3210. (c) Ugi, I.; Domling, A.; Horl, W. *Endeavour* **1994**, *18*, 115–122. (d) Domling, A. *Curr. Opin. Chem. Biol.* **2002**, *6*, 306–313.
- (2) (a) Hulme, C.; Gore, V. *Curr. Med. Chem.* **2003**, *10*, 51–80. (b) Domling, A. *Chem. Rev.* **2006**, *106*, 17–89. (c) Domling, A.; Wang, W.; Wang, K. *Chem. Rev.* **2012**, *112*, 3083–3135. (d) Zarganes-Tzitzikas, T.; Chandgude, A. L.; Domling, A. *Chem. Rec.* **2015**, *15*, 981–996.
- (3) Koopmanschap, G.; Ruijter, E.; Orru, R. V. A. *Beilstein J. Org. Chem.* **2014**, *10*, 544–598.
- (4) Sunderhaus, J. D.; Martin, S. E. *Chem. - Eur. J.* **2009**, *15*, 1300–1308.
- (5) (a) Ruijter, E.; Scheffelaar, R.; Orru, R. V. A. *Angew. Chem., Int. Ed.* **2011**, *50*, 6234–6246. (b) El Kaim, L.; Grimaud, L. *Tetrahedron* **2009**, *65*, 2153–2171.
- (6) Ugi, I. *Angew. Chem., Int. Ed. Engl.* **1962**, *1*, 8–21.
- (7) Heck, S.; Domling, A. *Synlett* **2000**, 424–426.
- (8) El Kaim, L.; Grimaud, L.; Oble, J. *Angew. Chem., Int. Ed.* **2005**, *44*, 7961–7964.
- (9) Barthelon, A.; El Kaim, L.; Gizolme, M.; Grimaud, L. *Eur. J. Org. Chem.* **2008**, *2008*, 5974–5987.

(10) Akin, K.; Gauriot, M.; Totobenazara, J.; Deguine, N.; Deprez-Poulain, R.; Deprez, B.; Charton, J. *Tetrahedron Lett.* **2012**, *53*, 458–461.

(11) Neo, A. G.; Castellano, T. G.; Marcos, C. F. *Synthesis* **2015**, *47*, 2431–2438.

(12) (a) Chandgude, A. L.; Domling, A. *Org. Lett.* **2016**, *18*, 6396–6399. (b) Soeta, T.; Kojima, Y.; Ukaji, Y.; Inomata, K. *Org. Lett.* **2010**, *12*, 4341–4343. (c) Soeta, T.; Matsuzaki, S.; Ukaji, Y. *Chem. - Eur. J.* **2014**, *20*, 5007–5012. (d) Soeta, T.; Ukaji, Y. *Chem. Rec.* **2014**, *14*, 101–116.

(13) (a) Salaun, A.; Potel, M.; Roisnel, T.; Gall, P.; Le Grel, P. *J. Org. Chem.* **2005**, *70*, 6499–6502. (b) Gunther, R.; Hofmann, H. J. *J. Am. Chem. Soc.* **2001**, *123*, 247–255.

(14) Lelais, G.; Seebach, D. *Helv. Chim. Acta* **2003**, *86*, 4152–4168.

(15) (a) Klosterman, H. J.; Lamoureux, G. L.; Parsons, J. L. *Biochemistry* **1967**, *6*, 170–177. (b) McKinney, D. C.; Basarab, G. S.; Cocozaki, A. I.; Foulk, M. A.; Miller, M. D.; Ruvinsky, A. M.; Scott, C. W.; Thakur, K.; Zhao, L.; Buurman, E. T.; Narayan, S. *ACS Med. Chem. Lett.* **2015**, *6*, 930–935. (c) Tanzawa, K.; Ishii, M.; Ogita, T.; Shimada, K. *J. Antibiot.* **1992**, *45*, 1733–1737.

(16) (a) Bordessa, A.; Keita, M.; Marechal, X.; Formicola, L.; Lagarde, N.; Rodrigo, J.; Bernadat, G.; Bauvais, C.; Soulier, J. L.; Dufau, L.; Milcent, T.; Crousse, B.; Reboud-Ravaux, M.; Onger, S. *Eur. J. Med. Chem.* **2013**, *70*, 505–524. (b) Laurencin, M.; Amor, M.; Fleury, Y.; Baudy-Floc’h, M. *J. Med. Chem.* **2012**, *55*, 10885–10895. (c) Suc, J.; Tumir, L. M.; Glavas-Obrovac, L.; Jukic, M.; Piantanida, L.; Jeric, I. *Org. Biomol. Chem.* **2016**, *14*, 4865–4874.

(17) Vickers, S.; Stuart, E. K.; Hucker, H. B.; Vandenheuevel, W. J. A. *J. Med. Chem.* **1975**, *18*, 134–138.

(18) Guy, L.; Vidal, J.; Collet, A.; Amour, A.; Reboud-Ravaux, M. *J. Med. Chem.* **1998**, *41*, 4833–4843.

(19) (a) Suc, J.; Baric, D.; Jeric, I. *RSC Adv.* **2016**, *6*, 99664–99675. (b) Hashimoto, T.; Kimura, H.; Kawamata, Y.; Maruoka, K. *Angew. Chem., Int. Ed.* **2012**, *51*, 7279–7281. (c) Krasavin, M.; Bushkova, E.; Parchinsky, V.; Shumsky, A. *Synthesis* **2010**, *2010*, 933–942. (d) Busnel, O.; Bi, L. R.; Dali, H.; Cheguillaume, A.; Chevance, S.; Bondon, A.; Muller, S.; Baudy-Floc’h, M. *J. Org. Chem.* **2005**, *70*, 10701–10708.

(20) Article published during review process of this article: Mercalli, V.; Nyadanu, A.; Cordier, M.; Tron, G. C.; Grimaud, L.; El Kaim, L. *Chem. Commun.* **2017**, *53*, 2118–2121.

(21) Kikugawa, Y.; Aoki, Y.; Sakamoto, T. *J. Org. Chem.* **2001**, *66*, 8612–8615.

(22) (a) Mullard, A. *Nat. Rev. Drug Discovery* **2013**, *12*, 173–175. (b) Karawajczyk, A.; Hamza, D.; Kalliokoski, T.; Pouwer, K.; Morgentini, R.; Nelson, A.; Müller, G.; Piechot, D.; Tzalis, D. *Drug Discovery Today* **2015**, *20*, 1310–1316. (c) Paillard, G.; Cochrane, P.; Jones, P. S.; Caracoti, A.; van Vlijmen, H.; Pannifer, A. D. *Drug Discovery Today* **2016**, *21*, 97–102. (d) Besnard, J.; Jones, P. S.; Hopkins, A. L.; Pannifer, A. D. *Drug Discovery Today* **2015**, *20*, 181–186. (e) Nelson, A.; Roche, D. *Bioorg. Med. Chem.* **2015**, *23*, 2613.