

University of Groningen

Clinical examination for diagnosing circulatory shock

Hiemstra, Bart; Eck, Ruben J; Keus, Frederik; van der Horst, Iwan C.C.

Published in:
Current opinion in critical care

DOI:
[10.1097/MCC.0000000000000420](https://doi.org/10.1097/MCC.0000000000000420)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hiemstra, B., Eck, R. J., Keus, F., & van der Horst, I. C. C. (2017). Clinical examination for diagnosing circulatory shock. *Current opinion in critical care*, 23(4), 293-301. DOI: 10.1097/MCC.0000000000000420

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Clinical examination for diagnosing circulatory shock

Bart Hiemstra, Ruben J. Eck, Frederik Keus, and Iwan C.C. van der Horst

Purpose of review

In the acute setting of circulatory shock, physicians largely depend on clinical examination and basic laboratory values. The daily use of clinical examination for diagnostic purposes contrasts sharp with the limited number of studies. We aim to provide an overview of the diagnostic accuracy of clinical examination in estimating circulatory shock reflected by an inadequate cardiac output (CO).

Recent findings

Recent studies showed poor correlations between CO and mottling, capillary refill time or central-to-peripheral temperature gradients in univariable analyses. The accuracy of physicians to perform an educated guess of CO based on clinical examination lies around 50% and the accuracy for recognizing a low CO is similar. Studies that used predefined clinical profiles composed of several clinical examination signs show more reliable estimations of CO with accuracies ranging from 81 up to 100%.

Summary

Single variables obtained by clinical examination should not be used when estimating CO. Physician's educated guesses of CO based on unstructured clinical examination are like the 'flip of a coin'. Structured clinical examination based on combined clinical signs shows the best accuracy. Future studies should focus on using a combination of signs in an unselected population, eventually to educate physicians in estimating CO by using predefined clinical profiles.

Keywords

cardiac output, circulatory shock, clinical examination, critical illness, diagnostic accuracy, physical examination, shock

INTRODUCTION

Many critically ill patients suffer from circulatory shock, which places them at increased risks of multi-organ failure, long-term morbidity and mortality [1,2]. Combinations of clinical, hemodynamic and biochemical variables are recommended for diagnosing shock [3,4].

Daily use of clinical examination (in any patient) for diagnostic purposes contrasts with the limited number of studies, so that the level of evidence in the critically ill is considered best practice [4]. Much remains unknown about the value of clinical examination in diagnosing shock, reflected by an inadequate cardiac output (CO) or maldistribution of blood flow. More knowledge on this topic could assist physicians in the diagnostic process and guide interventions. Previous overviews have evaluated the value of physical examination in sepsis patients [5], cardiovascular patients [6^{••}] and in hemodynamically unstable patients for predicting fluid responsiveness [7[•]]. We aim to provide an overview of the diagnostic test accuracy of clinical

examination findings for estimating CO in critically ill patients.

BACKGROUND

'Clinical examination' of the cardiovascular system has been performed for a long time. The first evaluations of heart rate by palpation of the arterial pulse date back as far as approximately 335–280 B.C.

Department of Critical Care, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

Correspondence to Iwan C.C. van der Horst, Department of Critical Care, University of Groningen, University Medical Center Groningen, Hanzeplein 1, P.O. Box 30.001, 9713 GZ Groningen, The Netherlands. Tel: +31 50 361 5617; e-mail: i.c.c.van.der.horst@umcg.nl

Curr Opin Crit Care 2017, 23:293–301

DOI:10.1097/MCC.0000000000000420

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

KEY POINTS

- Clinical examination findings are poorly associated with CO in single-variable and multivariable analyses.
- The physician's accuracy to subjectively estimate CO based on clinical examination equals the flip of a coin.
- Physicians are likely insufficiently capable to recognize a low CO by using clinical examination.
- Estimating CO by using a predefined combination of clinical signs seems the most accurate method to diagnose shock.
- Future studies on estimating CO should be conducted in a representative population, use standardized clinical examination and use appropriate statistical indices of diagnostic accuracy.

[8]. Around the second century A.D., physicians recognized the value of pulse rate in diagnosing diseases. Pulse quality and quantity were extensively evaluated and distinctions were made in pulse fullness, rate, rhythm and size [9]. However, it would still take hundreds of years before the clinical assessment of circulatory shock 'had evolved' into the way as it is conducted today. In 1941, Ebert *et al.* [10] elaborately described the complexity of symptoms seen in systemic and peripheral circulatory failure in septic shock patients. He encountered the same clinical picture that we still face today:

(..) All the patients studied presented a similar clinical picture. They were stuporous or comatose. The rectal temperatures ranged from 36.1 to 41.3 degrees Celsius. The skin was pale and often covered with perspiration. The extremities were cold, and this finding usually preceded the fall in arterial pressure. The skin of the body was usually warm, although in terminal stages it too became cool. The radial pulse was feeble or impalpable. The pulse rate was rapid. (..)

For years, clinical examination was considered the cornerstone for diagnosing shock. Reliance on examination declined when Swan *et al.* [11] introduced pulmonary artery catheterization (PAC) in 1970. PAC allowed a wide range of pressure and flow-based hemodynamic measurements, including variables such as pulmonary capillary wedge pressure, systemic vascular resistance and CO [12]. Several studies concluded that the use of PAC frequently resulted in change of therapy compared with clinical examination [13–18]. However, PAC remained controversial because of its invasiveness in the absence of any clinical benefit [19–22].

Today, PAC has largely been replaced by less-invasive methods for assessment of CO, ranging from echo to pulse pressure analysis devices [23–26].

Despite these technological improvements, clinical examination still holds a prominent position in diagnosing circulatory shock [4,27]. We aimed to provide an overview of the diagnostic accuracy of clinical examination for the assessment of circulatory shock measured by CO or cardiac index (CI). We only included studies that estimated CO using clinical examination based on a one-time snapshot. Physicians mostly use changes in clinical examination findings as proxy for changes in CO to guide their interventions. To evaluate the diagnostic accuracy of changes in clinical examination in relation to changes in CO was beyond the scope of this review. In this review, we were mainly interested which clinical examination findings may accommodate clinical needs, because in daily practice these snapshot measurements guide treatment decisions as triggers for interventions.

METHODS

A sensitive search strategy was used to identify eligible studies (Appendix 1, <http://links.lww.com/COCC/A17>). In addition, we used the snowball and citation search methods on the selected articles. We attempted to include all studies that provided results on clinical examination findings in relation to CO. We excluded prognostic studies. We separated studies that evaluated univariable associations from studies that used multivariable analyses. Varying statistical indices for describing diagnostic test accuracy as well as a varying prevalence of low CO were encountered, limiting interstudy comparison. Whenever available, we used likelihood ratios as the preferred modality to describe diagnostic accuracy. Likelihood ratios may provide valuable information on disease probability in an individual and do not change with pretest probability (i.e. the prevalence of disease) [28–30]. We calculated sensitivity, specificity, predictive values and likelihood ratios of clinical examination for the detection of low CO whenever possible.

RESULTS

Our search resulted in 8128 hits of which 28 publications were selected. An additional six publications were identified through snowballing. After selection, we included 34 publications in this overview.

UNIVARIABLE STUDIES

Thirteen studies evaluated univariable associations of clinical examination variables with CO, including

skin temperature or temperature gradients ($n=8$) [31–38], capillary refill time (CRT; $n=1$) [39], temperature gradient and CRT ($n=1$) [40], mottling ($n=1$) [41], heart rate and mean arterial pressure ($n=1$) [42] and central venous pressure ($n=1$; Table 1) [31–43]. The method used for measuring CO varied, including, for example thermodilution with the PAC or Doppler wave with transesophageal or transthoracic echocardiography.

Circulatory shock may lead to compensatory vasoconstriction of nonvital, peripheral tissues such as the skin. Peripheral perfusion can easily be evaluated by measurement of skin temperature, CRT and degree of skin mottling. Two studies demonstrated that a subjectively cool skin temperature was associated with a lower CO [31,32]. Studies evaluating the correlation between objective temperature measurements and CO showed conflicting results; some observed moderate correlations [33,35,40], whereas most observed no correlation [34–38]. Skin temperature measurement methods differ widely and are likely influenced by several factors: age, ambient temperature, hypothermia, peripheral vascular disease, vasopressors, pain and anxiety have all been proposed as influencing circumstances [44,45]. This may explain the conflicting results and may limit its usefulness for estimating CO in clinical practice. Several studies have emphasized the prognostic value of prolonged CRT and mottling of the skin [39,41,46–49], but only three studies have evaluated their associations with CO and found no relevant correlations [39–41].

Prospective studies on systemic hemodynamic variables showed that heart rate, mean arterial pressure and central venous pressure were not directly correlated to CO [42,43,50]. Only during episodes of deep hypotension, one study observed a moderate correlation between mean arterial pressure and CO [42]. These systemic hemodynamic variables seem to be poor indicators of CO, which supports the common conception that low blood pressure is a late sign of circulatory shock and should not be relied on for early diagnosis [4,51].

MULTIVARIABLE STUDIES

Twenty-one studies evaluated multivariable associations of clinical variables with CO. Because of the differing methods of estimating CO, we subdivided our results into studies that evaluated the capacity of physicians to estimate CO ($n=17$; Table 2) [13–18, 52–61, 62^{***}] and studies that constructed clinical profiles based on multiple variables ($n=3$) or a multivariable model ($n=1$) to correlate clinical examination findings with CO (Table 3) [63–66]. Furthermore, we could calculate the diagnostic test

accuracy for physician's estimation of low CO in nine studies (Table 2).

PHYSICIAN'S CAPACITY TO ESTIMATE CO BASED ON CLINICAL EXAMINATION

Seventeen studies evaluated the accuracy of physician's estimates or 'educated guesses' of CO as compared to objectively measured CO. Estimates were based on clinical examination, with or without knowledge of medical history, biochemical values and/or radiological imaging (Table 2). Some studies used a categorical variable for CO estimates (e.g. 'low', 'normal' or 'high'), whereas others used a continuous scale (e.g. 1–12 l per min) [15,17, 62^{***}]. Physician's estimates were correct in 42–62% of the time [13–18, 52–61]. Moderate-to-reasonable correlations and a high percentage error were found when physician's estimates of continuous CO were compared to objectively measured CO [15,16, 62^{***}]. Moderate-to-very poor agreements were found in studies that used weighted κ statistics to address agreement occurring by chance [55,59,60,67]. In addition, two studies reported that 21 and 26% of the CO estimations were completely disparate (an estimated high CO when the objective CO was low or vice versa) [55,59].

Nine studies provided enough data for calculation of the diagnostic accuracy of physician's estimates for detecting low CO. The overall results appeared disappointing [13,14,16,17,53,54,56,58, 60] (Table 2). Furthermore, two studies concluded that physicians more frequently overestimated (31–33%) rather than underestimated (18–23%) CO [14,57], implicating that physicians were more prone to miss an insufficient CO. Perel *et al.* [62^{***}] found the opposite when physicians were asked to estimate CO on a continuous scale.

These results suggest that physicians are not very capable to subjectively estimate CO based on clinical examination. The widely varying diagnostic accuracies are probably the result of different populations or cutoffs for a low CO, but overall it seems that physician's estimates are 'an inaccurate diagnostic test'. This is in accordance with two studies of Saugel *et al.* [67,68], which both demonstrate the incapability of physicians to reliably assess volume status using simple clinical signs. Furthermore, five out of six studies concluded that predictions of senior staff members were equally bad as those of residents or fellows [13,18,54,61,62^{***},69]. Finally, one study found that the accuracy of estimates was unrelated to the level of confidence physicians had in their assessment [69].

Several important limitations apply. Many studies did not elaborate their methods of clinical

Table 1. Prediction of cardiac output using a single variable

Author, year	Patients	Population	Variables of interest	Measurement method	Results	
					Nonsignificant	Significant
Peripheral temperature						
Kaplan <i>et al.</i> 2001 [31]	264 ^a	Surgical ICU patients	Temp, subjective: foot ('cool' or 'warm')	PAC, technique not mentioned	–	'Cool' : $CI=2.9 \pm 1.2$ 'Warm' : $CI=4.3 \pm 1.2$
Schey <i>et al.</i> 2009 [32]	10 ^a	Post cardiac surgery	Temp, subjective: foot: ('cool' or 'cool-warm' or 'warm') Temp, objective of foot	PAC, thermomodulation	T_{skin} , objective: $r=0.11$	'Cool' : $CO=3.71$ 'Cool-warm': $CO=4.83$ 'Warm' : $CO=5.12$
Joly <i>et al.</i> 1969 [33]	100	Circulatory shock	Temp, objective: toe ΔT : toe – ambient (ΔT_{p-a})	Indicator dilution technique	–	T_{skin} , objective: $r=0.71$ ΔT_{p-a} : $r=0.73$
Woods <i>et al.</i> 1987 [34]	26 ^a	Circulatory shock	ΔT : central – toe (ΔT_{c-p})	PAC, thermomodulation	ΔT_{c-p} : no correlation	
Vincent <i>et al.</i> 1988 [35]	15 ^a	Cardiogenic and septic shock	ΔT : toe – ambient (ΔT_{p-a})	PAC, thermomodulation	ΔT_{p-a} in septic shock: no correlation	ΔT_{p-a} in cardiogenic shock: $r=0.63$
Bailey <i>et al.</i> 1990 ^b [40]	40 ^a	Post cardiac surgery	ΔT : central – toe (ΔT_{c-p})	PAC, thermomodulation	ΔT_{c-p} day of operation: no correlation	ΔT_{c-p} postoperative day 1: $r=-0.60$
Sommers <i>et al.</i> 1995 [36]	21 ^a	Post cardiac surgery	T_{skin} , objective: axillary, groin, knee, ankle, toe	PAC, thermomodulation	T_{skin} , objective: no correlation in any site	–
Boerma <i>et al.</i> 2008 [37]	35	Sepsis and septic shock	ΔT : central – foot (ΔT_{c-p})	TEE, Doppler wave	ΔT_{c-p} : $r=-0.15$	–
Bourcier <i>et al.</i> 2016 [38]	103 ^a	Sepsis and septic shock	ΔT : toe – ambient (ΔT_{p-a})	TTE, technique not mentioned	ΔT_{p-a} : no correlation	–
Capillary refill time						
Bailey <i>et al.</i> 1990 ^b [40]	40 ^a	Post cardiac surgery	CRT: site not mentioned	PAC, thermomodulation	CRT: no correlation	–
Ait-Ofella <i>et al.</i> 2014 [39]	59	Septic shock	CRT: index finger	FloTrac, arterial pressure waveform analysis	CRT: no correlation	–
Skin mottling						
Ait-Ofella <i>et al.</i> 2011 [41]	60	Septic shock	Mottling score: knee	TTE, Doppler wave	Mottling score: no correlation	–
Systemic hemodynamic variables						
Wo <i>et al.</i> 1993 [42]	256 ^a	Severe injury and critically ill postoperative	HR, MAP	PAC, thermomodulation	HR: $r=0.27$, $r^2=0.07$, MAP: $r=-0.01$, $r^2=0.0001$,	MAP during severe hypotension: $r=0.50$, $r^2=0.25$
Kuntischer <i>et al.</i> 2006 [43]	16 ^a	Major burns	Central venous pressure	Thermal dye double indicator dilution	–	Central venous pressure: $r=0.40$

^a=repeated measurements in each patient.^b=same study population. ΔT_{c-p} , central-to-peripheral temperature gradient ($^{\circ}C$); ΔT_{p-a} , peripheral-to-ambient temperature gradient ($^{\circ}C$); CI , cardiac index ($l/min/m^2$); CO , cardiac output (l/min); CRT, capillary refill time (s); HR, heart rate (beats/min); MAP, mean arterial pressure (mmHg); PAC, pulmonary artery catheter; TEE, transoesophageal echocardiography; Temp, temperature ($^{\circ}C$); TTE, transthoracic echocardiography.

Table 2. Physician's capacity to estimate cardiac output based on clinical examination

Author, year	Patients	Setting	Variables of interest		Measurement method	Estimation	Results
			Classification	Estimation based on			
Connors <i>et al.</i> 1983 [13]	62 ^a	ICU	CI categorical: < 2.5; 2.5–3.5; > 3.5	Clinical assessment, laboratory and X-ray	PAC, thermodilution	44% correct estimation	Sens 58% (45–68%); Spec 60% (48–71%) PPV 58% (49–65%); NPV 60% (52–67%) LR+ 1.43 (1.02–2.00); LR– 0.71 (0.51–0.98)
Eisenberg <i>et al.</i> 1984 [14]	97	ICU	CO categorical: < 4.5; 4.5–7.5; > 7.5	Not described	PAC, thermodilution	51% correct estimation	Sens 71% (54–85%); Spec 56% (43–69%) PPV 48% (39–57%); NPV 78% (66–86%) LR+ 1.64 (1.15–2.33); LR– 0.51 (0.29–0.89)
Tuchschmidt <i>et al.</i> 1987 [15]	35	ICU	CO continuous	Clinical assessment and X-ray	PAC, thermodilution	$r = 0.72$	–
Connors <i>et al.</i> 1990 [17]	461	ICU	CI dichotomous: < 2.2; ≥ 2.2 CI continuous	Clinical assessment, laboratory, X-ray and ECG	PAC, thermodilution	64% correct estimation Mean CI difference in CI = 1.0 ± 0.9	Sens 49% (40–57%); Spec 70% (65–75%) PPV 43% (38–49%); NPV 74% (71–77%) LR+ 1.62 (1.28–2.05); LR– 0.73 (0.62–0.87)
Caloria <i>et al.</i> 1990 ^b [16]	114	Surgical ICU	CO categorical: < 4; 4–8; > 8	Clinical assessment, laboratory and X-ray	PAC, thermodilution	51% correct estimation $r = 0.47$	Sens 67% (30–93%); Spec 80% (71–87%) PPV 22% (14–34%); NPV 97% (92–99%) LR+ 3.33 (1.83–6.07); LR– 0.42 (0.16–1.05)
Steingrub <i>et al.</i> 1991 ^b [53]	152	Surgical and medical ICU	CO categorical: < 4; 4–8; > 8	Clinical assessment, laboratory and X-ray	PAC, thermodilution	51% correct estimation	Sens 54% (37–70%); Spec 73% (63–81%) PPV 40% (31–51%); NPV 82% (76–87%) LR+ 1.96 (1.29–2.98); LR– 0.64 (0.44–0.91)
Mimoz <i>et al.</i> 1994 [18]	112	ICU	Combinations of CI, PAOP and SVRI	Clinical assessment, laboratory, X-ray and echocardiography	PAC, thermodilution	56% correct estimation	–
Staudinger <i>et al.</i> 1998 [54]	149	ICU	CI categorical: < 2.0; 2.0–4.0; > 4.0	Clinical assessment, medical history, laboratory and X-ray	PAC, thermodilution	62% correct estimation	–
Rodriguez <i>et al.</i> 2000 [55]	33	ED + respiratory distress or hypotension	CI categorical: < 2.6; 2.6–4.0; > 4.0	Clinical assessment, medical history, laboratory, X-ray and ECG	TEE, Doppler wave	$\kappa 1 = -0.04$ (95% CI –0.31–0.24) $\kappa 2 = 0.07$ (95% CI –0.17–0.31)	–
Linton <i>et al.</i> 2002 [56]	50	Post cardiac surgery	CI categorical: < 1.9; 1.9–3.5; > 3.5	Not described	LiDCO, indicator-dilution	54% correct estimation	Sens 42% (15–72%); Spec 74% (57–87%) PPV 33% (18–54%); NPV 80% (71–87%) LR+ 1.58 (0.67–3.72); LR– 0.79 (0.47–1.32)
Iregui <i>et al.</i> 2003 [57]	105	ICU	CI categorical: < 2.5; 2.5–4.5; > 4.5	Clinical assessment, laboratory and X-ray	TEE, Doppler wave	44% correct estimation	–
Veale <i>et al.</i> 2005 [58]	68	ICU	CI categorical: < 2.5; 2.5–4.2; > 4.5	Not described	BioZ CO monitor, impedance cardiography	42% correct estimation	Sens 22% (6–48%); Spec 66% (51–79%) PPV 19% (8–38%); NPV 70% (63–76%) LR+ 0.65 (0.25–1.68); LR– 1.18 (0.86–1.62)
Rodriguez <i>et al.</i> 2006 [59]	31	ED + endotracheal intubation	CI categorical: ranges not specified	Clinical assessment, medical history, laboratory and X-ray	TEE, Doppler wave	$\kappa = 0.57$ (95% CI 0.36–0.77)	–
Nowak <i>et al.</i> 2011 [60]	38	ED + respiratory distress	CO categorical: < 4.0; 4.0–8.0; > 8.0	Clinical assessment and medical history	Nexfin, ABP waveform analysis	50% correct estimation $\kappa = -0.02$ (95% CI –0.25–0.20)	Sens 33% (4–78%); Spec 63% (44–79%) PPV 14% (5–36%); NPV 83% (73–90%) LR+ 0.89 (0.26–3.00); LR– 1.07 (0.57–2.00)
Duan <i>et al.</i> 2014 [61]	132	ICU	CI categorical: < 3; 3–5; > 5	Not described	PiCCO, thermodilution	50% correct estimation	–
Peter <i>et al.</i> 2016 [62 ^{***}]	206 ^a	ICU	CO continuous	Clinical assessment	PiCCO, thermodilution	Percentage error = 66% Absolute mean difference in CO = –1.5 ± 2.2	–

^a = repeated measurements in each patient.^b = overlapping study populations.95% CI, 95% confidence intervals; CI, cardiac index (l/min/m²); CO, cardiac output (l/min); ECG, electrocardiography; ICU, intensive care unit; LiDCO, lithium dilution cardiac output; LR–, negative likelihood ratio; LR+, positive likelihood ratio; NPV, negative predictive value; PAC, pulmonary artery catheter; PAOP, pulmonary artery catheter; pulse contour cardiac output; PPV, positive predictive value; Sens, sensitivity; Spec, specificity; SVRI, systemic vascular resistance index (dynes · s/cm⁵ · min²); TEE, transesophageal echocardiography.

Table 3. Combined signs of clinical examination for estimation of CO

Author, year	Patients	Population	Variables of interest			CO-measurement	Results
			Clinical profile	Clinical profile based on			
Combined clinical profiles							
Ramo <i>et al.</i> 1970 [63]	98	AMI	I (normal CI): no signs of HF II (normal CI): mild-to-moderate HF III (low CI): overt pulmonary edema IV (low CI): cardiogenic shock	Mean arterial pressure, cool extremities, urine output, mental status, third heart sound gallop rhythm and rales		PAC, indicator-dilution technique	I (normal CI): 23 of 45 (51%) II (normal CI): 19 of 30 (63%) III (low CI): 10 of 10 (100%) IV (low CI): 13 of 13 (100%)
Forrester <i>et al.</i> 1977 [64]	200	AMI	I (normal CI): no pulmonary congestion or peripheral hypoperfusion II (normal CI): pulmonary congestion only III (low CI): hypoperfusion only IV (low CI): both	Heart rate, blood pressure, cool extremities, urine output and mental status		PAC, thermodilution	Overall: 81% correct estimations of CI I & II (normal CI): 84 of 95 (88%) III & IV (low CI): 76 of 105 (72%)
Grissom <i>et al.</i> 2009 [65]	405	ALI	I: All three clinical signs aberrant II: Any one clinical sign aberrant	Capillary refill time, knee mottling and cool extremities		PAC, thermodilution	92% correct estimations of CI in class I: Sens 12% (3–28%); Spec 98% (97–99%) PPV 40% (17–69%); NPV 93% (92–93%) LR+ 7.52 (2.23–25.3); LR– 0.89 (0.79–1.01) 75% correct estimations of CI in class II: Sens 52% (34–69%); Spec 78% (73–82%) PPV 17% (12–23%); NPV 95% (93–96%) LR+ 2.31 (1.58–3.38); LR– 0.62 (0.44–0.89)
Multivariable analysis							
Sasse <i>et al.</i> 1996 [66]	23 ^a	ICU patients	CO continuous	Heart rate, respiratory rate, mean arterial pressure and temperature		PAC, thermodilution	Heart rate: $R^2 = 0.05$ Respiratory rate: $R^2 = 0.14$ Mean arterial pressure: $R^2 = 0.03$

^a=repeated measurements in each patient.ALI, acute lung injury; AMI, acute myocardial infarction; CI, cardiac index (l/min/m²); CO, cardiac output (l/min); HF, heart failure; LR–, negative likelihood ratio; LR+, positive likelihood ratio; NPV, negative predictive value; PAC, pulmonary artery catheter; PPV, positive predictive value; Sens, sensitivity; Spec, specificity.

examination in terms of variables used and definitions employed, leaving variability at the physician's discretion so that these studies cannot be reproduced. PAC was used in most studies, but only in selected patients who failed to respond to initial therapy or in whom clinical examination alone was deemed insufficient, so that evaluation of the accuracy of clinically estimated *CO* will be biased by definition. Likewise, many other studies also used convenience samples, which hampers generalizability of their results. Clinical examination should be performed in a standardized fashion, according to a protocol, to maximize interobserver agreement and generalizability.

COMBINED SIGNS OF CLINICAL EXAMINATION FOR ESTIMATION OF *CO*

Three studies have compared predefined clinical profiles based upon clinical examination with objectively measured *CI* (Table 3). Forrester *et al.* [64] found a good agreement in patients with acute myocardial infarction (AMI). In their study, 75% of patients with low *CI* and 96% of patients with very low *CI* had clinical signs of peripheral hypoperfusion, such as decreased skin temperature, confusion or oliguria in conjunction with either arterial hypotension or tachycardia. Ramo *et al.* [63] observed 100% correct estimation of low *CI* when patients with AMI had overt signs of pulmonary edema or signs of cardiogenic shock. In their study, clinical signs of overt pulmonary edema were defined by rales or a third heart sound gallop rhythm and cardiogenic shock was diagnosed by the presence of a systolic blood pressure below 90 mmHg, oliguria, cold extremities and disorientation. These findings suggest that physicians can diagnose cardiogenic shock in patients with AMI using clinical examination. Accurate estimation of *CO* for diagnosing shock in all critically ill patients based on clinical examination might appear much more difficult because of large interindividual differences. Grissom *et al.* [65] combined CRT, mottling and skin temperature to predict *CI* in an unselected cohort of patients with acute lung injury. The presence of all three physical signs had a high specificity (98%) but a low sensitivity (12%) for diagnosing shock, suggesting that these three signs accurately rule in, but inaccurately rule out circulatory shock. Varying types of shock are probably associated with varying clinical signs [70], so that a 'one size fits all' approach seems inappropriate. Roughly, one-third of all patients with circulatory shock suffer from a low *CO*, whereas two-thirds have distributive shock with associated high *CO* [1,71]. Especially in the latter, clinical examination may indicate

inadequate circulation regardless of the height of *CO* and it is difficult to establish how much *CO* is sufficient for each individual patient.

PREDICTING *CO* USING A MULTIVARIABLE MODEL

One study used multivariable regression analyses to estimate *CO* based on heart rate, respiratory rate, mean arterial pressure and central temperature (Table 3) [66]. These multivariable results confirm that systemic hemodynamic variables do not correspond well with *CO*. Future diagnostic studies of *CO* should therefore incorporate all clinical and hemodynamic variables in a multivariable model.

CONCLUSION

Clinical examination findings are poorly associated with *CO* in single-variable and multivariable analyses. Physicians seem to be insufficiently capable to estimate *CO* or recognize a low *CO* using their clinical examination. The most promising results were found when *CO* was estimated by using predefined profiles composed of combined clinical examination signs. However, most studies were conducted in highly selected populations and the details of estimations were not specified. On the basis of current evidence, using clinical examination to diagnose *CO* can, to our opinion, not be considered best practice. Future studies on this topic should be conducted in a representative population, use standardized clinical examination and use appropriate statistical indices of diagnostic accuracy. Ultimately, these results should guide education of physicians to estimate *CO* using predefined clinical profiles.

Acknowledgements

None.

Financial support and sponsorship

None.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. De Backer D, Biston P, Devriendt J, *et al.* Comparison of dopamine and norepinephrine in the treatment of shock. *N Engl J Med* 2010; 362:779–789.
2. Vincent JL, Marshall JC, Namendys-Silva SA, *et al.* Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *Lancet Respir Med* 2014; 2:380–386.

3. De Backer D, Donadello K, Sakr Y, *et al*. Microcirculatory alterations in patients with severe sepsis: impact of time of assessment and relationship with outcome. *Crit Care Med* 2013; 41:791–799.
 4. Cecconi M, De Backer D, Antonelli M, *et al*. Consensus on circulatory shock and hemodynamic monitoring. Task force of the European Society of Intensive Care Medicine. *Intensive Care Med* 2014; 40:1795–1815.
 5. Postelnicu R, Evans L. Monitoring of the physical exam in sepsis. *Curr Opin Crit Care* 2017; 23:232–236.
 6. Elder A, Japp A, Verghese A. How valuable is physical examination of the cardiovascular system? *BMJ* 2016; 354:i3309.
- This review extensively elaborates the diagnostic accuracy of physical examination of the cardiovascular system.
7. Bentzer P, Griesdale DE, Boyd J, *et al*. Will this hemodynamically unstable patient respond to a bolus of intravenous fluids? *JAMA* 2016; 316:1298–1309.
- This review provides a comprehensive overview of studies that measure hypovolemia.
8. Billman GE. Heart rate variability: a historical perspective. *Front Physiol* 2011; 2:86.
 9. Bedford DE. The ancient art of feeling the pulse. *Br Heart J* 1951; 13:423–437.
 10. Ebert RV, Stead EA. Circulatory failure in acute infections. *J Clin Invest* 1941; 20:671–679.
 11. Swan HJ, Ganz W, Forrester J, *et al*. Catheterization of the heart in man with use of a flow-directed balloon-tipped catheter. *N Engl J Med* 1970; 283:447–451.
 12. Ganz W, Donoso R, Marcus HS, *et al*. A new technique for measurement of cardiac output by thermodilution in man. *Am J Cardiol* 1971; 27:392–396.
 13. Connors AF Jr, McCaffree DR, Gray BA. Evaluation of right-heart catheterization in the critically ill patient without acute myocardial infarction. *N Engl J Med* 1983; 308:263–267.
 14. Eisenberg PR, Jaffe AS, Schuster DP. Clinical evaluation compared to pulmonary artery catheterization in the hemodynamic assessment of critically ill patients. *Crit Care Med* 1984; 12:549–553.
 15. Tuschmidt J, Sharma OP. Impact of hemodynamic monitoring in a medical intensive care unit. *Crit Care Med* 1987; 15:840–843.
 16. Celoria G, Steingrub JS, Vickers-Lahti M, *et al*. Clinical assessment of hemodynamic values in two surgical intensive care units. Effects on therapy. *Arch Surg* 1990; 125:1036–1039.
 17. Connors AF Jr, Dawson NV, Shaw PK, *et al*. Hemodynamic status in critically ill patients with and without acute heart disease. *Chest* 1990; 98:1200–1206.
 18. Mimoz O, Rauss A, Rekik N, *et al*. Pulmonary artery catheterization in critically ill patients: a prospective analysis of outcome changes associated with catheter-prompted changes in therapy. *Crit Care Med* 1994; 22:573–579.
 19. Robin ED. Death by pulmonary artery flow-directed catheter. Time for a moratorium? *Chest* 1987; 92:727–731.
 20. Shah MR, Hasselblad V, Stevenson LW, *et al*. Impact of the pulmonary artery catheter in critically ill patients: meta-analysis of randomized clinical trials. *JAMA* 2005; 294:1664–1670.
 21. Marik PE. Obituary: pulmonary artery catheter 1970 to 2013. *Ann Intensive Care* 2013; 3:38–43.
 22. Rajaram SS, Desai NK, Kalra A, *et al*. Pulmonary artery catheters for adult patients in intensive care. *Cochrane Database Syst Rev* 2013; (2):CD003408.
 23. Vincent JL, Rhodes A, Perel A, *et al*. Clinical review: update on hemodynamic monitoring – a consensus of 16. *Crit Care* 2011; 15:229.
 24. Alhashemi JA, Cecconi M, Hofer CK. Cardiac output monitoring: an integrative perspective. *Crit Care* 2011; 15:214.
 25. Schloghofer T, Gilly H, Schima H. Semi-invasive measurement of cardiac output based on pulse contour: a review and analysis. *Can J Anaesth* 2014; 61:452–479.
 26. Teboul JL, Saugel B, Cecconi M, *et al*. Less invasive hemodynamic monitoring in critically ill patients. *Intensive Care Med* 2016; 42:1350–1359.
 27. Sevransky J. Clinical assessment of hemodynamically unstable patients. *Curr Opin Crit Care* 2009; 15:234–238.
 28. Sackett DL. The rational clinical examination. A primer on the precision and accuracy of the clinical examination. *JAMA* 1992; 267:2638–2644.
 29. McGee S. Simplifying likelihood ratios. *J Gen Intern Med* 2002; 17:646–649.
 30. Sackett D, Simel D. A primer on the precision and accuracy of the clinical examination & an updated summary. In: Keitz S, editor. *The rational clinical examination: evidence-based clinical diagnosis*. New York: McGraw-Hill Professional; 2009. pp. 1–16.
 31. Kaplan LJ, McPartland K, Santora TA, Trooskin SZ. Start with a subjective assessment of skin temperature to identify hypoperfusion in intensive care unit patients. *J Trauma* 2001; 50:620–627; discussion 627–8.
 32. Schey BM, Williams DY, Bucknall T. Skin temperature as a noninvasive marker of hemodynamic and perfusion status in adult cardiac surgical patients: an observational study. *Intensive Crit Care Nurs* 2009; 25:31–37.
 33. Joly HR, Weil MH. Temperature of the great toe as an indication of the severity of shock. *Circulation* 1969; 39:131–138.
 34. Woods I, Wilkins RG, Edwards JD, *et al*. Danger of using core/peripheral temperature gradient as a guide to therapy in shock. *Crit Care Med* 1987; 15:850–852.
 35. Vincent JL, Moraine JJ, van der Linden P. Toe temperature versus transcutaneous oxygen tension monitoring during acute circulatory failure. *Intensive Care Med* 1988; 14:64–68.
 36. Sommers MS, Stevenson JS, Hamlin RL, Ivey TD. Skin temperature and limb blood flow as predictors of cardiac index. *Clin Nurs Res* 1995; 4: 22–37.
 37. Boerma EC, Kuiper MA, Kingma WP, *et al*. Disparity between skin perfusion and sublingual microcirculatory alterations in severe sepsis and septic shock: a prospective observational study. *Intensive Care Med* 2008; 34:1294–1298.
 38. Bourcier S, Pichereau C, Boelle PY, *et al*. Toe-to-room temperature gradient correlates with tissue perfusion and predicts outcome in selected critically ill patients with severe infections. *Ann Intensive Care* 2016; 6: 63–71.
 39. Ait-Oufella H, Bige N, Boelle PY, *et al*. Capillary refill time exploration during septic shock. *Intensive Care Med* 2014; 40:958–964.
 40. Bailey JM, Levy JH, Kopel MA, *et al*. Relationship between clinical evaluation of peripheral perfusion and global hemodynamics in adults after cardiac surgery. *Crit Care Med* 1990; 18:1353–1356.
 41. Ait-Oufella H, Lemoine S, Boelle PY, *et al*. Mottling score predicts survival in septic shock. *Intensive Care Med* 2011; 37:801–807.
 42. Wo CC, Shoemaker WC, Appel PL, *et al*. Unreliability of blood pressure and heart rate to evaluate cardiac output in emergency resuscitation and critical illness. *Crit Care Med* 1993; 21:218–223.
 43. Kuntscher MV, Germann G, Hartmann B. Correlations between cardiac output, stroke volume, central venous pressure, intra-abdominal pressure and total circulating blood volume in resuscitation of major burns. *Resuscitation* 2006; 70:37–43.
 44. Kholoussy AM, Sufian S, Pavlides C, Matsumoto T. Central peripheral temperature gradient. Its value and limitations in the management of critically ill surgical patients. *Am J Surg* 1980; 140:609–612.
 45. Schey BM, Williams DY, Bucknall T. Skin temperature and core-peripheral temperature gradient as markers of hemodynamic status in critically ill patients: a review. *Heart Lung* 2010; 39:27–40.
 46. Champion HR, Sacco WJ, Hannan DS, *et al*. Assessment of injury severity: the triage index. *Crit Care Med* 1980; 8:201–208.
 47. van Genderen ME, Lima A, Akkerhuis M, *et al*. Persistent peripheral and microcirculatory perfusion alterations after out-of-hospital cardiac arrest are associated with poor survival. *Crit Care Med* 2012; 40: 2287–2294.
 48. Coudroy R, Jamet A, Frat JP, *et al*. Incidence and impact of skin mottling over the knee and its duration on outcome in critically ill patients. *Intensive Care Med* 2015; 41:452–459.
 49. de Moura EB, Amorim FF, da Cruz Santana AN, *et al*. Skin mottling score as a predictor of 28-day mortality in patients with septic shock. *Intensive Care Med* 2016; 42:479–480.
 50. Lattik R, Couture P, Denault AY, *et al*. Mitral Doppler indices are superior to two-dimensional echocardiographic and hemodynamic variables in predicting responsiveness of cardiac output to a rapid intravenous infusion of colloid. *Anesth Analg* 2002; 94:1092–1099; table of contents.
 51. van Genderen ME, Bartels SA, Lima A, *et al*. Peripheral perfusion index as an early predictor for central hypovolemia in awake healthy volunteers. *Anesth Analg* 2013; 116:351–356.
 52. Connors AF Jr, Dawson NV, McCaffree R, *et al*. Assessing hemodynamic status in critically ill patients: do physicians use clinical information optimally? *J Crit Care* 1987; 2:174–180.
 53. Steingrub JS, Celoria G, Vickers-Lahti M, *et al*. Therapeutic impact of pulmonary artery catheterization in a medical/surgical ICU. *Chest* 1991; 99:1451–1455.
 54. Staudinger T, Locker GJ, Lacizka K, *et al*. Diagnostic validity of pulmonary artery catheterization for residents at an intensive care unit. *J Trauma* 1998; 44:902–906.
 55. Rodriguez RM, Berumen KA. Cardiac output measurement with an esophageal Doppler in critically ill emergency department patients. *J Emerg Med* 2000; 18:159–164.
 56. Linton RA, Linton NW, Kelly F. Is clinical assessment of the circulation reliable in postoperative cardiac surgical patients? *J Cardiothorac Vasc Anesth* 2002; 16:4–7.
 57. Iregui MG, Prentice D, Sherman G, *et al*. Physicians' estimates of cardiac index and intravascular volume based on clinical assessment versus transthoracic Doppler measurements obtained by critical care nurses. *Am J Crit Care* 2003; 12:336–342.
 58. Veale WN Jr, Morgan JH, Beatty JS, *et al*. Hemodynamic and pulmonary fluid status in the trauma patient: are we slipping? *Am Surg* 2005; 71:621–625; discussion 625–6.
 59. Rodriguez RM, Lum-Lung M, Dixon K, Nothmann A. A prospective study on esophageal Doppler hemodynamic assessment in the ED. *Am J Emerg Med* 2006; 24:658–663.
 60. Nowak RM, Sen A, Garcia AJ, *et al*. The inability of emergency physicians to adequately clinically estimate the underlying hemodynamic profiles of acutely ill patients. *Am J Emerg Med* 2012; 30:954–960.
 61. Duan J, Cong LH, Wang H, *et al*. Clinical evaluation compared to the pulse indicator continuous cardiac output system in the hemodynamic assessment of critically ill patients. *Am J Emerg Med* 2014; 32:629–633.

62. Perel A, Saugel B, Teboul JL, *et al.* The effects of advanced monitoring on hemodynamic management in critically ill patients: a pre and post questionnaire study. *J Clin Monit Comput* 2016; 30:511–518.

This large multicenter study included a broad ICU cohort and used sophisticated statistical measures to correlate CO to clinical examination.

63. Ramo BW, Myers N, Wallace AG, *et al.* Hemodynamic findings in 123 patients with acute myocardial infarction on admission. *Circulation* 1970; 42:567–577.
64. Forrester JS, Diamond GA, Swan HJ. Correlative classification of clinical and hemodynamic function after acute myocardial infarction. *Am J Cardiol* 1977; 39:137–145.
65. Grissom CK, Morris AH, Lanken PN, *et al.* Association of physical examination with pulmonary artery catheter parameters in acute lung injury. *Crit Care Med* 2009; 37:2720–2726.
66. Sasse SA, Chen PA, Mahutte CK. Relationship of changes in cardiac output to changes in heart rate in medical ICU patients. *Intensive Care Med* 1996; 22:409–414.
67. Saugel B, Ringmaier S, Holzapfel K, *et al.* Physical examination, central venous pressure, and chest radiography for the prediction of transpulmonary thermodilution-derived hemodynamic parameters in critically ill patients: a prospective trial. *J Crit Care* 2011; 26:402–410.
68. Saugel B, Wagner JY, Wendon J, Perel A. Getting the full diagnostic picture in intensive care medicine: a plea for 'physiological examination'. *Ann Am Thorac Soc* 2015; 12:1738–1739.
69. Dawson NV, Connors AF Jr, Speroff T, *et al.* Hemodynamic assessment in managing the critically ill: is physician confidence warranted? *Med Decis Making* 1993; 13:258–266.
70. Vazquez R, Gheorghe C, Kaufman D, Manthous CA. Accuracy of bedside physical examination in distinguishing categories of shock: a pilot study. *J Hosp Med* 2010; 5:471–474.
71. Sakr Y, Reinhart K, Vincent JL, *et al.* Does dopamine administration in shock influence outcome? Results of the Sepsis Occurrence in Acutely Ill Patients (SOAP) Study. *Crit Care Med* 2006; 34:589–597.