

University of Groningen

## Modulation of content and composition of glucosinolates in Brassica upon abiotic stress

Aghajanzadeh, Tahereh A.; De Kok, L.J.

*Published in:*  
Berichte aus dem Julius Kühn-Institut

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2017

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Aghajanzadeh, T. A., & De Kok, L. J. (2017). Modulation of content and composition of glucosinolates in Brassica upon abiotic stress. *Berichte aus dem Julius Kühn-Institut*, 191, 7.

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## **Modulation of content and composition of glucosinolates in *Brassica* upon abiotic stress**

**Tahereh A. Aghajanzadeh<sup>1</sup> and Luit J. De Kok<sup>2</sup>**

<sup>1</sup>*Department of Biology, Faculty of Basic Science, University of Mazandaran, Babolsar, Iran (E-mail:t.aghajanzade@umz.ac.ir):* <sup>2</sup>*Laboratory of Plant Physiology, Groningen Institute for Evolutionary Life Sciences, University of Groningen, P.O. Box 11103, 9700 CC Groningen, the Netherlands*

Plants elicit multiple responses when exposed to a variety of biotic and abiotic stress factors. These stress factors induce signaling cascades that activate ion channels, kinases, production of reactive oxygen species (ROS), and accumulation of plant hormones. These signals affect eventually both the primary and secondary metabolism resulting in a substantial variation in the plant metabolome. Plant secondary metabolism shows a high phenotypically plasticity in response to both biotic and abiotic stress factors. Glucosinolates are secondary sulfur compounds, which occur in high levels in Brassica vegetables and which are responsible for their characteristic flavor and odor and maybe involved in the defense against insects and pathogens. Moreover, they also have high nutraceutical and pharmacological value. Currently more than 130 different glucosinolates have been identified in plants and more than 30 of them are present in Brassica species. Their content and composition in plants is strongly affected during plant development and is affected by various environmental factors, *viz.* nutrient availability (S, N, K, Se and B) temperature, drought, UV-B, as well as fungal and bacterial pathogens. In the current study, the impact of chloride and sulfate salinity and high levels of Cu, Zn and Ni on the content and composition of glucosinolates was investigated in seedlings of *Brassica* species.