

University of Groningen

Real-time conformational changes and controlled orientation of native proteins inside a protein nanoreactor

Van Meervelt, Veerle; Soskine, Misha; Singh, Shubham; Schuurman-Wolters, Gea; Wijma, Hein J; Poolman, Bert; Maglia, Giovanni

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/jacs.7b10106](https://doi.org/10.1021/jacs.7b10106)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Van Meervelt, V., Soskine, M., Singh, S., Schuurman-Wolters, G., Wijma, H. J., Poolman, B., & Maglia, G. (2017). Real-time conformational changes and controlled orientation of native proteins inside a protein nanoreactor. *Journal of the American Chemical Society*, 139(51), 18640-18646. [jacs.7b10106]. <https://doi.org/10.1021/jacs.7b10106>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Real-Time Conformational Changes and Controlled Orientation of Native Proteins Inside a Protein Nanoreactor


Veerle Van Meervelt,^{†,‡,Ⓜ} Misha Soskine,[‡] Shubham Singh,[‡] Gea K. Schuurman-Wolters,[‡] Hein J. Wijma,[‡] Bert Poolman,[‡] and Giovanni Maglia^{*,‡,Ⓜ}

[†]Department of Chemistry, University of Leuven, Leuven B-3001, Belgium

[‡]Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Groningen 9747 AG, The Netherlands

Supporting Information

ABSTRACT: Protein conformations play crucial roles in most, if not all, biological processes. Here we show that the current carried through a nanopore by ions allows monitoring conformational changes of single and native substrate-binding domains (SBD) of an ATP-Binding Cassette importer in real-time. Comparison with single-molecule Förster Resonance Energy Transfer and ensemble measurements revealed that proteins trapped inside the nanopore have bulk-like properties. Two ligand-free and two ligand-bound conformations of SBD proteins were inferred and their kinetic constants were determined. Remarkably, internalized proteins aligned with the applied voltage bias, and their orientation could be controlled by the addition of a single charge to the protein surface. Nanopores can thus be used to immobilize proteins on a surface with a specific orientation, and will be employed as nanoreactors for single-molecule studies of native proteins. Moreover, nanopores with internal protein adaptors might find further practical applications in multianalyte sensing devices.



INTRODUCTION

Protein conformational dynamics play key roles in molecular recognition^{1–3} and regulation of protein activity.⁴ The lack of understanding of macromolecule dynamics is a fundamental problem in today's biochemistry. Single-molecule techniques allow probing the conformational heterogeneity and structural dynamics of proteins in real-time. However, such approaches typically require labeling with fluorophores and/or tethering proteins to a surface, which in turn may perturb the very same dynamic behavior that is intended to be studied.⁵

Ionic currents through individual nanopores can be employed to sample the activity of unlabeled molecules at the single-molecule level. Initial work focused on observing the traversing of polymers across single nanopores,^{6,7} or the binding⁸ or reacting of small molecules with a nanopore in real-time.⁹ More recently, nanopore currents have been employed to sample enzymatic reactions.¹⁰ Notable examples are the visualization of the intermediate mechanical steps during an helicase catalyzed unwinding of DNA,¹¹ or the sampling of the kinetic intermediates of the GroEL-assisted protein refolding reaction with an engineered co-chaperon GroES-nanopore.¹²

A more generic approach to nanopore enzymology, however, is sampling proteins inside a nanopore. Work with solid-state nanopores showed that many features of proteins that are rapidly diffusing across a nanopore, including their approximate shape, dipole moment and conformation can be identified.^{13,14} We reported that single proteins can be incorporated and sampled inside Cytolysin A (ClyA) biological nanopores from *Salmonella typhi* (Figure 1) for tens of seconds,^{15–19} which

allowed sampling the ubiquitination of proteins in solution¹⁹ or the binding of ligands to the proteins inside the nanopore.^{15,20}

Here, we use substrate-binding domains (SBD1 and SBD2) of the ABC importer Gln PQ from *Lactococcus lactis* (Figure 1) as model systems to probe the conformational dynamics of proteins within a nanopore. SBD1 and SBD2 differ in amino acid specificity and dynamics of closing and opening upon ligand binding as described previously.²¹ Substrate-binding proteins undergo large conformational changes upon binding to their ligands. SBD1 and SBD2 have been previously characterized by several analytical techniques,^{22,23} including single-molecule Förster resonance energy transfer (smFRET),²¹ making them ideal to characterize this new experimental setup. We found that nanopore currents can efficiently report the conformational changes of SBD1 and SBD2. Remarkably, the kinetic constants for ligand binding to the proteins inside the nanopore are almost identical to those measured in solution. Thus, nanopores can be used as nanoreactors to report the conformational dynamics of proteins lodged inside the nanopore in real-time. Our technique does not require bulky tags and proteins are sampled with high bandwidth for tens of seconds, which is challenging in single-molecule fluorescence experiments.

Received: September 27, 2017

Published: December 5, 2017




Figure 1. Trapping proteins inside the ClyA nanopore. (Left) Surface representation of Type I ClyA-AS (blue) in the bilayer (gray) with SBD1 (PDB-ID = 4KPT) in open conformation lodged inside the nanopore. (Right) SBD2 in open (PDB-ID = 4KR5) and closed (PDB-ID = 4KQP) conformation. SBDs are colored according to the residue type: basic residues are colored blue, acidic residues red, polar residues green, and nonpolar residues white. Created with VMD²⁴ while ClyA was created by homology modeling using the *E. coli* ClyA crystal structure.^{16,25}

RESULTS AND DISCUSSION

Conformational Dynamics of SBD1 Inside ClyA Nanopores. SBD1 (74 nM) added to the *cis* side of single type I ClyA-AS nanopores (hereafter ClyA, Figure 1) entered the nanopore under negative applied potentials. The longest average residence time of SBD1 was under -60 mV (4.2 ± 1.8 s, $n = 291$ events, Figure 2A). Individual blockades showed a main current level L_O with a residual current percent ($I_{res\%} = I_B / I_O \times 100$, with I_B the blocked pore and I_O the open pore current level) equal to $67.6 \pm 0.1\%$ ($N = 5$ independent nanopore experiments). The blockades switched with a frequency of 0.7 ± 0.2 s⁻¹ to a second level (L_C , $I_{res\%} = 66.8 \pm 0.4\%$, Figure 2B). Then L_C switched back to L_O with an average rate of 15.1 ± 4.1 s⁻¹ ($n = 441$). The addition of asparagine (0.4 μ M, Figure 2B) to the *cis* side of the nanopore increased the frequency of L_C blockades; and at 50 μ M asparagine L_C was the most prominent current blockade (Figure 2B). By contrast, SBD1(E184W), a variant that does not show any asparagine transport in *L. lactis* cells (Figure S1 of the Supporting Information, SI), showed only one single level ($I_{res\%} = 66.8 \pm 0.1\%$, $N = 3$) that did not respond to asparagine (up to 1 mM, Figure 2C). Together, these results suggest that L_O corresponds to the open conformation of SBD1 and L_C to the closed asparagine-bound state of the protein.²¹ The current blockades allowed calculating the equilibrium dissociation constant (K_d^{app}) by plotting the relative dwell times of the closed population [$L_C / (L_C + L_O)$, Figure 2D]. The K_d^{app} calculated by this method (0.47 ± 0.03 μ M) is similar to the values obtained by isothermal titration calorimetry (ITC, 0.2 μ M) and smFRET (0.35 μ M).²¹ The ligand-induced closing rate constant ($k_{closing} = 1.4 \pm 0.8 \times 10^7$ s⁻¹ M⁻¹) could be determined from the L_O to L_C transition and was similar to the values measured by smFRET (2.2×10^7 s⁻¹ M⁻¹, Figure 2E).²¹ The L_C to L_O transition showed a slight variation with the ligand concentration (Figure 2E). At high asparagine concentration the transition rate (4.8 ± 0.5 s⁻¹, 2.8 μ M

asparagine) was similar to the rate constants measured by smFRET (4.2 s⁻¹).²¹

Conformational Dynamics of SBD2 Inside ClyA Nanopores. The average residence time of SBD2 inside ClyA (3.9 ± 0.7 s, $n = 225$) was similar to that of SBD1, albeit at -100 mV. Contrary to SBD1, however, individual blockades showed three levels: L_{OA} ($I_{res\%} = 61.9 \pm 0.3\%$, rate = 3.6 ± 0.4 s⁻¹, $n = 2900$, $N = 3$), L_{OB} ($I_{res\%} = 59.3 \pm 0.2\%$, rate = 6.1 ± 1.3 s⁻¹, $n = 2900$, $N = 3$, Figure 3A), and L_C ($I_{res\%} = 60.7 \pm 0.2\%$, $k_{opening} = 55.6 \pm 3.1$ s⁻¹, $n = 856$, $N = 3$). The addition of the substrate glutamine increased the frequency of L_C current levels, and reduced the dwell time of the L_{OA} and L_{OB} levels (Figure 3A). When the inactive D417F variant of SBD2²¹ was used instead of SBD2, L_C events were not observed at any glutamine concentration tested (up to 200 μ M, Figure 3B). Therefore, L_C reflects the glutamine-bound closed conformation of SBD2 from which the equilibrium dissociation constant ($K_d^{app} = 0.83 \pm 0.10$ μ M, Figure 3C) could be measured. The K_d^{app} for internalized SBD2 proteins corresponds well to the values measured in solution by smFRET and ITC experiments ($K_d^{app} = 1.1 \pm 0.1$ μ M and $K_d^{app} = 0.9 \pm 0.2$ μ M, respectively).²¹ As observed for SBD1, the resulting opening rate showed a slight decrease with the ligand concentration (Figure 3D). At high ligand concentration the opening rate ($k_{opening} = 31.8 \pm 3.7$ s⁻¹, 2.3 μ M glutamine) was about 2-fold faster than the value measured by smFRET ($k_{opening} = 17.2$ s⁻¹). The rate constants measured from L_{OA} and L_{OB} were almost identical ($4.3 \pm 0.1 \times 10^7$ s⁻¹ M⁻¹ and $4.2 \pm 0.1 \times 10^7$ s⁻¹ M⁻¹, respectively), and corresponded well to the closing rate constant measured with smFRET (3.8×10^7 s⁻¹ M⁻¹), suggesting that they are related to the closing of SBD2.²¹

SBD2 Has a Fixed Orientation Inside ClyA That Can Be Controlled. L_{OA} and L_{OB} could reflect the protein lodging inside ClyA at a deep and a shallow residence site within the nanopore, as it was previously shown for Dendra2 fluorescent protein,¹⁵ AlkB²⁶ and thrombin.^{15,17} However, for these proteins the relative time spent between the two sites was




Figure 2. Conformational dynamics of SBD1 inside ClyA-AS. (A) Typical ionic current blockades provoked by the capture of SBD1 (74 nM, *cis*) by the ClyA-AS nanopore at -60 mV. The open pore (I_O) and SBD-blocked (L) current levels are indicated, with L_O and L_C corresponding to the open and closed state of SBD1. (B) Details of SBD1 current blockade before and after addition of $0.40 \mu\text{M}$ and $50 \mu\text{M}$ asparagine (added *cis*). (C) Typical current blockades of the inactive SBD1 (E184W) variant before and after addition of $50 \mu\text{M}$ asparagine showing no ligand-induced current blockades. (D) The K_d^{app} value of SBD1 for asparagine was obtained from the areas of the histograms of the open (L_O) and ligand-bound (L_C) populations as $[L_C/(L_C + L_O)]$. The K_d^{app} is the concentration of substrate at 50% signal saturation. (E) Dependency of the opening and closing rates of SBD1 on asparagine concentration. The data in E were fitted to eq S5C (opening rates) and S6C (closing rates) as described in the SI. Current traces were collected in 150 mM NaCl, 15 mM Tris-HCl, pH 7.5 at 24°C by applying a Bessel low-pass filter with a 2 kHz cutoff and sampled at 10 kHz. A postacquisition Gaussian filter of 100 Hz was applied. Experiments were performed at -60 mV.

strongly voltage-dependent, and the difference in residual current between the two levels was large (between 13.6% and 56.5%). By contrast, the relative distribution of L_{OA} and L_{OB} showed no voltage dependence (Figure S2) and relatively small ΔI_{res} values ($2.6 \pm 0.4\%$, -100 mV). Alternatively, L_{OA} and L_{OB} might reflect two different SBD2 open conformations. However, previous smFRET experiments²¹ did not show any indication for the presence of two open conformations. In addition, glutamine binding rate constants measured from L_{OA} and L_{OB} were identical (Figure 3C), which suggests they both reflect the same open conformation. Further, the inactive SBD2 (D417F), which rests solely in the open conformation²¹ still showed L_{OA} and L_{OB} blockades. Hence, it is unlikely that L_{OA} and L_{OB} correspond to two different open conformations of SBD2. We propose that L_{OA} and L_{OB} reflect two orientations of SBD2 inside the nanopore. SBD2 has two globular domains or lobes (lobe A, residues 253–343 and 441–481, lobe B, residues 348–437) connected by a flexible linker (residues 344–347 and 438–440, Figure 4A). The two lobes of SBD2 do not show a large imbalance in the spatial distribution of the

surface charges as shown by the protein dipole moment that is aligned to the latitudinal axis of the protein (Figures 4B and S3). Thus, within the electric field inside ClyA the probability of SBD2 to be oriented with either lobes toward the *cis* or *trans* opening is likely similar (Figure 4A). In contrast, SBD1 has its dipole moment aligned to the longitudinal axis of the protein (Figure S3), suggesting that inside the nanopore SBD1 may be oriented. This is consistent with the observation that SBD1 shows one main current level (Figure 2A, B). Note that as measured from the ratio of the area of the histogram of L_{OA} and L_{OB} in a SBD2 blockade, orientation A is about 1.7-fold more represented than orientation B (Figure 4B, top), despite the dipole of SBD2 would predict a 50% distribution between the two orientations. This effect is possibly related to the asymmetry of the ClyA nanocavity, which is formed by chiral amino acids that might favor one of the two orientations of SBD2.

Next, we imposed asymmetry between the lobes of SBD2 via point mutations that grafted a charged residue at the apical positions in each lobe of the protein surface. We reasoned that




Figure 3. Orientation and dynamics of SBD2 measured by nanopore experiments. (A) Typical current blockade provoked by the capture of SBD2 (72 nM, *cis*) by the ClyA-AS nanopore. Left, apo-SBD2, middle current blockades after the addition of 0.40 μM of glutamine (*cis*); and right, blockades after adding 50 μM glutamine (*cis*). (B) Typical current blockades provoked by the capture of inactive SBD2(D417F) (70 nM) before and after the addition of 200 μM glutamine to the *cis* side. Red asterisks represent the restoration of the open pore current after SBD2(D417F) exited from the pore. (C) K_d^{app} value of SBD2 for glutamine obtained by fitting to a binding isotherm, using the relative closed population [$L_C/(L_{OA} + L_{OB} + L_C)$] at the indicated substrate concentrations. (D) Opening and closing rates of SBD2 versus the glutamine concentration. The data was fitted by eq S5C (opening rates) and S6C (closing rates) as described in the SI. Experiments were performed at -100 mV in 150 mM NaCl, 15 mM Tris-HCl, pH 7.5 at 24 $^\circ\text{C}$ by applying a Bessel low-pass filter with a 2 kHz cutoff and sampled at 10 kHz. A postacquisition Gaussian filter of 100 Hz was then applied.

tuning the dipole moment of the protein would allow the electric field inside the nanopore to align the protein with one preferred orientation. Residues in both lobes were selected *in silico* by supercharging of SBD2, followed by molecular dynamics simulations in order to minimize the impact on protein structure (Figure 4C and SI). Two nonconserved residues that were furthest apart from the center of mass of SBD2 were selected: T256 on lobe A and S358 on lobe B (Figure 4C). In remarkable agreement to our view, when a negative charged residue was introduced to lobe A (T256E) or a positive charged residue was introduced in lobe B (S358 K), blockades provoked by SBD2 variants were mainly L_{OA} (Figure 4B, Tables S1 and S2). Accordingly, inverse arrangement of the additional charges resulted in mainly L_{OB} blockades (Figure 4B, Tables S1 and S2). All tested variants could bind glutamine, although the SBD2 variants that favored L_{OB} orientation, showed a slightly reduced affinity (Figures S4 and S5, additional discussion in SI). Therefore, L_{OA} and L_{OB} current levels describe two orientations of SBD2 relative to ClyA, with L_{OA} corresponding to lobe A facing the *cis* side of ClyA (orientation A) and L_{OB} corresponding to the opposite orientation (orientation B, Figure 4A, B).

SBD1 and SBD2 Adopt Multiple Open and Closed Conformations. Two textbook mechanisms are often evoked to explain molecular recognition in proteins: Koshland's "induced fit" hypothesis²⁷ and the "conformational selection" model.^{28,29} The "induced fit" model treats the unliganded protein as if it exists in a low-energy, single, and stable

conformation that switches to the liganded-conformation upon binding the substrate. However, NMR and crystallographic studies revealed that proteins are inherently dynamic and may sample a vast ensemble of conformations even in the absence of ligands,³⁰ which may play important roles in molecular recognition.²⁸ The "conformational selection" model accommodates this thermally accessed conformational heterogeneity by arguing that weakly populated, higher energy conformations recognize and bind ligands with high(er) affinity with subsequent population shift toward these conformers.³¹ However, experimental evidence to prove models of substrate recognition is difficult, mainly because the ability of measuring the affinity of ligands to multiple protein structures can only be obtained using single-molecule methods.

Several crystal structures of substrate-binding proteins revealed that they may exist in at least four states or conformations: a ligand-free (O) or ligand-bound (OL) open state, and a ligand-free (C) or ligand-bound (CL) closed state.^{32–36} The O, CL, and the C states of SBD1 and SBD2 were directly observed by both single-molecule FRET²¹ and nanopore analysis (Figures 2 and 3). At saturating substrate concentrations (50 μM) an open state of SBD1 and SBD2 could still be observed (Figures 2B, 3A, and S6), suggesting that the OL state also exists for SBD1 and SBD2. Open-liganded states were not observed by smFRET at 50 μM ligand concentration, possibly because the sampling rate was close to the resolution of the event.

Notably, the nanopore analysis also revealed that the opening rates of SBD1 and SBD2 changed with the

two closed states (Scheme 1 and SI). Fitting the kinetic data for SBD1 and SBD2 to kinetic equations derived from such model (SI) revealed that both SBD proteins open about two-folds slower and close about 100- to 1000-folds faster in the presence of the ligand (Scheme 1). In turn this suggests a mechanism in which the ligands induce a conformational change that results in the closing of the protein around the ligand.

ClyA Nanopores As Nanoreactors for Single Molecule Enzymology. The results presented in this work show that nanopore currents can be used to sample the conformational dynamics of proteins. This approach samples native proteins, does not require expensive equipment or lengthy sample preparations, and allows long recordings at high sampling frequency. Molecular dynamics simulations suggested that the stability and activity of a folded macromolecule in a confined space are different from bulk because of entropic effects.^{37–41} Therefore, the kinetics and thermodynamics of proteins trapped inside the nanopore might be different from bulk, although experimental evidence for these predictions is scarce. Surprisingly, we found that inside the nanopore the conformational dynamics of SBD1 and SBD2 were similar to that in solution, suggesting that the effect of confinement for these proteins is negligible. In any case, even if other proteins inside the nanopore will show a different dynamics behavior than in bulk, it should be noticed that the cell is highly crowded, and the interaction of proteins with nearby macromolecules can impact protein stability and influence the recognition of ligands.^{42,43} Thus, the interaction of proteins with the nanopore inner surface may actually reproduce more faithfully the crowded environment that proteins face inside cells than the highly diluted conditions that are sampled with other single-molecule techniques.

Inside the nanopore proteins experience a relatively strong electric field ($\sim 10^6$ V/m). The effect of electric fields on protein stability, activity and orientation is not well understood, with some reports suggesting that proteins might even unfold when subjected to relatively moderate EFs (5×10^2 V/m),⁴⁴ or inside nanopores.⁴⁵ However, inside cells high transmembrane fields ($2\text{--}3 \times 10^7$ V/m) preserve the nonequilibrium chemical steady-state or allow the cell-to-cell signaling in complex organisms. Our results suggest that electric fields in the order of 10^6 V/m do not have a strong effect on the activity and dynamics of proteins with weak dipoles such as SBD1 and SBD2. Nonetheless, we also showed that inside nanopores proteins can be oriented, and switching a single charge can control such orientation. This is intriguing, because recently it has been proposed that a relatively strong ($5 \times 10^5\text{--}3 \times 10^6$ V/m) electric field exists inside the cytoplasm of cells.^{46,47} Thus, in-cell proteins might be oriented or sorted by intracellular electric fields, with post-translational modification such as phosphorylation controlling such processes.

CONCLUSIONS

In summary, we show that nanopore currents can be used to monitor the conformational dynamics of proteins. This approach is simple, low-cost, requires only microgram amounts of proteins, and allows high bandwidth sampling with no intrinsic limitations on the observation time. Importantly, it is also the first method that allows the analysis of native proteins at the single-molecule level. We found that the dynamics, kinetics and equilibrium constants of substrate binding to SBD1 and SBD2 inside the nanopore were almost identical to those in solution, suggesting that the electric field and nanoconfinement

of water inside the nanopore most likely have no effect on the activity or stability of proteins. The latter finding is surprising, considering that the transport of water in nanoconfined geometries is expected to differ from bulk phase.⁴⁸ In any case, proteins inside the nanopore are likely to mimic more faithfully the tightly packed environment inside cells and might be used to sample other biological confined reactions such as the refolding of proteins inside the GroEL nanochamber.

This innovative technology is likely to have applications in basic and applied science. The single-molecule data showed that the proteins adopt multiple open and closed conformations. The kinetic analysis is compatible with a mechanism in which the binding of ligands to the open protein conformation triggers a conformational change that induces the closing of the protein around the substrate. Further, we showed unambiguously that the proteins inside the nanopore align with the electric field, and that a single change in the protein surface charge can switch the orientation. Thus, the orientation and subsequent activity of proteins inside cellular electric fields can be controlled post-translationally by a single modification. Finally, we show here that nanopores are capable of transducing protein conformational changes and ligand binding into electrical signals. Thus, nanopores with a panel of protein adaptors might be integrated in low-cost and portable devices for real-time detection of biologically relevant analytes from biological samples.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.7b10106.

Additional results and discussion, materials and methods, figures, tables of ligands SBD1 and SBD2, and additional references (PDF)

AUTHOR INFORMATION

Corresponding Author

*g.maglia@rug.nl

ORCID

Veerle Van Meervelt: 0000-0002-2727-367X

Giovanni Maglia: 0000-0003-2784-0811

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

V.V.M. thanks the Research Foundation Flanders (FWO) for the doctoral fellowship. The work of G.M. was funded by an ERC starting grant (SMEN, # 260884). The work of B.P. was funded by a NWO TOPGO (#L.10.060) and ERC Advanced grant (ABCVolume, #670578). We thank Joris M.H. Goudsmits for performing the autocorrelation analysis.

REFERENCES

- (1) Boehr, D. D.; Nussinov, R.; Wright, P. E. *Nat. Chem. Biol.* **2009**, *5* (11), 789.
- (2) Popovych, N.; Sun, S.; Ebright, R. H.; Kalodimos, C. G. *Nat. Struct. Mol. Biol.* **2006**, *13* (9), 831.
- (3) Carroll, M. J.; Mauldin, R. V.; Gromova, A. V.; Singleton, S. F.; Collins, E. J.; Lee, A. L. *Nat. Chem. Biol.* **2012**, *8* (3), 246.
- (4) Tzeng, S.-R.; Kalodimos, C. G. *Nature* **2009**, *462* (7271), 368.
- (5) Toseland, C. P. *J. Chem. Biol.* **2013**, *6* (3), 85.

- (6) Bezrukov, S. M.; Vodyanoy, I.; Parsegian, V. A. *Nature* **1994**, *370* (6487), 279.
- (7) Kasianowicz, J. J.; Brandin, E.; Branton, D.; Deamer, D. W. *Proc. Natl. Acad. Sci. U. S. A.* **1996**, *93* (24), 13770.
- (8) Bayley, H.; Cremer, P. S. *Nature* **2001**, *413* (6852), 226.
- (9) Shin, S. H.; Luchian, T.; Cheley, S.; Braha, O.; Bayley, H. *Angew. Chem., Int. Ed.* **2002**, *41* (19), 3707.
- (10) Willems, K.; Van Meervelt, V.; Wloka, C.; Maglia, G. *Philos. Trans. R. Soc., B* **2017**, *372*, 20160230.
- (11) Craig, J. M.; Laszlo, A. H.; Brinkerhoff, H.; Derrington, I. M.; Noakes, M. T.; Nova, I. C.; Tickman, B. I.; Doering, K.; de Leeuw, N. F.; Gundlach, J. H. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114*, 11932.
- (12) Ho, C.-W.; Van Meervelt, V.; Tsai, K.-C.; De Temmerman, P.-J.; Mast, J.; Maglia, G. *Sci. Adv.* **2015**, *1* (11), e1500905.
- (13) Yusko, E. C.; Bruhn, B. R.; Eggenberger, O. M.; Houghtaling, J.; Rollings, R. C.; Walsh, N. C.; Nandivada, S.; Pindrus, M.; Hall, A. R.; Sept, D.; Li, J.; Kalonia, D. S.; Mayer, M. *Nat. Nanotechnol.* **2016**, *12* (4), 360.
- (14) Waduge, P.; Hu, R.; Bandarkar, P.; Yamazaki, H.; Cressiot, B.; Zhao, Q.; Whitford, P. C.; Wanunu, M. *ACS Nano* **2017**, *11*, 570610.1021/acsnano.7b01212
- (15) Soskine, M.; Biesemans, A.; Moeyaert, B.; Cheley, S.; Bayley, H.; Maglia, G. *Nano Lett.* **2012**, *12* (9), 4895.
- (16) Soskine, M.; Biesemans, A.; De Maeyer, M.; Maglia, G. *J. Am. Chem. Soc.* **2013**, *135* (36), 13456.
- (17) Van Meervelt, V.; Soskine, M.; Maglia, G. *ACS Nano* **2014**, *8* (12), 12826.
- (18) Biesemans, A.; Soskine, M.; Maglia, G. *Nano Lett.* **2015**, *15* (9), 6076.
- (19) Wloka, C.; Van Meervelt, V.; Van Gelder, D.; Danda, N.; Jager, N.; Williams, C. P.; Maglia, G. *ACS Nano* **2017**, *11* (5), 4387.
- (20) Soskine, M.; Biesemans, A.; Maglia, G. *J. Am. Chem. Soc.* **2015**, *137* (17), 5793.
- (21) Gouridis, G.; Schuurman-Wolters, G. K.; Ploetz, E.; Husada, F.; Vietrov, R.; de Boer, M.; Cordes, T.; Poolman, B. *Nat. Struct. Mol. Biol.* **2014**, *22* (1), 57.
- (22) Schuurman-Wolters, G. K.; Poolman, B. *J. Biol. Chem.* **2005**, *280* (25), 23785.
- (23) Fulyani, F.; Schuurman-Wolters, G. K.; Žagar, A. V.; Guskov, A.; Slotboom, D. J.; Poolman, B. *Structure* **2013**, *21* (10), 1879.
- (24) Humphrey, W.; Dalke, A.; Schulten, K. *J. Mol. Graphics* **1996**, *14* (1), 33.
- (25) Mueller, M.; Grauschopf, U.; Maier, T.; Glockshuber, R.; Ban, N. *Nature* **2009**, *459* (7247), 726.
- (26) Soskine, M.; Biesemans, A.; Maglia, G. *J. Am. Chem. Soc.* **2015**, *137* (17), 5793.
- (27) Koshland, D. E. *Proc. Natl. Acad. Sci. U. S. A.* **1958**, *44*, 98.
- (28) Frauenfelder, H.; Sligar, S. G.; Wolynes, P. G. *Science (Washington, DC, U. S.)* **1991**, *254* (5038), 1598.
- (29) Ma, B.; Kumar, S.; Tsai, C. J.; Nussinov, R. *Protein Eng., Des. Sel.* **1999**, *12* (9), 713.
- (30) Boehr, D. D.; Nussinov, R.; Wright, P. E. *Nat. Chem. Biol.* **2009**, *5* (11), 789.
- (31) Tsai, C. J.; Ma, B.; Nussinov, R. *Proc. Natl. Acad. Sci. U. S. A.* **1999**, *96* (18), 9970.
- (32) Shouldice, S. R.; Skene, R. J.; Dougan, D. R.; Snell, G.; McRee, D. E.; Schryvers, A. B.; Tari, L. W. *J. Bacteriol.* **2004**, *186* (12), 3903.
- (33) Duan, X.; Quiocho, F. A. *Biochemistry* **2002**, *41* (3), 706.
- (34) Berntsson, R. P.-A.; Smits, S. H. J.; Schmitt, L.; Slotboom, D.-J.; Poolman, B. *FEBS Lett.* **2010**, *584* (12), 2606.
- (35) Flocco, M. M.; Mowbray, S. L. *J. Biol. Chem.* **1994**, *269* (12), 8931.
- (36) Oswald, C.; Smits, S. H. J.; Höing, M.; Sohn-Bösser, L.; Dupont, L.; Le Rudulier, D.; Schmitt, L.; Bremer, E. *J. Biol. Chem.* **2008**, *283* (47), 32848.
- (37) Zhou, H. X. *J. Mol. Recognit.* **2004**, *17* (5), 368.
- (38) Klimov, D. K.; Newfield, D.; Thirumalai, D. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, *99* (12), 8019.
- (39) Cheung, M. S.; Thirumalai, D. *J. Mol. Biol.* **2006**, *357* (2), 632.
- (40) Marino, K. A.; Bolhuis, P. G. *J. Phys. Chem. B* **2012**, *116*, 11872.
- (41) Vaitheeswaran, S.; Thirumalai, D. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105* (46), 17636.
- (42) van den Berg, J.; Boersma, A. J.; Poolman, B. *Nat. Rev. Microbiol.* **2017**, *15* (5), 309.
- (43) Boersma, A. J.; Zuhorn, I. S.; Poolman, B. *Nat. Methods* **2015**, *12* (3), 227.
- (44) Bekard, I.; Dunstan, D. E. *Soft Matter* **2014**, *10* (3), 431.
- (45) Talaga, D. S.; Li, J. *J. Am. Chem. Soc.* **2009**, *131* (26), 9287.
- (46) Tyner, K. M.; Kopelman, R.; Philbert, M. *Biophys. J.* **2007**, *93* (4), 1163.
- (47) Andreev, V. P. *PLoS One* **2013**, *8* (4), e6188410.1371/journal.pone.0061884.
- (48) Majolino, D.; Corsaro, C.; Crupi, V.; Venuti, V.; Wanderlingh, U. *J. Phys. Chem. B* **2008**, *112* (13), 3927.