

Gait & Posture
Original Article

A cross-sectional study of age-related changes in plantar pressure distribution between 4-7 years: a comparison of regional and pixel-level analyses

Jill Phethean¹, Todd C Pataky^{2*}, Christopher J. Nester¹, Andrew H. Findlow¹
**Corresponding author*

¹School of Health Sciences, University of Salford, Manchester, M6 6PU, UK

² Department of Bioengineering, Shinshu University, Tokida 3-15-1 Ueda, Nagano, 386-8567, Japan

Corresponding author contact details:

Todd Pataky, Ph.D.
Shinshu University, Department of Bioengineering
Nagano-ken, Ueda-shi, Tokida 3-15-1
Japan 386-8567
email: tpataky@shinshu-u.ac.jp

Acknowledgements—None.

Abstract

Quantifying morphological and functional development in children's feet, and thereby establishing development norms is difficult. In addition to practical challenges of experimentation on children, measurement equipment like plantar pressure (PP) platforms are almost exclusively geared toward adult-sized feet. These PP quantification problems may be exacerbated by typical regional data analysis techniques, which further reduce spatial resolution. The goal of this study was to quantify PP distributions in developing children, and also to compare the results obtained from typical (regional) techniques with those obtained from a higher-resolution (pixel-level) technique. Ninety-eight children between four and seven years of age were assessed in a cross-sectional design. Maximum PP distributions were collected for each child, and these pressures were linearly regressed against age in days. Present results agree with previous investigations in that maximum pressures and maximum pressure changes occurred in the forefoot. However, results from the present pixel-level technique suggest that these changes are limited to the central metatarsals, and that regional methods can suggest significance where none exists in the actual raw (pixel-level) data, due to signal aliasing and, in particular, to conflation of regional boundaries. We postulate that increased central metatarsal pressures are reflective of the coupling between generalized joint laxity decreases and relatively increasingly inclined central metatarsal bones with age.

Keywords – Foot morphology; Walking; Gait; Morphological and functional development; Children; Statistical parametric mapping

Research Highlights

- We examined age-related plantar pressure changes in young children
- Regional pressures increased with age at the central metatarsals and medial midfoot
- Pixel data disagreed with regional midfoot and heel data and were also more detailed
- Pixel data showed that children's feet tend to become longer and narrower with age
- Pixel-level analysis overcomes regional analysis' inadequate spatial resolution

1. Introduction

Plantar pressure (PP) distributions can be used to characterise foot function and have been shown to change during normal development towards adulthood. For example, several studies report that maximum PP (PP_{\max}) increases under the total foot, the hindfoot and the forefoot between the commencement of independent walking and 13 years of age, but not under the midfoot [1-5]. Age-specific foot loading patterns for toddlers demonstrate the highest PP_{\max} under the hallux; whereas in 7 year olds the highest PP_{\max} is under the hindfoot [2,5]. These clearly identified foot loading patterns are considered to be part of the normal developmental stages of the growing foot [1,4,6].

Whilst there are child-specific variations in the precise timing of growth milestones, the period between four and seven years of age demonstrates considerable neuromuscular and skeletal change [7-9], including changes in foot structure, morphology and function [3,5,10,11]. This period spans the transition from immature to mature gait [12] and changes in bone length and structural composition occur alongside rotational changes in the transverse and frontal planes [7,8,10,11,13]. Changes in gait are seen in the temporal-spatial parameters, and in kinematics, electromyography and kinetics related to ankle joint motion [12,14,15]. Consequently, the changes occurring in this four to seven year old period suggest potential differences in the foot loading patterns within this period.

There are two potential problems with work to-date that both stem from potentially inadequate spatial (anatomical) resolution. This first is that the midfoot and forefoot were previously reported as whole regions of interest (ROI) [3,5], rather than with

mediolateral subdivisions. This is a potential problem because we know that the adult foot demonstrates differences in the mobility of the medial and lateral midfoot and forefoot [16]. Thus smaller ROIs, and their more finely detailed anatomical information, may allow us to draw more definitive conclusions regarding the relation between PP and the maturation of the foot's multiple structures and articulations.

The second potential problem is that previously employed ROIs may have impinged upon neighbouring anatomical structures. For example, midfoot ROI definitions typically extend posteriorly toward the anterior heel and anteriorly toward the posterior forefoot. The problem is that there is typically a large pressure increase from the midfoot to both the anterior heel and the posterior forefoot. Therefore if midfoot ROIs are not carefully defined then midfoot results could be driven by the heel or forefoot, or both. Indeed, this phenomenon has been shown to have the potential to create, eradicate, and even reverse statistical and biomechanical conclusions [17].

We presently refer to the aforementioned problems as “intra-region variation” and “inter-region conflation”, respectively. The cause of both is inadequate spatial resolution in ROI data. Healthy and pathological adult PP data have spatial wavelengths on the order of 30 mm [18] and 20 mm [19], respectively. Plantar pressures must therefore be measured with (Nyquist) resolutions of at least 15 mm and 10 mm, respectively, to avoid signal aliasing. Although original pixel-level data typically have adequate resolutions, on the order of 5 mm, ROI-level data typically have resolutions on the order of 50 mm, which are too gross. Problems associated with inadequate spatial resolution have been addressed elsewhere for adult feet

[18,19], but spatial resolution has not, to our knowledge, been previously explored for children's data.

The purpose of this study was to investigate the effect of spatial resolution on the conclusions regarding PP development amongst children aged four to seven years of age. To this end we presently compare the results of hypothesis testing on pixel-level data vs. ROI-level data. The present null hypothesis was that age has no effect on PP distribution.

2. Methods

2.1. Participants

Subsequent to approval from the ethics committee at the University of Salford 46 girls and 52 boys aged between four and seven years old were recruited from nurseries, schools, university staff and students, and by word of mouth. Although not analyzed categorically, mean (\pm st.dev.) ages for four-, five-, six- and seven-year olds were: 48.5 \pm 1.1, 60.7 \pm 1.0, 72.2 \pm 0.8 and 84.4 \pm 1.0 months, respectively. Parents and guardians gave informed consent in all cases. Since we previously found that gender failed to predict plantar pressure differences amongst children [20] genders were presently pooled.

Children met four criteria to be included: (1) normal gestational period of 37-42 weeks; (2) correct development of locomotion skills; (3) within 0.4th-99.6th centile for height and weight in relation to their chronological age; and (4) no gross gait abnormalities during visual inspection e.g. tiptoe walking, limping, tripping, excessive in/out-toed gait. Normal attainment of locomotion included walking prior to 17

months of age and appropriate age-related development of locomotion skills assessed by movement and balance, and stair tasks in accordance with the ‘Schedule of Growing Skills’ [21]. Stature and weight reference curves [22] were used to determine height and weight centiles.

2.2. Procedure

PP data were collected within 60 days of each child’s birthday, barefoot in mid-gait at natural walking speed [23] using an optical pedobarograph system (Biokinetics Inc, Bethesda, USA; now defunct). The pressure plate surface dimensions, resolution, and sampling frequency were: 57cm x 48.5cm, 3mm x 2mm, and 25 Hz, respectively. A familiarisation period minimised targeting of the plate and determined a start position that ensured five to six steps were taken prior to plate contact. Trials were excluded if the foot failed to contact close to the plate center, and also if gait was observed to be irregular (e.g. targeting, excessive acceleration/deceleration, irregular cadence/stride length). Five trials were collected from each foot.

2.3. Region of Interest (ROI) analysis

Nine anatomical ROIs were manually defined on each PP_{max} image: Heel, MedMF and LatMF (medial and lateral midfoot), Hallux, and MH1-5 (the five metatarsal heads). Constant-sized ellipsoids were used: 3.38cm² (Heel), 2.16cm² (MedMF and LatMF), 1.69cm² (Hallux and MH1), and 1.09 cm² (MH2-MH5). The first set of ROI analyses (“Method A”) merged: (1) the MF region and (2) the MH region (Fig.1a) to mimic previous developmental studies that did not sub-divide the midfoot or metatarsal regions [3,5]. The second set of analyses (“Method B”) assessed the nine ROIs separately (Fig.1b). Although there was little or no foot contact in the medial

midfoot region in some subjects, we employed a medial midfoot ROI to follow convention [24].

Maximum pressure values were extracted from each ROI, were averaged within-subjects, and were then linearly regressed against age. The threshold for significance was set at $\alpha=0.05$, and Bonferroni corrections of 0.013 and 0.00568 were used to correct for multiple comparisons across the four and nine ROIs, respectively. The left and right feet were analysed separately for cross-validation.

2.4 Whole-foot (pixel-level) analysis

PP_{\max} distributions for each step were first registered (i.e. spatially aligned) within-subjects and within-feet using a rapid frequency-based technique [25]. The resulting average distributions (one per-subject, per foot) were registered between-subjects using an optimal linear (scaling) transform. The chronologically first seven-year-old's feet were used as the registration templates, and then, to avoid potential bias from template peculiarities, all images were re-registered to the resulting between-subject mean. A linear transformation was utilized to retain subjects' original morphology.

At this point all images were registered to a common coordinate system, with a total of 196 mean images (one per subject, per foot). Identical to the regional data above, these images were regressed (in a pixel-wise manner) against age, yielding two linear correlation coefficient (r) distributions, one for each foot. The r distributions were then transformed into t distributions via the identity:

$$t = r (n - 2)^{0.5} (1 - r^2)^{-0.5} \quad (1)$$

where n is the number of subjects and t has the Student's t distribution with degrees of freedom: $(n-2)$. Since t is presently both parametric and pixel-specific it is referred to as a 'statistical parametric map' (SPM) [26]. Statistical significance was determined topologically as the probability that a purely random (Gaussian) spatial process, with identical smoothness, will produce suprathreshold pixel-clusters of the observed size [26]; presently a threshold of $|t| > 2$ was selected because this is the lowest threshold that has been shown to be valid for plantar pressure data [27]. Smoothness was estimated from the regression residuals using the average spatial gradient [26].

3. Results

Between-subject average pressure distributions (Fig.2) qualitatively indicate: (i) increased pressure with age, especially at the heel, central metatarsal heads, and hallux, and (ii) a tendency for the foot to become more slender with age (i.e. relatively longer and narrower). Whole-foot peak pressure averaged 258, 280, 293, and 287 kPa (all at the Heel) for the 4-7 yearly intervals on the left (Table 1, Fig.3), with comparable values on the right (Supplementary Material).

Regression analysis of the Method A ROI data (Fig.1a) reached significance only for the MH region, with r^2 values of 0.007, 0.001, 0.077 and 0.001 and p values of 0.403, 0.896, 0.006 and 0.861 for the Heel, MF, MH and Hallux regions, respectively.

Regression analysis of the Method B ROI data (Fig.1b) yielded significant positive correlations between pressure and age in both the left (Fig.3) and right feet (Supplementary Material), but only for MH3 ($p < 0.001$) and MedMF ($p < 0.005$).

Regression analysis of the pixel-level data (Fig.4a) echoed both aforementioned mean-image trends (Fig.2). In particular, pressures tended to be positively correlated with age at the heel, central metatarsals, and hallux. Additionally, the broad rings of negative correlation around the feet (Fig.4a) indicate that the foot tended to become more slender with age. Statistical inference (Fig.4b) confirmed that central metatarsal pressures were positively correlated with age ($p=0.002$), but failed to find significance at either the heel or hallux. The ring of negative correlation observed around both feet reached significance only at the lateral heel in the left foot ($p=0.004$).

4. Discussion

4.1. Discrepancies amongst analysis methods in descriptions of age-related changes

The present midfoot, metatarsals, and heel results were dependent on the analysis method employed. ROI Method A (Fig.1a) failed to find a midfoot effect, but Method B (Fig.1b) found increasing medial midfoot pressures. The former is consistent with qualitative observations of the mean PP distributions (Fig.2), but the latter is not. This apparent discrepancy is explained by inter-region conflation (see Introduction). Since the foot becomes increasingly slender with age (Fig.2; c.f. [5]), we were presently forced to draw the medial midfoot region increasingly more laterally in order to maintain a morphologically consistent mediolateral delineation. That is, the MedMF region can be drawn nearly in line with the lateral edge of the hallux at four years, but it must be positioned more laterally at seven years. Thus the medial midfoot data (Fig.3), instead of truly reflecting pressure changes with age, may actually reflect increasing conflation with the lateral midfoot. The true midfoot trend is observed most clearly at the pixel-level (Fig.4a); the pixel-level data resolve the ROI Method A-B discrepancy.

The present study employed manual ROI definitions, so it is possible that more objective, anatomically-defined ROIs [28,29] may produce more consistent results. However, it has also been shown that midfoot ROI data are less reliable than midfoot pixel-level data [30] because small changes in ROI boundary definitions can greatly affect the underlying pressure values [31]. Thus, while anatomically defined ROIs can be expected to produce more consistency, no ROI is expected to be as consistent as the pixel-level data.

All three methods found metatarsal pressure increases but the interpretations of each are different because their anatomical resolutions are different. For Method A the interpretation is simply that forefoot pressures increase with age, which echoes previous results that employed a full-forefoot ROI [3,5]. For Method B the interpretation, with an uncorrected threshold of $p=0.05$, is that the three medial metatarsal heads' pressure increase with age (Fig.3), and thus that the foot is loaded increasingly medially with age. With a Bonferroni correction of $p=0.00568$, the interpretation would change to the pixel-level interpretation of increased central metatarsal pressures (Fig.4b). The statistical signal clearly peaked at MH3 and tapered off medially and laterally from there; thus any region drawn medial or lateral to MH3 is driven by pressure trends on its border closest to MH3, where the statistical signal is strongest. This can be seen in the regional p values (Fig.3): $p=0.029, 0.009, 0.000, 0.117, 0.992$ for MH1-5, respectively.

None of the three methods found a significant positive correlation between heel pressure and age, and only the pixel-level approach found significant negative

correlation (Fig.4b), but only on the lateral heel periphery. This apparent discrepancy is explained by intra-region variation. The heel clearly becomes relatively thinner with age (Fig.2), a change which is reflected as a cloud of negative correlation around the low-pressure periphery of the foot (Fig.4a). However, this morphological change could not be captured with ROI data because the highest pressures (i.e. the central heel) drives the ROI results. Thus a single heel ROI has insufficient spatial resolution to discriminate amongst multiple, and indeed opposite within-heel trends. Although the pixel-level analyses presently failed to find similar evidence of morphological changes in the right foot, or at any other peripheral foot location in general, the qualitative trend suggests increasingly slenderness with age (Fig.2), and this trend is captured by the raw statistical image (Fig.4a). This increasing slenderness would be expected to reach significance with a longitudinal study because between-subject morphological variability is essentially eliminated in longitudinal analyses.

One may argue that the subjectivity of the present manual ROI definitions may have affected the results. We do not dispute this, but we'd argue that automated or otherwise algorithmic ROI definition is equally susceptible to the aforementioned risks. All ROIs can lie on sharp pressure gradients [17,31] and are thus susceptible to inter-region conflation. Furthermore, no ROI scalar can represent multiple within-ROI trends. More formally, ROI techniques risk aliasing because they sample the PP distribution with insufficient spatial resolution. It has been estimated that statistical signals in PP data have spatial wavelengths on the order of 20 mm [18,19,27]; at this spatial frequency each ROI should have a maximum size of 10 mm to avoid aliasing.

Last, we'd argue that the pixel-level approach offers two non-trivial advantages over the ROI approach. First, its spatial resolution is far-superior (Fig.4 vs. Fig.3), and its anatomical resolution is therefore also superior. Second, pixel-level results are superimposed directly on the foot (Fig.4), so whole-foot effects can be appreciated at a single glance. With the ROI results (Fig.3) it is much more difficult to appreciate how effects are distributed across the foot.

4.2. Comparison with previous studies

The present pressure values match well with three previous studies [2,3,5] (Table 1); we are unaware of other studies that provide data for four-seven year olds. All studies report the greatest pressure under the heel, and higher hallux than forefoot pressure. The main difference between the present and previous study is that the midfoot pressures tended to be lower. A possible explanation is conflation; the cited studies employed automated regionalization, which has been shown elsewhere to be highly susceptible to both anterior heel and posterior midfoot conflation [17,31]. It is possible that the present manually defined ROIs avoided conflation to a greater degree, but since ROI definition information was not provided in the cited studies we are unable to explore this possibility further.

The present forefoot pressures were similar to previous reports [5]. Other studies reported higher forefoot pressure for children aged five years and older [2,3]. The large inter-individual variations in the forefoot region, which were both observed presently (Fig. 3) and reported previously [5] suggest that these discrepancies may simply be an artefact of sampling.

Finally, although we cannot compare our data directly, it is notable that significant age-related pressure increases have been previously reported for the calcaneus and hallux between the ages of one and seven [2]. The absence of similar increases in the present dataset suggests that these changes are likely to occur prior to four years of age.

4.3. Morphological interpretation

The present results (Figs.2,4) support previous reports that the developing foot becomes increasingly slender with age [5]. While the cited study demonstrated this morphological change explicitly via a decrease in the foot's length-width ratio from 0.4 ± 0.02 in four year-olds to 0.38 ± 0.02 in 7 year-olds, the present results are the first, to our knowledge, that report the same result implicitly, via a systematic PP distribution change, and in particular via a cloud of negative correlation around the foot periphery (Fig.4a).

4.4. Functional interpretation

The present results (Fig.4b) suggest that forefoot PP increases [2,3,5] are limited to the central metatarsals. It is possible that this is caused by the interaction amongst three factors: (1) generalized decreases in joint laxity with age [32], (2) increasingly inclined metatarsal bone postures, secondary to a reduction in talar head decline [13], and (3) progression towards adult foot kinematics, which involve less central metatarsal movement compared to the 1st and 5th metatarsals [16,33]. Therefore, this inclined central metatarsal posture coupled with increased joint stiffness would likely increase pressures under the metatarsal heads, and the differences in the central vs.

peripheral metatarsal postures [16,31] could explain the presently observed pressure distribution changes (Figs.2,4).

While the present study focused on maximal plantar pressures, we acknowledge that many other morphological and functional changes occur during maturation of the skeleton and gait between ages 4-7. These are not all fully described in the literature and thus it is not possible to detail how concurrent morphological and functional changes contribute to the observations in this study. However, a variety of other parameters, like center-of-pressure progression, must be considered to fully appreciate the biomechanical bases for the present results.

Conclusion

The present data are broadly in line with previous investigations that report forefoot pressure increases in the developing child's foot as a function of age. These results suggest that PP increases are limited to the central metatarsals, which we speculate is caused by increased central metatarsal joint stiffness secondary to increased weight-bearing demands. The present results comparing regional and pixel methods reiterate previous findings that regional data can distort the underlying PP distribution due to signal aliasing. Since pixel level techniques provide much better anatomical resolution than regional techniques, it is recommended that regionalization should be avoided when analysing PP data, especially in children where pixel sizes are comparatively large with respect to foot size.

References:

- [1] Bosch K, Gerb J, Rosebaum D. Preliminary normative values for foot loading parameters of the developing child. *Gait Posture* 2007; 26:238–247.
- [2] Bosch K, Nagel A, Weigend L, Rosenbaum D. From first to last steps in life—pressure patterns of three generations. *Clin Biomech (Bristol Avon)* 2009; 24:676–681.
- [3] Bosch K, Gerb J, Rosebaum D. Development of healthy children’s feet—Nine-year results of a longitudinal investigation of plantar loading patterns. *Gait Posture* 2010; 32(4):564–571.
- [4] Halleman A, D’Aout K, De Clercq D, Aerts P. Pressure distribution patterns under the feet of new walkers: The first two months of independent walking. *Foot Ankle Intl* 2003; 24(5); 444–453.
- [5] Müller S, Carlsohn A, Müller J, Baur H, Mayer F. Static and dynamic foot characteristics in children aged 1-13 years: a cross-sectional study. *Gait Posture* 2012; 35(3):389-94.

- [6] Bertsch C, Unger H, Winkelmann W, Rosenbaum D. Evaluation of early walking patterns from plantar pressure distribution measurements. First year results of 42 children. *Gait Posture* 2004; 19(4):235-242.
- [7] Anderson M, Blais M, Green WT. Growth of the normal foot during childhood and adolescence; length of the foot and interrelations of foot, stature and lower extremity as seen in serial records of children between 1-18 years of age. *Am J Phys Anthropol* 1956; 14(2):287-308.
- [8] Salenius P, Vankka E. The development of the tibiofemoral angle in children. *J Bone Joint Surg* 1975; 57-A(2):259-261.
- [9] Sobera M, Siedlecka B, Syczewska M. Posture control development in children aged 2-7 years old, based on the changes of repeatability of the stability indices. *Neurosci Lett* 2011; 491(1):13-7.
- [10] Staheli LT. Rotational problems in children. *J Bone Joint Surg* 1993; 75A(6):939-949.
- [11] Pfeffier M, Kotz R, Ledl T, Hauser G, Sluga M. Prevalence of flat-foot in preschool children. *Paediatrics* 2006; 118(2):643-649.
- [12] Sutherland D. The development of mature gait. *Gait Posture* 1997; 6(2):163-170.

- [13] Vanderwilde R, Staheli L, Chew DE, Malagon V. Measurements on radiographs of the foot in normal infants and children. *J Bone Joint Surg Am* 1988; 70(3):407-415.
- [14] Chester VL, Tingley M, Edmund NB. A comparison of kinetic gait parameters for 3-13 year olds. *Clin Biomech* 2006; 21(7):726-732.
- [15] Cupp T, Oeffinger D, Tylkowski C, Augsburg S. Age-related kinetic changes in normal pediatrics. *J Ped Orthop* 1999; 19(4):475-478.
- [16] Lundgren, P, C Nester, A Liu, A, Arndt, R Jones, Stacoff, A, Wolf, P. Lundberg, A. Invasive in vivo measurement of rear, mid and forefoot motion during walking. *Gait Posture* 2008; 28(1):93-100.
- [17] Pataky TC, Goulermas JY. Pedobarographic statistical parametric mapping (pSPM): A pixel-level approach to foot pressure image analysis. *J Biomech* 2008; 41: 2136–2143.
- [18] Pataky TC. Spatial resolution in plantar pressure measurement revisited. *J Biomech* 2012; 45(12):2116–2124.
- [19] Lord M. Spatial resolution in plantar pressure measurement. *Med Eng Phys* 1997; 19(2):140-144.

- [20] Phethean J, Nester CJ. The influence of body weight, body mass index and gender on plantar pressures: Results of a cross-sectional study of healthy children's feet. *Gait Posture* 2012; 36:287-290.
- [21] Bellman M, Longman S, Aukett A. *Schedule of Growing Skills in Practice*. Windsor:NFER-Nelson; 1997.
- [22] Freeman JV, Cole TJ, Chinn S, Jones PRM, White EM, Preece MA. Cross sectional stature and weight reference curves for the UK, 1990. *Arch Dis Child* 1995; 73:17-24.
- [23] Oladeji PT, Stackhouse C, Gracely E, Orlin M. Comparison of the two-step and midgait methods in children. *J Am Podiatr Med Assoc* 2008; 98(4): 268-277.
- [24] Cavanagh PR, Ulbrecht JS. Clinical plantar pressure measurement in diabetes: rationale and methodology. *The Foot* 1994; 4:123–135.
- [25] Oliveira FPM, Pataky TC, Tavares JMRS. Registration of pedobarographic image data in frequency domain. *Computer Methods in Biomechanics and Biomedical Engineering* 2010; 13(6): 262-269.
- [26] Friston KJ, Ashburner JT, Kiebel SJ, Nichols TE, Penny WD, Eds. *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. London: Elsevier; 2007.

[27] Pataky TC. Assessing the significance of pedobarographic signals using random field theory. *J Biomech* 2008; 41(11):2465–2473.

[28] Giacomozzi C, Macallari V, Leardini A, Benedetti MG. Integrated pressure-force-kinematics measuring system for the characterisation of plantar foot loading during locomotion. *Med Biol Eng Comput* 2000; 38:156-63.

[29] Stebbins J, Giacomozzi C, Theologis T. Correlation between plantar pressure and Oxford Foot Model kinematics. *J Foot Ankle Res* 2008; 1:O22.

[30] Keijsers N, Stolwijk N, Pataky TC. Linear dependence of peak, mean, and pressure—time integral values in plantar pressure images. *Gait Posture* 2010; 31: 140–142.

[31] Pataky TC, Caravaggi P, Savage R, Crompton, R. Regional peak plantar pressures are highly sensitive to region boundary definitions. *J Biomech* 2008; 41:2772–2775.

[32] Lin C-J, Lai K-A, Kuan T-S, Chou Y-L. Correlating factors and clinical significance of flexible flatfoot in preschool children. *J Ped Orthop* 2001; 21:378-382.

[33] Nester CJ, Liu AM, Ward E, Howard D, Cocheba J, Derrick T, Patterson P. In vitro study of foot kinematics using a dynamic walking simulator. *J Biomech* 2007; 40(9):1927-1937.

Table 1. Barefoot peak plantar pressures for children aged 4-7 years (kPa).

	Current study	Bosch et al, 2010	Müller et al, 2011	Current study	Bosch et al, 2010	Müller et al, 2011	Current study	Bosch et al, 2010	Müller et al, 2011	Current study	Bosch et al, 2010	Müller et al, 2011	Bosch et al, 2009
	4 years	4.2 years	4 years	5 years	5.2 year	5 years	6 years	6.2 years	6 years	7 years	7.2 years	7 years	7 years
	mean	median	mean	mean	median	mean	mean	median	mean	mean	median	mean	mean
Heel	258	213	267	280	246	280	293	359	286	287	370	301	384
MedMF	22	67	87	17	64	82	28	74	80	31	70	77	83
LatMF	50			43			48			48			
MH1	134	154	159	162	184	175	147	231	188	172	247	208	256
MH2	156			145			185			198			
MH3	132			148			160			186			
MH4	109			116			116			122			
MH5	80			90			96			80			
Hallux	223	164		224	191		217	251		216	282		272




Figure 1: Region of interest (ROI) definitions: (a) Method A, and (b) Method B.




Figure 2: Average peak pressure distributions for the four age groups.




Figure 3: Linear regression results for the region of interest (ROI) data (left foot).




Figure 4: Linear regression results for the pixel-level data. (a) The raw statistical parametric map (SPM) was computed by conducting separate linear regression at all pixels; t values indicate the strength of linear regression (Eqn.1). (b) Inference images, depicting the portions of the feet which reached (or failed to reach) statistical significance; $|t| > 2$.

SUPPLEMENTARY




Figure S1: Linear regression results for the region of interest (ROI) data (right foot).