

University of Groningen

The life cycle of radio galaxies as seen by LOFAR

Brienza, Marisa

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Brienza, M. (2018). The life cycle of radio galaxies as seen by LOFAR [Groningen]: Rijksuniversiteit Groningen

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



/ rijksuniversiteit
groningen

The life cycle of radio galaxies as seen by LOFAR

Proefschrift

ter verkrijging van de graad van doctor aan de
Rijksuniversiteit Groningen
op gezag van de
rector magnificus prof. dr. E. Sterken
en volgens besluit van het College voor Promoties.

De openbare verdediging zal plaatsvinden op
vrijdag 23 maart 2018 om 14.30 uur

door

Marisa Brienza

geboren op 11 november 1987
te Varese, Italië

Promotor

Prof. dr. R. Morganti

Copromotor

Dr. G. Heald

Beoordelingscommissie

Prof. dr. P.D. Barthel

Prof. dr. G. Giovannini

Prof. dr. C.P. O'Dea

*To my families,
for giving me roots and wings.*

*Alle mie famiglie,
per avermi dato le radici e le ali.*



Netherlands Institute for Radio Astronomy



The research leading to this thesis has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Advanced Grant RADIOLIFE-320745.

LOFAR, the Low Frequency Array designed and constructed by ASTRON (Netherlands Institute for Radio Astronomy), has facilities in several countries, that are owned by various parties (each with their own funding sources), and that are collectively operated by the International LOFAR Telescope (ILT) foundation under a joint scientific policy. The Westerbork Synthesis Radio Telescope is operated by ASTRON with support from the Netherlands Foundation for Scientific Research (NWO).

The Sardinia Radio Telescope is funded by the Department of University and Research (MIUR), Italian Space Agency (ASI), and the Autonomous Region of Sardinia (RAS) and is operated as National Facility by the National Institute for Astrophysics (INAF). The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.


GMRT is run by the National Centre for Radio Astrophysics of the Tata Institute of Fundamental Research.

Cover design: Francesco Santoro, inspired by Pawel Kuczynski - New Technology

Printed by: Gildeprint

ISBN: 978-94-034-0457-8

ISBN: 978-94-034-0456-1 (electronic version)



Contents

| | | |
|----------------------|--|-----------|
| 1 | Introduction | 1 |
| 1.1 | Active galactic nuclei and their role in galaxy evolution | 3 |
| 1.2 | Jetted active galactic nuclei | 4 |
| 1.3 | The life cycle of radio galaxies | 8 |
| 1.3.1 | Remnant radio galaxies | 12 |
| 1.3.2 | Restarted radio galaxies | 16 |
| 1.4 | The Low Frequency Array | 19 |
| 1.5 | This Thesis | 21 |
| 1.5.1 | Thesis outline | 24 |
| References | | 26 |
| 2 | LOFAR discovery of a 700-kpc remnant radio galaxy at low redshift | 35 |
| 2.1 | Introduction | 37 |
| 2.2 | Data | 40 |
| 2.2.1 | LOFAR observations | 40 |
| 2.2.2 | WSRT observations | 42 |
| 2.2.3 | Archival radio data | 44 |
| 2.3 | Results | 45 |
| 2.3.1 | Morphology | 45 |
| 2.3.2 | Optical identification and galactic environment | 45 |
| 2.3.3 | Source energetics | 48 |
| 2.3.4 | Spectral properties and age | 49 |
| 2.4 | Discussion | 52 |

| | | |
|----------|--|------------|
| 2.4.1 | Evidence in favour of the remnant radio galaxy interpretation | 52 |
| 2.4.2 | Merger scenario and triggering mechanism | 53 |
| 2.4.3 | Spectral comparison with other remnant radio galaxies and implications for remnant evolution | 54 |
| 2.5 | Implications for selection of remnant radio galaxies | 56 |
| | References | 60 |
| 3 | Search and modelling of remnant radio galaxies in the LOFAR Lockman Hole field | 65 |
| 3.1 | Introduction | 67 |
| 3.2 | Lockman Hole data | 69 |
| 3.2.1 | New low-resolution catalogues | 69 |
| 3.3 | Selection of remnant radio galaxies | 74 |
| 3.3.1 | Ultra-steep spectral index selection | 74 |
| 3.3.2 | Spectral curvature selection | 75 |
| 3.3.3 | Morphology selection | 75 |
| 3.3.4 | Low radio core prominence selection | 77 |
| 3.3.5 | Results of remnant radio galaxy selection process . . | 79 |
| 3.3.6 | Implications from the selection | 85 |
| 3.4 | Simulating the population of active and remnant FRI radio galaxies | 86 |
| 3.4.1 | Dominant classes of radio source in our sample: SKADS Simulations | 87 |
| 3.4.2 | Simulation approach | 88 |
| 3.4.3 | Simulation results | 96 |
| 3.4.4 | Discussion of the Monte Carlo simulation results . . | 100 |
| 3.5 | Summary and conclusions | 103 |
| | References | 106 |
| 4 | The duty cycle of the radio galaxy B2 0258+35 | 111 |
| 4.1 | Introduction | 113 |
| 4.2 | Overview on the source B2 0258+35 | 115 |
| 4.3 | Data | 117 |
| 4.3.1 | LOFAR HBA observations and data reduction . . . | 120 |
| 4.3.2 | SRT observations and data reduction | 121 |
| 4.3.3 | VLA P-band observations and data reduction . . . | 122 |

| | | |
|--------------|--|------------|
| 4.3.4 | GMRT observations and data reduction | 123 |
| 4.4 | Results | 124 |
| 4.4.1 | Morphology | 124 |
| 4.4.2 | Spectral properties of the outer lobes | 125 |
| 4.4.3 | Energetics and age of the outer lobes | 129 |
| 4.4.4 | The central compact source | 131 |
| 4.5 | Discussion | 132 |
| 4.5.1 | Source characteristics | 134 |
| 4.5.2 | Source evolutionary scenarios | 136 |
| 4.5.3 | Comparison with other radio galaxies | 139 |
| 4.6 | Conclusions and future work | 141 |
| Appendix 4.A | Image of the full LOFAR field of view | 144 |
| References | | 145 |
| 5 | The restarted radio galaxy 3C388 | 151 |
| 5.1 | Introduction | 153 |
| 5.2 | Data | 157 |
| 5.2.1 | VLA observations at 350 MHz and data reduction . | 157 |
| 5.2.2 | VLA observations at 1400 and 4850 MHz and data reduction | 159 |
| 5.2.3 | LOFAR observations at 150 MHz and data reduction | 159 |
| 5.2.4 | GMRT observations at 612 MHz and data reduction | 160 |
| 5.3 | Results | 162 |
| 5.3.1 | Morphology | 162 |
| 5.3.2 | Spectral analysis | 162 |
| 5.3.3 | Spectral age modelling | 165 |
| 5.4 | Discussion | 169 |
| 5.4.1 | Spectral index distribution | 170 |
| 5.4.2 | Spectral ages | 172 |
| 5.4.3 | Interpretative scenarios | 173 |
| 5.5 | Conclusions | 175 |
| References | | 178 |
| 6 | A LOFAR view on the duty cycle of young radio galaxies | 183 |
| 6.1 | Introduction | 185 |
| 6.2 | The LOFAR Two-metre Sky Survey | 187 |
| 6.3 | The sample | 189 |
| 6.4 | Results | 194 |

| | | |
|--|--|------------|
| 6.4.1 | J1203+4803 | 195 |
| 6.4.2 | J1241+5458 | 199 |
| 6.4.3 | 1310+4653 | 200 |
| 6.4.4 | 1414+4554 | 202 |
| 6.5 | Discussion and conclusion | 205 |
| Appendix 6.A Radio images and spectra of the sources in the sample | | 208 |
| References | | 216 |
| 7 | Conclusions and future prospects | 219 |
| 7.1 | Conclusions chapter by chapter | 220 |
| 7.2 | General conclusions | 225 |
| 7.3 | Future prospects | 229 |
| References | | 231 |
| Samenvatting | | 233 |
| Riassunto | | 243 |
| Acknowledgements | | 253 |