

FAMILIES OF LOCALLY SEPARATED HAMILTON PATHS

János Körner

e-mail: korner@di.uniroma1.it

Angelo Monti

e-mail: [<monti@di.uniroma1.it>](mailto:monti@di.uniroma1.it)

Sapienza University of Rome

ITALY

Abstract

We improve by an exponential factor the lower bound of Körner and Muzi for the cardinality of the largest family of Hamilton paths in a complete graph of n vertices in which the union of any two paths has maximum degree 4. The improvement is through an explicit construction while the previous bound was obtained by a greedy algorithm. We solve a similar problem for permutations up to an exponential factor.

Keywords Hamilton paths, graph-difference, permutations, intersection problems

AMS Subject classification numbers 05D99, 05C35, 05C62, 94A24

1 Introduction

Greedy algorithms play a central role in combinatorial problems related to information theory. The classical example for this is the code distance problem asking for the maximum cardinality of a set of binary sequences of length n any two of which differ in at least αn coordinates where α is a fixed constant. Gilbert [4] and Varshamov [12] showed independently that for $\alpha \in [0, 1/2)$ the maximum cardinality is at least $\exp_2 n(1 - h(\alpha))$ where h is the binary entropy function $h(t) = -t \log_2 t - (1 - t) \log_2 (1 - t)$. This result, called the Gilbert–Varshamov bound, is proved by a greedy algorithm. Considerable effort went into trying to strengthen this bound, and although it was recently improved by a linear factor [6], [13], it is believed by many to be exponentially tight.

In an attempt to generalise Shannon’s zero–error capacity problem [10] from fixed–length sequences of elements of a finite alphabet to permutations and then to subgraphs of a fixed complete graph, cf. [7], [2] and [8], one of the main construction techniques is again based on greedy algorithms. It is crucial to understand the true role of greedy algorithms in our context. We will study two different combinatorial problems of information–theoretic flavour and discuss the constructions obtained by the greedy algorithm in both cases. For the general framework and especially for Shannon’s zero–error capacity problem we refer the reader to the monograph [3].

2 Hamilton paths

The first problem we are to deal with is about Hamilton paths in the fixed complete graph K_n with vertex set $[n] = \{1, 2, \dots, n\}$. We will say that two such paths are crossing if their union has maximum degree 4. Let us denote by $Q(n)$ the maximum cardinality of a family of Hamilton paths from K_n in which any pair of paths is crossing. It was proved in [8] (in Theorem 3) that

$$\frac{(n-2)!}{\lfloor n/2 \rfloor! (1 + \sqrt{2})^n} \leq Q(n) \leq \frac{n!}{\lfloor n/2 \rfloor! 2^{\lfloor n/2 \rfloor}}.$$

The lower bound was achieved by a greedy algorithm. The core of this proof was an upper bound on the number of Hamilton paths not crossing a fixed path. Here we give an explicit construction resulting in a larger lower bound. The main message of this result is that for the present problem the greedy algorithm is not tight. The improvement is in the second term of the asymptotics of $Q(n)$ and it could not be otherwise for the leading terms in the lower and upper bounds for $Q(n)$ in [8] coincide. In fact, both of the bounds in the above theorem are of the form $2^{cn}(dn)!$ where c and d are constants. The leading term is the factorial and the constant d in it is the same in both of the bounds. The difference is in the constant in the exponential factor. Our new result increases the constant in the exponential factor in the lower bound in Theorem 3 in [8].

More precisely, we have

Theorem 1

$$\frac{(\lfloor n/2 \rfloor - 1)!}{2^{\lfloor n/4 \rfloor}} \leq Q(n).$$

Before proving our result we observe that

$$\frac{(n-2)!}{\lfloor n/2 \rfloor!(1+\sqrt{2})^n} < \left(\frac{2}{1+\sqrt{2}}\right)^n \cdot \lceil n/2 \rceil! \leq \frac{(\lfloor n/2 \rfloor - 1)!}{2^{\lfloor n/4 \rfloor}}$$

where the first inequality follows since

$$\frac{(n-2)!}{\lfloor n/2 \rfloor!} \leq \frac{n!}{\lfloor n/2 \rfloor!} = \binom{n}{\lfloor n/2 \rfloor} (\lceil n/2 \rceil)! < 2^n (\lceil n/2 \rceil)!$$

while the last inequality is obvious as $\frac{2}{1+\sqrt{2}} < 2^{-1/4}$. Thus the new lower bound of this theorem improves on the one in [8] quoted above.

Proof.

We consider the complete bipartite graph $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$. For the ease of notation let us suppose that this graph is defined by the bipartition of $[n]$ into the sets A and B where $A = \lfloor n/2 \rfloor$ is the set of the first $\lfloor n/2 \rfloor$ natural numbers while $B = [n] - A$. Let us fix a permutation of the elements of B . Then any permutation of A defines a Hamilton path in $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$ where the first element of the path is the first element of the permutation of B . The path alternates the elements of A and B in such a way that the elements of B are in the odd positions along the path and in the order defined by the permutation of B we have fixed. Likewise, the relative order of the elements of A following the path is the one given by the permutation of A . Every Hamilton path in this family can thus be identified with some permutation of the set A . We represent each permutation as a linear order of A .

Notice now that if there is an element $a \in A$ such that in two different linear orders its position differs by at least 2 while none of these positions is the last one, then the vertex a has disjoint pairs of adjacent vertices in the two Hamilton paths. In other words, the vertex a has degree 4 in the union of these paths. In this case we will say that the corresponding pair of permutations is *two-different*. We will now construct a sufficiently large family of pairwise two-different permutations. To this end, we first observe that if two permutations are two-different, then in their respective inverses there is some natural number whose images differ by at least two. Körner, Simonyi and Sinaimeri have shown (cf. Theorem 1 in [9]) that for any natural number m the set $[m]$ has a family of exactly $\frac{m!}{2^{\lfloor m/2 \rfloor}}$ permutations such that for any two of them there is at least one element of $[m]$ that has images differing by at least two. Hence the inverses of these permutations, seen as linear orders, form a family satisfying our original condition of having a same element a in positions differing by at least 2, except for the fact that one of these two positions might be the last position in one of the permutations. To exclude this event, we choose a subfamily of our permutations in which the last element in all the permutations is the same. Choosing $m = |A|$ we complete the proof.

□

The construction in the last proof seems sufficiently natural for us to believe that it might be asymptotically optimal at least for Hamilton paths in a balanced complete bipartite graph. In other words, let $B(n)$ denote the largest cardinality of a family of Hamilton paths in $K_{\lfloor n/2 \rfloor, \lfloor n/2 \rfloor}$ any two members of which are crossing. If the answer is positive, one might go even further and ask whether the lower bound in Theorem 1 is asymptotically tight.

3 Intersecting vs. separated families

In [8] where the previous problem was introduced, it was presented within a fairly general framework for extremal combinatorics with an information-theoretic flavour. This framework is in terms of two (mostly but not necessarily disjoint) families of graphs on the same vertex set $[n] = \{1, 2, \dots, n\}$. If we denote these families by \mathcal{F} and \mathcal{D} , respectively, then $M(\mathcal{F}, \mathcal{D}, n)$ stands for the largest cardinality of a subfamily $\mathcal{G} \subseteq \mathcal{F}$ with the property that the union of any two of its different members belongs to \mathcal{D} . Here the union of two graphs on the same vertex set is the graph whose edge set is the union of those of the two graphs. This framework is the fruit of an attempt to describe a consistent part of extremal combinatorics where information theoretic methods seem to be relevant. In [8] it was emphasised that when the families \mathcal{F} and \mathcal{D} are disjoint, the question about $M(\mathcal{F}, \mathcal{D}, n)$ is fairly information-theoretic in nature and there never is a natural candidate for an optimal construction; much as it happens for the graph capacity problem of Shannon's. As a matter of fact, in each of these cases one is interested in the largest cardinality of a family of mathematical objects (graphs or strings) any pair of which are different in some specific way. Intuitively, the objects in these families are not only different, but also distinguishable in some well-defined sense inherent to the problem. In other words, the optimal constructions are, in a vague and broad sense, similar to the channel codes in information theory.

If, however, $\mathcal{F} = \mathcal{D}$, the solution of our problems seems to be of a completely different nature, and, in particular, any analogy with codes disappears. Rather, the optimal solutions often are so-called kernel structures, and even when this is not the case, kernel structures give good constructions for the problem. Kernel structures are families of objects having in common a fixed projection, called the kernel. This means that in a vague and broad sense, these objects are similar. We are trying to build up a dichotomy distinguishing those problems for which kernel structures can be defined (and often are even optimal) and those where the optimal structures are like "codes" in information theory. It would be highly interesting to get a better understanding of this dichotomy.

At the surface, the dichotomy for the problems of determining $M(\mathcal{F}, \mathcal{D}, n)$ is based on whether or not $\mathcal{F} = \mathcal{D}$. However, this view is too simplistic. As a matter of fact, our next problem will illustrate the lack of an easy way to distinguish between the two cases. More precisely, we will give an example where the families \mathcal{F} and \mathcal{D} are disjoint, and the

solution of the problem still has a kernel structure. To explain our example, let \mathcal{F}_{cy} be the family of Hamilton cycles in the complete graph K_n . However, let \mathcal{D}_3 be the family of graphs having at least one vertex of degree 3. The relation of having a vertex of degree 3 in the union of two graphs is clearly irreflexive when restricted to cycles. In fact, the problem of determining $M(\mathcal{F}_{\text{cy}}, \mathcal{D}_3, n)$ is, on the surface, no different from the one which is the subject of Theorem 1. However, this impression is wrong and the present problem is completely different from the previous one. To understand why, it suffices to realise that the union of two Hamilton cycles in K_n has a vertex of degree 3 if and only if they share a common edge. This makes it possible to invoke kernel structures. Moreover, not surprisingly, a quasi-optimal solution for this problem has a kernel structure. In fact, we have

Proposition 1

$$M(\mathcal{F}_{\text{cy}}, \mathcal{D}_3, n) \geq (n - 2)!$$

and this bound is tight for odd n . For even n we have

$$M(\mathcal{F}_{\text{cy}}, \mathcal{D}_3, n) \leq (n - 1)[(n - 3)!]$$

Proof.

To prove the lower bound, consider the family of all Hamilton cycles in K_n containing a fixed edge. To establish the upper bound, let first n be odd. By a classical construction of Walecki (1890), cited in [1], for every odd n , the family of the Hamilton cycles of K_n can be partitioned into $(n - 2)!$ classes where the cycles belonging to the same class are disjoint. (Each class is a partition of the edges of K_n into $\frac{n-1}{2}$ Hamilton cycles). This gives our upper bound for odd n .

For even n we can only have families of $\frac{n-2}{2}$ pairwise disjoint Hamilton cycles. The total number of Hamilton cycles is $\frac{(n-1)!}{2}$ and each of these is contained, by symmetry, in the same number of families of pairwise edge-disjoint cycles. Thus, by double-counting, the maximum number of pairwise edge-intersecting Hamilton cycles is at most the $\frac{n-2}{2}$ 'th part of the total which gives the weaker upper bound for this case.

□

4 Permutations

A permutation of $[n]$ is nothing but a (consecutively oriented) Hamilton path in the symmetrically complete directed graph W_n . (We denote by W_n the directed graph in which every pair of distinct vertices is connected by two edges going in opposite direction.) In the previous part of the paper we were dealing with problems about Hamilton paths in (the undirected complete graph) K_n such that for any pair of them, for at least one vertex in K_n the union of its 1-neighbourhoods in the two paths satisfied some disjointness condition. We now turn to an analogous problem about directed paths.

In a directed graph, the (open) 2-out-neighbourhood of a vertex $a \in [n]$ consists of the vertices reachable from a by a (consecutively oriented) path of at most two edges. We will say that two directed Hamilton paths in W_n are *two-separated* if for at least one vertex of W_n the union of its 2-out-neighbourhoods in the two paths contains exactly 4 vertices. Formulating this condition in terms of permutations means that for some $a \in [n]$ its two immediate successors in the two linear orders defining the two respective permutations are axy and avw where x, y, v, w are four different elements of $[n]$. This relation is seemingly very similar to the one underlying Theorem 1. As a matter of fact, in the permutation language, to two crossing Hamilton paths there correspond two permutations such that for some $a \in [n]$ its neighbours are completely different, meaning that we have xay and vaw in the two permutations and x, y, v, w are all different. If all pairs of paths in a Hamilton path family are crossing then for any one of the paths only one of the corresponding two permutations can be in the family, and it can be either one of the two that correspond to the same path. Somewhat surprisingly, the cardinality of the largest family of pairwise two-separated Hamilton paths is of a different order of magnitude from the one in Theorem 1. In fact, we have the following almost tight result.

Theorem 2 *Let $R(n)$ stand for the maximum cardinality of a family of pairwise two-separated Hamilton paths in W_n . Then*

$$\frac{n!}{n^{13}5^{n-10}} \leq R(n) \leq \frac{n!}{2^{\lfloor n/2 \rfloor}}.$$

Proof.

In this proof we will represent our directed Hamilton paths as permutations. We will say that two permutations are two-separated if they correspond to two-separated Hamilton paths. We associate with any permutation a sequence of unordered couples of elements of $[n]$. The permutation $\pi = \pi_1 \dots \pi_n$ is associated with the sequence of unordered and disjoint couples $\{\pi_1, \pi_2\} \dots \{\pi_{2i-1}, \pi_{2i}\} \dots \{\pi_{2\lfloor n/2 \rfloor - 1}, \pi_{2\lfloor n/2 \rfloor}\}$. We will call this sequence of couples the couple order of π . We then partition the set of all permutations of $[n]$ according to their couple order. In each class the permutations it contains have the same couple order and all the permutations with the same couple order belong to this class. It is sufficient to show that no two permutations in the same class are two-separated. Considering that the number of classes is $\frac{n!}{2^{\lfloor n/2 \rfloor}}$, we will thus obtain our upper bound.

In fact, suppose to the contrary that the permutations π' and π'' having the same couple order are two-separated. Let xy and vw be the two immediate successors of a in π' and π'' , respectively. We will distinguish two cases. In the first case a and x appear in the same couple $\{a, x\}$ of the couple order, while in the second case a is in a different couple from x and y , which then have to appear together in the succeeding couple. In the first case, if a and x appear in the same order (a, x) in both permutations, we are done. Otherwise, if their order is different, then y must appear in one of the two positions following a in π'' , contradicting our hypothesis that π' and π'' are two-separated. It

remains to see what happens if in the *couple order* of π' the number a is in the couple preceding $\{x, y\}$. This means that in the couple order a is in the same couple with some $z \in [n]$ different from both x and y . Since our two permutations have the same couple order, we conclude that at least one of x and y is in one of the two positions immediately following a in π'' , but then π' and π'' are not two-separated.

Let us now turn to lower bounding $R(n)$. To this end we will use a greedy algorithm to exhibit a large enough family of permutations with the required property. At each step in the algorithm we choose an arbitrary permutation and eliminate from the choice space all those not two-separated from the chosen one. For brevity, we will say that the latter are *incompatible* with the chosen permutation. This procedure goes on until the choice space becomes empty.

To analyse this algorithm, we need an upper bound on the number of those permutations which are not two-separated from a fixed permutation ι of $[n]$. Without loss of generality we assume that ι is the identity permutation mapping each number in itself. Let $\pi = \pi_1\pi_2 \dots \pi_n$ be a permutation incompatible with ι . We have that

$$\text{for each } j, 1 \leq j \leq n-2, \text{ if } \pi_j \notin \{n, n-1\} \text{ then } \{\pi_{j+1}, \pi_{j+2}\} \cap \{\pi_j + 1, \pi_j + 2\} \neq \emptyset \quad (1)$$

otherwise the two immediate successors of π_j in the two permutations ι and π are four different elements of $[n]$, contradicting that ι and π are incompatible.

At this point we have to introduce some terminology. If $\pi_{j+1} \in \{\pi_j + 1, \pi_j + 2\}$ or respectively $\pi_{j+2} \in \{\pi_j + 1, \pi_j + 2\}$ then we say that π_{j+1} or respectively π_{j+2} is a *close follower* of π_j . In this terminology property (1) means that at least one of π_{j+1} and π_{j+2} must be a close follower of π_j .

In the following we call *run of big jumps* in π any maximal (not extendable) sequence $\pi_j\pi_{j+1} \dots \pi_{j+t}$, $1 \leq j \leq n-t$ (of length at least two) such that $\pi_{j+1} - \pi_j \notin \{-2, -1, 1, 2, 3\}$ and $\pi_{k+1} - \pi_k \notin \{-1, 1, 2\}$, $j < k < j+t$. We will refer to the position j as the *head of the run* and to the positions k of the run with $k \geq j+2$ as *tail positions of the run*. We call *free positions* of π the first position of the entire permutation, the second position in any run and the two positions immediately following those of either n or $n-1$. All the other positions will be called *constrained*. This terminology is clarified by the following claim: in a constrained position we must have one of at most five possible values. To justify this claim we first notice that for any constrained position j not belonging to any run we have

$$\pi_j \in \{\pi_{j-1} - 2, \pi_{j-1} - 1, \pi_{j-1} + 1, \pi_{j-1} + 2, \pi_{j-1} + 3\} \quad (2)$$

The same is true if j is the head of a run since in this case either $\pi_j - \pi_{j-1} > 3$ or $\pi_j - \pi_{j-1} < -2$ would contradict the maximality of the run.

Consider now a constrained position j in the tail of a run. We claim that the number π_j in this position must be a close follower of at least one of π_{j-1} and π_{j-2} . By definition, position $j-2$ also belongs to the same run. Thus either $\pi_{j-1} - \pi_{j-2} \geq 3$ or $\pi_{j-1} - \pi_{j-2} \leq -2$. This implies that $\pi_{j-1} \notin \{\pi_{j-2} + 1, \pi_{j-2} + 2\}$ and from property (1) we

obtain $\pi_j \in \{\pi_{j-2} + 1, \pi_{j-2} + 2\}$. We have just proved our claim that for any constrained position j of π the value π_j satisfies (3).

Let us now consider the permutations incompatible with ι having exactly r runs of big jumps. In these permutations there are at most $r + 5$ free positions and at most $\binom{n-1}{r+2}$ possibilities for these free positions (remember that, by definition, the first position is always a free position and that the two free positions depending on n are adjacent to it and the same is true for the two free positions depending on $n - 1$). Further, let us recall that in a constrained position we can have one of only 5 possible values. Thus the number of these incompatible permutations is upper bounded by

$$\binom{n-1}{r+2} n^{r+5} 5^{n-(r+5)}. \quad (3)$$

We will prove that in any incompatible permutation there can be at most three runs of big jumps. This in turn implies that any incompatible permutation can have at most eight free positions and from bound (3) we have that the number of permutation incompatible with permutation ι is bounded above by

$$4 \binom{n-1}{5} n^8 5^{n-8}.$$

This means that the greedy algorithm will eliminate at most $n^{13} 5^{n-10}$ permutations at each step, yielding a family with the desired property and containing at least

$$\frac{n!}{n^{13} 5^{n-10}}$$

permutations, as claimed for the lower bound.

To conclude the proof it remains to show that in a permutation π incompatible with ι there are at most three runs of big jumps. To this aim we now prove that any run of big jumps of π that contains neither n nor $n - 1$ is a suffix of π . The proof is by contradiction. Let us assume that in π there is a run that contains neither n nor $n - 1$ and yet it is not a suffix. Let t be the last position of such a run.

We distinguish two cases. In the first case, we suppose

$$\pi_{t-1} > \pi_t.$$

Thus π_t is not a close follower of π_{t-1} whence, by property (1), we must have $\pi_{t+1} \in \{\pi_{t-1} + 1, \pi_{t-1} + 2\}$. Since π_t is in the run of big jumps we are considering, the last inequality can be satisfied only if

$$\pi_t \leq \pi_{t-1} - 2.$$

(Note that, by our assumption that the run we are considering is not a suffix of π we have that π_t is not the last element in π). But then

$$\pi_{t+1} - \pi_t \geq 3,$$

contradicting the assumption that π_t is in the last position of the run of big jumps.

In the second case we suppose

$$\pi_{t-1} < \pi_t.$$

This implies

$$\pi_t \geq \pi_{t-1} + 3. \tag{4}$$

Thus π_t is not a close follower of π_{t-1} so that once again, by property (1), we must have $\pi_{t+1} \in \{\pi_{t-1} + 1, \pi_{t-1} + 2\}$. In particular, we see that $\pi_{t+1} < \pi_t$. However, since π_{t+1} is not part of the run of big jumps, the only possibility left is

$$\pi_{t+1} = \pi_t - 1. \tag{5}$$

Since π_t is not a close follower of π_{t-1} this role must be taken by π_{t+1} . Hence from (5) and from (4) it follows that

$$\pi_t = \pi_{t-1} + 3 \tag{6}$$

Thus, by (5) and (6), we have

$$\pi_{t+1} = \pi_{t-1} + 2. \tag{7}$$

Let us see what we can say in the light of this about π_{t-2} . Note that $\pi_t - \pi_{t-1} = 3$ which implies that position $t - 1$ is not the head of the run of big jumps. This in turn implies that position $t - 2$ is in the run and $\pi_{t-1} - \pi_{t-2} \notin \{-1, 1, 2\}$. Hence we must have $\pi_{t-2} \leq \pi_{t-1} - 3$ or $\pi_{t-2} \geq \pi_{t-1} + 2$. Moreover, property (1) implies

$$\{\pi_{t-2} + 1, \pi_{t-2} + 2\} \cap \{\pi_{t-1}, \pi_t\} \neq \emptyset,$$

whence the only choice left is $\pi_{t-2} = \pi_{t-1} + 2$ (remember that $\pi_t = \pi_{t-1} + 3$), and this is impossible by equality (7). □

5 Problem classes

We have already mentioned the need for a meta-conjecture to describe for which problems within our framework does the optimal solution have a kernel structure. It seems to us that this can happen only in case of one-family problems and it will never be the case if the family \mathcal{D} in a two-family problem is not monotone. (A class \mathcal{F} of graphs is monotone if $G \in \mathcal{F}$ and $G \subseteq H$ imply $H \in \mathcal{F}$. Note, in particular, that the family \mathcal{D}_3 of graphs having a vertex of degree 3 is not monotone.) Our problems belong to a class of combinatorial problems introduced in [11] where they are referred to as intersection problems. In the eyes of the authors of [11] an intersection problem is in terms of the maximum cardinality of a family of objects such that the pairwise intersections of these objects have some prescribed property. (Even if, in this paper, we are considering pairwise unions, we might have considered, equivalently, the intersections of the complementary objects.) We feel that it is more appropriate to reserve the term intersection problem to the case $\mathcal{F} = \mathcal{D}$

when in fact we end up with an intersection problem in disguise, in the sense of [11]. In our opinion, an intersection problem is a one-family problem. In such problems one asks for the largest cardinality of a family of objects, typically subgraphs of K_n with a given property such that the intersection of any two members of the family is still in the family. One often considers as intersection problems only those about the maximum cardinality of a family of graphs such that the intersection of any two of them is non-empty.

A more delicate question is to understand when exactly can the optimal solution of a problem of our kind be obtained through a greedy algorithm. Our two main results can hopefully contribute to a better understanding of this issue. At a first glance one is tempted to believe that for an information-theoretic (genuinely two-family) problem the greedy algorithm often gives close-to-optimal constructions. Our question is not precise enough and we do not expect a precise answer at this stage.

6 Acknowledgement

We are grateful to Miki Simonovits for stimulating discussions about the true nature of intersection problems.

References

- [1] J. Akiyama, M. Kobayashi, G. Nakamura, Symmetric Hamilton cycle decompositions of the complete graph. *J. Combinatorial Designs*, **12**(2004), no. 1, pp. 39–45,
- [2] G. Brightwell, G. Cohen, E. Fachini, M. Fairthorne, J. Körner, G. Simonyi, and Á. Tóth, *Permutation capacities of families of oriented infinite paths*, *SIAM J. Discrete Math.* **24** (2010), 441–456,
- [3] I. Csiszár, J. Körner, *Information theory: coding theorems for discrete memoryless systems*, 2nd edition, Cambridge University Press, 2011,
- [4] E. N. Gilbert, A comparison of signalling alphabets, *Bell Syst. Tech. J.*, **31**(1952), pp. 504–522,
- [5] N. Hartfield, G. Ringel, *Pearls of graph theory*, Academic Press, New York, 1990,
- [6] Tao Jiang, A. Vardy: Asymptotic Improvement of the Gilbert-Varshamov Bound on the Size of Binary Codes, *IEEE Transactions on Information Theory* **50**(2004) no. 8, pp. 1655–1664
- [7] J. Körner, C. Malvenuto, Pairwise colliding permutations and the capacity of infinite graphs, *SIAM J. Discrete Mathematics*, **20** (2006), 203–212.
- [8] J. Körner, I. Muzi, Degree–doubling graph families, *SIAM J. on Discrete Mathematics*, **27** (2013), no. 3, pp. 1575–1583,
- [9] J. Körner, G. Simonyi, B. Sinaireri, On types of growth for graph–different permutations, *J. Comb. Theory Ser. A*, **116** (2009), 713–723.
- [10] C. E. Shannon, The zero–error capacity of a noisy channel, *IRE Trans. Inform. Theory*, **2** (1956), 8–19.
- [11] M. Simonovits and V. T. Sós, Intersection theorems on structures, *Annals of Discrete Mathematics*, 6(1980), pp. 301–313,
- [12] R. R. Varshamov, Estimate of the number of signals in error correcting codes, *Dokl. Acad. Nauk SSSR* **117**pp. 739–741.
- [13] Van Vu, Lei Wu, Improving the Gilbert–Varshamov bound for q -ary codes, *IEEE Transactions on Information Theory* **51**(2005) no. 9, pp. 3200–3208