

12

13 **ABSTRACT**

14 Different *in vitro* assays characterise most of the essential oils and phenolic compounds
15 as antioxidants. These molecules can be found in a variety of aromatic plants and have
16 been related to their bioactive properties. For the first time, a comparative study
17 between the antioxidant properties of essential oils and phenolic extracts from *Cistus*
18 *ladanifer* leaves, *Citrus latifolia* fruit peels, *Cupressus lusitanica* foliage and *Eucalyptus*
19 *gunnii* leaves was performed. Overall, the antioxidant properties of phenolic extracts
20 (unless scavenging activity of *Citrus latifolia*) were excellent and better than those
21 obtained from the essential oils extracts, and even for the standards BHA (2-*tert*-butyl-
22 4-methoxyphenol) and α -tocopherol. The better EC₅₀ values for all the assays
23 (scavenging activity, reducing power and lipid peroxidation inhibition) were obtained in
24 the *Eucalyptus gunnii* phenolic extract (less than 0.1 mg mL⁻¹). Among the essential oils
25 extracts, the best contribution was given by *Cistus ladanifer*.

26

27 **Keywords:** Aromatic plants; Essential oils; Phenolics; Antioxidant activity.

28

29 **1. Introduction**

30 Essential oils are volatile, natural compounds with a strong odour and formed by
31 aromatic plants as secondary metabolites. Since the middle ages, essential oils have
32 been widely used for bactericidal, virucidal, fungicidal, antiparasitical, insecticidal,
33 medicinal and cosmetic applications, especially nowadays in pharmaceutical, sanitary,
34 cosmetic, agricultural and food industries. Particularly, they are used in embalmmnt,
35 preservation of foods and as antimicrobial, analgesic, sedative, anti-inflammatory,
36 spasmolytic and locally anesthetic remedies. Because of the mode of extraction, mostly
37 by distillation (steam or hydro-distillation) from aromatic plants, they contain a variety
38 of volatile molecules such as terpenes and terpenoids, phenol-derived aromatic
39 components and aliphatic components. *In vitro* physicochemical assays characterise
40 most of them as antioxidants ([Bakkali et al., 2008](#)).

41 Phenolic compounds are aromatic hydroxylated compounds commonly found in
42 vegetables, fruits and many food sources that form a significant portion of our diet, and
43 some of which are among the most potent and therapeutically useful bioactive
44 substances. Natural phenolic compounds accumulate as end-products from the
45 shikimate and acetate pathways and can range from relatively simple molecules
46 (phenolic acids, phenylpropanoids, flavonoids) to highly polymerised compounds
47 (lignins, melanins, tannins), with flavonoids representing the most common and widely
48 distributed sub-group ([Bravo, 1998](#)). In our diet, they might provide health benefits
49 associated with reduced risk of chronic diseases that may be due to their ability to
50 reduce agents by donating hydrogen and quenching singlet oxygen ([Nijveldt et al.,](#)
51 [2001](#)). Antioxidant properties of phenolic compounds also play a vital role in the

52 stability of food products, as well as in the antioxidative defence mechanisms of
53 biological systems (Macheix and Fleuriert, 1998).

54 In Portugal, *Cistus ladanifer* (systematic family Cistaceae) is widely distributed, being
55 one of the most abundant species in the southern part of the country, occurring in large
56 areas as pure dense stands (Amaral, 1971; Teixeira et al., 2007). The “Cistus” products
57 are particularly appreciated for their balsamic odour, as well as for their fixative
58 properties (Moyler and Clery, 1997). *Cistus* species are used as an antidiarrheics, as
59 general remedies in folk medicine for treatment of various skin diseases, and as anti-
60 inflammatory agents (Attaguile et al., 2000). Several research works have been reported
61 in the literature on *C. ladanifer* L. volatile compounds, due to the great importance of
62 this raw material for the fragrance industry (Teixeira et al., 2007). Phytochemical
63 studies on different *Cistus* species have also revealed the presence of several flavonoid
64 compounds that are considered to be chain-breaking antioxidants (Danne et al., 1994).

65 *Cupressus lusitanica*, commonly known as cedar of Goa, Mexican cypress and
66 Portuguese cypress, belongs to the systematic family Cupressaceae. The leaves of this
67 plant are used in indigenous practice to treat catarrh and headache. The essential oil of
68 the leaves is used against rheumatism, whooping cough, and styptic problems (Kuiate et
69 al., 2006). The chemical analysis of three samples of this oil from Portugal showed that
70 it contains monoterpenes, sesquiterpenes and diterpenes, with abietadiene as major
71 component (Adams et al., 2001). Nevertheless, its phenolic composition and antioxidant
72 activity was not reported yet.

73 *Eucalyptus* species are fast growing trees exploited mainly for paper pulp but also as a
74 source for various essential oils. For the production of phytopharmaceuticals, essential
75 oils rich in 1,8-cineole (called also “eucalyptol”), are of special importance. These

76 products are applied for relief of head colds, rheumatism, muscular pain, and as
77 expectorant in cases of bronchitis (added to cough syrups) (Lassak and McCarthy,
78 1983). Only the antioxidant activity and phenolic contents of *Eucalyptus globulus* bark
79 aqueous extracts were reported (Vázquez et al., 2009).

80 Lime is the second most important citrus fruit, in both fresh consumption and industrial
81 uses; it is the fifth largest crop in harvested area worldwide. The Persian lime (*Citrus*
82 *latifolia*) is one of the main varieties in the production of this fruit. Total phenolics
83 content and antioxidant activity of this lime species peels were reported by Urbando-
84 Rivera et al. (2005), while nothing was described in its essential oils composition.

85 In this work, the antioxidant properties of four different aromatic plants (*Cistus*
86 *ladanifer* leaves, *Citrus latifolia* fruit peels, *Cupressus lusitanica* foliage and *Eucalyptus*
87 *gunnii* leaves) were evaluated, and compared, for the first time, considering two
88 fractions: volatile fraction (essential oils) and phenolic fraction. To access the different
89 contributions of both extracts, their DPPH (2,2-diphenyl-1-picrylhydrazyl) radical
90 scavenging activity, reducing power, and inhibition of lipid peroxidation by decreasing
91 the β -carotene bleaching were evaluated.

92

93 **2. Materials and Methods**

94

95 *2.1. Standards and reagents*

96 All the solvents were of analytical grade purity; methanol was supplied by Lab-Scan
97 (Lisbon, Portugal). The standards used in the antioxidant activity assays: BHA (2-*tert*-
98 butyl-4-methoxyphenol), TBHQ (tert-butylhydroquinone), L-ascorbic acid, α -
99 tocopherol, gallic acid and (+)-catechin were purchased from Sigma (St. Louis, MO,

100 USA). 2,2-Diphenyl-1-picrylhydrazyl (DPPH) was obtained from Alfa Aesar (Ward
101 Hill, MA, USA). The standard butylated hydroxytoluene (BHT) was purchased from
102 Merck (Darmstadt, Germany). All other chemicals were obtained from Sigma Chemical
103 Co. (St. Louis, MO, USA). Water was treated in a Milli-Q water purification system
104 (TGI Pure Water Systems, USA).

105

106 2.2. *Plant material*

107 *Cistus ladanifer* fresh leaves from flowering steams were collected randomly, from wild
108 plants growing in the Natural Park of Montesinho (Northeastern Portugal) in July 2008.

109 *Cupressus lusitanica* foliage and *Eucalyptus gunnii* matured alternate leaves were
110 randomly gathered from the crown of trees cultivated in the experimental farm of the
111 school of agriculture (Escola Superior Agrária, ESA), in Bragança (Northeastern
112 Portugal) in July 2008. *Citrus latifolia* fruits were obtained from a local supermarket in
113 December 2008, and the bioactive compounds were extracted from the peels.

114

115 2.3. *Essential oils extraction*

116 The essential oil samples were isolated from the fresh material (~100 g leaves/peels plus
117 350 mL of distilled ultra pure water) by hydro-distillation for 3 h, using a Clevenger-
118 type apparatus. The extracts were dried with anhydrous sulphate and concentrated under
119 reduced pressure by rotatory evaporator, until water evaporation (Fakhari et al., 2005).
120 The extraction yield was calculated in mL of oil per 100 g of fresh material. The
121 collected oil was weighed, dissolved in methanol at a concentration of 500 mg mL⁻¹,
122 and stored in sealed vials at -20°C for further use.

123

124 *2.4. Phenolics extraction*

125 Lyophilized (Ly-8-FM-ULE, Snijders, HOLLAND) powdered samples (typically 3 g)
126 were extracted by stirring with 50 mL of methanol at 25°C at 150 rpm for 12h and
127 filtered through Whatman n° 4 paper. The residue was then extracted with one
128 additional 50 mL portion of the methanol. The extracts were evaporated (rotary
129 evaporator Büchi R-210) to dryness and redissolved in methanol at a concentration of 5
130 mg mL⁻¹, and stored at 4°C for further use.

131 For phenolics estimation, the extract solution (1 mL) was mixed with Folin and
132 Ciocalteu's phenol reagent (1 mL). After 3 min, saturated sodium carbonate solution (1
133 mL) was added to the mixture and adjusted to 10 mL with distilled water (Singleton and
134 Rossi, 1965). The reaction was kept in the dark for 90 min, after which the absorbance
135 was read at $\lambda=725$ nm (Analytikijena 200-2004 spectrophotometer). Gallic acid was
136 used to calculate the standard curve (0.01-0.4 mM; $Y=2.8557X-0.0021$; $R^2=0.9999$) and
137 the results were expressed as mg of gallic acid equivalents (GAE) per g of extract.

138

139 *2.5. Antioxidant activity*

140 *2.5.1. DPPH radical-scavenging activity*

141 Various concentrations of the extracts (0.3 mL) were mixed with 2.7 mL of methanolic
142 solution containing DPPH radicals (6×10^{-5} mol L⁻¹). The mixture was shaken vigorously
143 and left to stand for 60 min in the dark (until stable absorption values were obtained).
144 The reduction of the DPPH radical was determined by measuring the absorption at 517
145 nm (Hatano et al., 1988). The radical scavenging activity (RSA) was calculated as a
146 percentage of DPPH discoloration using the equation: $\% \text{ RSA} = [(A_{\text{DPPH}} - A_{\text{S}}) / A_{\text{DPPH}}] \times$
147 100, where A_{S} is the absorbance of the solution when the sample extract has been added

148 at a particular level, and A_{DPPH} is the absorbance of the DPPH solution. The extract
149 concentration providing 50% of radicals scavenging activity (EC_{50}) was calculated by
150 interpolation from the graph of RSA percentage against extract concentration. BHA and
151 α -tocopherol were used as standards.

152

153 *2.5.2. Reducing power*

154 Various concentrations of the extracts (1.0 mL) were mixed with 1.0 mL of 200 mmol
155 L^{-1} sodium phosphate buffer (pH 6.6) and 1.0 mL of 1% potassium ferricyanide. The
156 mixture was incubated at 50°C for 20 min. After 1.0 mL of 10% trichloroacetic acid
157 (w/v) were added, the mixture was centrifuged at 1000 rpm for 8 min (Centorion
158 K24OR- 2003 refrigerated centrifuge). The upper layer (1 mL) was mixed with 1 mL of
159 deionised water and 0.2 mL of 0.1% of ferric chloride, and the absorbance was
160 measured spectrophotometrically at 700 nm: higher absorbance indicates higher
161 reducing power (Oyaizu, 1986). The extract concentration providing 0.5 of absorbance
162 (EC_{50}) was calculated by interpolation from the graph of absorbance at 700 nm against
163 extract concentration. BHA and α -tocopherol were used as standards.

164

165 *2.5.3. Inhibition of β -carotene bleaching*

166 The antioxidant activity of the extracts was evaluated by the β -carotene linoleate model
167 system. A solution of β -carotene was prepared by dissolving β -carotene (2 mg) in
168 chloroform (10 mL). Two millilitres of this solution were pipetted into a round-bottom
169 flask. After the chloroform was removed at 40°C under vacuum, linoleic acid (40 mg),
170 Tween 80 emulsifier (400 mg), and distilled water (100 mL) were added to the flask
171 with vigorous shaking. Aliquots (4.8 mL) of this emulsion were transferred into
172 different test tubes containing different concentrations of the extracts (0.2 mL). The

173 tubes were shaken and incubated at 50°C in a water bath. As soon as the emulsion was
174 added to each tube, the zero time absorbance was measured at 470 nm using a
175 spectrophotometer. Absorbance readings were then recorded at 20-min intervals until
176 the control sample had changed colour. A blank, devoid of β -carotene, was prepared for
177 background subtraction (Shon et al., 2003). Lipid peroxidation (LPO) inhibition was
178 calculated using the following equation: LPO inhibition = (β -carotene content after 2h
179 of assay/initial β -carotene content) \times 100. The extract concentration providing 50%
180 antioxidant activity (EC_{50}) was calculated by interpolation from the graph of antioxidant
181 activity percentage against extract concentration. TBHQ was used as standard.

182

183 2.6. Statistical analysis

184 For each one of the plants three samples were analysed and also all the assays were
185 carried out in triplicate. The results are expressed as mean values and standard deviation
186 (SD). The statistical differences represented by letters (**Table 1**) were obtained through
187 one-way analysis of variance (ANOVA) followed by Tukey's honestly test, coupled
188 with Welch's statistical analysis using SPSS software version 16.0.

189

190 3. Results and Discussion

191 Radical scavenging effects of phenolic and essential oils extracts of the aromatic plants
192 *Cistus ladanifer*, *Citrus latifolia*, *Cupressus lusitanica* and *Eucalyptus gunnii* were
193 examined and compared (**Figure 1**). Results are expressed as a percentage of the ratio
194 of the decrease in the absorbance at 517 nm to the absorbance of DPPH solution in the
195 absence of sample at 517 nm. The extracts scavenging effects on DPPH radicals
196 increase with the concentration and the results are very good for phenolic extracts,

197 specially for *Eucalyptus gunnii* sample ($95.96 \pm 0.14\%$ at 0.31 mg mL^{-1}), being much
198 higher than the scavenging effects of the standards BHA (96% at 3.6 mg mL^{-1}) and α -
199 tocopherol (95% at 8.6 mg mL^{-1}) (**Figure 1**). Nevertheless, this effect drastically
200 decreases in the corresponding essential oils extracts and, curiously, the lowest values
201 were obtained for the *Eucalyptus gunnii* sample ($68.18 \pm 1.86\%$ at $500.00 \text{ mg mL}^{-1}$). In
202 the essential oils extracts, the best contribution to the antioxidant activity was achieved
203 by *Cistus ladanifer* ($86.88 \pm 0.76\%$ at $153.60 \text{ mg mL}^{-1}$).

204 Reducing power of the aromatic plants was examined as a function of their
205 concentration in phenolics or essential oils (**Figure 2**). Reducing power of the samples
206 increased with the increase of concentration. A high value of absorbance at 700 nm is
207 related to a high reducing power. For phenolic extracts the absorbance values at 700 nm
208 were higher than the ones observed for essential oils extracts. The reducing power of
209 *Eucalyptus gunnii* phenolic extract was the highest (1.77 ± 0.07 at 0.31 mg mL^{-1}) while
210 for the lime extract was the lowest (0.02 ± 0.00 at 0.31 mg mL^{-1}). For essential oils
211 extracts the best reducing power was obtained for *Cistus ladanifer* sample (0.60 ± 0.00
212 at 4.80 mg mL^{-1}) and the worst was obtained for *Eucalyptus gunnii* (0.13 ± 0.00 at 4.80
213 mg mL^{-1}). Most of the samples were better than the standards (only lime extract gave a
214 higher EC_{50} value than one of the standards, BHA); reducing power of BHA at 3.6 mg
215 mL^{-1} and α -tocopherol at 8.6 mg mL^{-1} was only 0.12 and 0.13 , respectively.

216 Lipid peroxidation inhibition, measured by the bleaching of β -carotene, is also
217 presented in **Figure 3**. β -Carotene undergoes a rapid decolourization in the absence of
218 an antioxidant since the free linoleic acid radical attacks the β -carotene molecule, which
219 loses the double bonds and, consequently loses its orange colour. The results obtained
220 for phenolic extracts were excellent and even better than the standard TBHQ (82.2% at

221 2.00 mg mL⁻¹). For *Eucalyptus gunnii*, phenolic extracts gave the highest β-carotene
222 bleaching inhibition percentages (e.g. 84.71 ± 6.96% at 0.31 mg mL⁻¹) while *Cupressus*
223 *lusitanica* gave the lowest (e.g. 43.51 ± 0.37% at 0.31 mg mL⁻¹). Once more, in the
224 essential oils extracts, the best contribution was given by *Cistus ladanifer* (66.32 ±
225 1.39% at 0.60 mg mL⁻¹) and the worst by *Citrus latifolia*.

226

227 For an overview of the results, EC₅₀ values (mg mL⁻¹) obtained in the different
228 antioxidant activity assays performed for phenolic and essential oils extracts from the
229 aromatic plants are shown in **Table 1**. Both kind of extracts of *Cistus ladanifer*, *Citrus*,
230 *Cupressus lusitanica* and *Eucalyptus gunnii* revealed interesting antioxidant properties.
231 Nevertheless, the phenolic extracts revealed better antioxidant properties (significantly
232 lower EC₅₀ values; $p \leq 0.05$) than essential oils extracts. After distillation extraction from
233 aromatic plants, several volatile molecules can be present in the samples such as
234 terpenes and terpenoids, aliphatic components, but also phenol-derived aromatic
235 components. Nevertheless, it is already known that thermal degradation of compounds
236 is one of the most significant disadvantages of the distillation extraction methodology
237 (Teixeira et al., 2007). In fact, our research group also reported the destruction of the
238 structures of polyphenols by heat which causes a decrease in the antioxidant activity
239 (Barros et al., 2007). This could explain the decrease in the antioxidant activity of
240 essential oils fraction (loss of phenolics and remain of the other volatile compounds)
241 relatively to the phenolic fraction (extracted at room temperature). Also, phenols might
242 be better antioxidants than terpenes, terpenoids or aliphatic components, since they
243 easily donate hydrogen atoms to quench the radicals formed in the DPPH and lipid

244 peroxidation inhibition assays, or electrons to reduce Fe³⁺/ferricyanide complex to the
245 ferrous form (Fe²⁺) in the reducing power assay.

246 The extraction yields obtained in essential oils extractions were much lower than those
247 obtained for the phenolic extraction (**Table 1**). Nevertheless, it was not observed any
248 relation between the extraction yields and the antioxidant activity EC₅₀ values obtained
249 in the different assays. Phenolic extracts (unless for DPPH scavenging properties of
250 lime extracts) revealed excellent antioxidant properties, even better than the well-known
251 standards BHA and TBHQ. These synthetic antioxidant compounds are added to an
252 extensive variety of foods in order to prevent or retard oxidation, so they are widely
253 used in the food industry (Adegoke et al., 1998) and are included in the human diet
254 (Leclercq et al., 2000). Nevertheless, the use of naturally occurring antioxidants has
255 been promoted because of concerns regarding the safety of synthetic antioxidants. The
256 possible activity of synthetic antioxidants as promoters of carcinogenesis has become a
257 concern. BHA and related antioxidants have been suggested to have toxic effects like
258 liver damage and mutagenesis (Grice, 1986; Wichi, 1988). Therefore, replacing
259 synthetic antioxidants with natural alternatives has attracted great interest over the last
260 years, and aromatic plants seem to be good candidates.

261 *Eucalyptus gunnii* phenolic extract revealed EC₅₀ values lower than 0.1 mg mL⁻¹, which
262 can be explained by the higher phenolic contents (176.07 ± 0.18 mg GAE g⁻¹; **Figure**
263 **4**). This value is similar to the content found in a previous study (Vázquez et al., 2009)
264 on aqueous *Eucalyptus globulus* bark extracts from Galicia (180.9 mg GAE g⁻¹). *Citrus*
265 *latifolia* revealed the lowest phenolic content (14.19 ± 0.48mg GAE g⁻¹; **Figure 4**)
266 which is in agreement with the highest EC₅₀ values obtained in all the antioxidant
267 activity assays. This value was lower than the phenolics found in another study

268 (Urbando-Rivera et al., 2005) in *Citrus latifolia* obtained from Mexico (20 mg GAE g⁻¹)
269 ¹). The EC₅₀ values obtained with the studied aromatic plants are very promissory being
270 better than the values obtained by our research group in different natural products such
271 as mushrooms (Barros et al., 2008), honey (Ferreira et al., 2009) or chestnut fruits
272 (Barreira et al., 2008). Furthermore, there are several reports on antioxidant properties
273 of phenolics and essential oils but, as far as we know, this is the first study comparing
274 the antioxidant properties of both extracts of these four aromatic plants.

275

276 **Acknowledgements**

277 The authors are grateful to the Foundation for Science and Technology, through CIMO
278 Research Centre for financial support of this work.

279

280 **References**

281 Adams, R.P., 2001. Identification of Essential Oils Components by Gas
282 Chromatography/Quadrupole Mass Spectrometry. Allured Publishing Coporation,
283 Carol Stream, IL, USA.

284 Adegoke, G.O., Kumar, M.V., Krishna, A.G.G., Varadaraj, M.C., Sambaiah, K., Lokesh,
285 B.R., 1998. Antioxidants and lipid oxidation in foods: A critical appraisal. J. Food
286 Sci. Technol. 35, 283-298.

287 Amaral, F.J., 1971. Nova flora de Portugal, vol. I, Continente e Açores, Lisboa.

288 Attaguile, L.G., Russo, A., Campisi, A., Savoca, F., Acquaviva, R., Ragusa, N., Vanella,
289 A. 2000. Antioxidant activity and protective effect on DNA cleavage of extracts
290 from *Cistus incanus* L. and *Cistus monspeliensis* L. Cell Biol. Toxicol. 16, 83-90.

291 Bakkali, F., Averbeck, S., Averbeck, D., Idaomar, M., 2008. Biological effects of
292 essential oils – A review. Food Chem. Toxicol. 46, 446-475.

293 Barreira, J.C.M., Ferreira, I.C.F.R., Oliveira, M.B.P.P., Pereira, J.A., 2008. Antioxidant
294 activities of the extracts from chestnut flower, leaf, skins and fruit. Food Chem.
295 107, 1106–1113.

296 Barros, L., Baptista, P., Correia, D.M., Morais, J.S., Ferreira, I.C.F.R., 2007. Effects of
297 conservation treatment and cooking on the chemical composition and antioxidant
298 activity of Portuguese wild edible mushrooms. J. Agric. Food Chem. 55, 4781-
299 4788.

300 Barros, L., Falcão, S., Baptista, P., Freire, C., Vilas-Boas, M., Ferreira, I.C.F.R., 2008.
301 Antioxidant activity of *Agaricus* sp. mushrooms by chemical, biochemical and
302 electrochemical assays. Food Chem. 111, 61–66.

303 Bravo, L. 1998. Polyphenols: chemistry, dietary sources, metabolism and nutritional
304 significance. Nutr. Rev. 11, 317–33.

305 Danne, A., Petereit, F., Nahrstedt, A., 1994. Flavan-3-ols, prodelphinidins and further
306 polyphenols from *Cistus salvifolius*. Phytochem. 37, 533-538.

307 Fakhari, A.R., Salehi, P., Heydari, R., Ebrahimi, S.N., Haddadb, P.R., 2005.
308 Hydrodistillation-headspace solvent microextraction, a new method for analysis of
309 the essential oil components of *Lavandula angustifolia* Mill. J. Chromatogr. A
310 1098, 14-18.

311 Ferreira, I.C.F.R., Aires, E., Barreira, J.C.M., Estevinho, L.M., 2009. Antioxidant
312 activity of Portuguese honey samples: Different contributions of the entire honey
313 and phenolic extract. Food Chem. 114, 1438-1443.

314 Grice, H.C., 1986. Safety evaluation of butylated hydroxytoluene (BHT) in the liver,
315 lung and gastrointestinal tract. Food Chem. Toxicol. 24, 1127-1130.

316 Hatano, T., Kagawa, H., Yashura, T., Okuda, T., 1988. Two new flavonoids and other
317 constituents in licorice root: their relative astringency and radical scavenging
318 effects. Chem. Pharm. Bull. 36, 2090-2097.

319 Kuiate, J.R., Bessièrè, J.M., Zollo, P.H.A., Kuate, S.P., 2006. Chemical composition and
320 antidermatophytic properties of volatile fractions of hexanic extract from leaves of
321 *Cupressus lusitanica* Mill. from Cameroon. J. Ethnopharmacol. 103, 160-165.

322 Lassak, E.V., McCarthy, T. 1983. Australian Medicinal Plants. North Ryde, Methuen
323 Australia, 240 p.

324 Leclercq, C., Arcella, D., Turrini, A., 2000. Estimates of the theoretical maximum daily
325 intake of erythorbic acid, gallates, butylated hydroxyanisole (BHA) and butylated
326 hydroxytoluene (BHT) in Italy: A stepwise approach. Food Chem. Toxicol. 38,
327 1075-1084.

328 Macheix, J., Fleuriet, A., 1998. Flavonoids in Health and Disease. Marcel Dekker Inc,
329 New York. 839-868.

330 Moyler, D.A., Clery, R.A., 1997. Plant Volatile Analysis, Springer, Berlin, p. 96.

331 Nijveldt, R., Nood, E., Hoorn, D., Boelens, P., Norren, K., Leeuwen, P., 2001.
332 Flavonoids: a review of probable mechanisms of action and potential applications.
333 Amer. J. Clin. Nutr. 74, 418-425.

334 Oyaizu, M., 1986. Studies on products of browning reactions: Antioxidative activities of
335 products of browning reaction prepared from glucosamine. Jap. J. Nutr. 44, 307-
336 315.

337 Shon, M.Y., Kim, T.H., Sung, N.J., 2003. Antioxidants and free radical scavenging
338 activity of *Phellinus baumii* (Phellinus of Hymenochaetaceae) extracts. Food
339 Chem. 82, 593-597.

340 Singleton, V.L., Rossi, J.A. jr, 1965. Colorimetric of total phenolics with
341 phosphomolybdic-phosphotungstic acid reagents. Am. J. Enol. Vit. 16, 144-158.

342 Teixeira, S., Mendes, A., Alves, A., Santos, L., 2007. Simultaneous distillation-
343 extraction of high volatile compounds from *Cistus ladanifer* L. Anal. Chim. Acta
344 584, 439-446.

345 Urbando-Rivera, J., Navarro-Ocaña, A., Valdivia-López, M.A., 2005. Mexican lime
346 peel: comparative study on contents of dietary fibre and associated antioxidant
347 activity. Food Chem. 89, 57-61.

348 Vázquez, G., González-Alvarez, J., Santos, J., Freire, M.S., Antorrena, G., 2009.
349 Evaluation of potential applications for chestnut (*Castanea sativa*) shell and
350 eucalyptus (*Eucalyptus globulus*) bark extracts. Ind. Crop. Prod. 29, 364-370.

351 Wichi, H.P., 1988. Enhanced tumour development by butylated hydroxyanisole (BHA)
352 from the prospective of effect on forestomach and oesophageal squamous
353 epithelium. Food Chem. Toxicol. 26, 717-723.


354

355


356

357 **Table 1.** Extraction yields (%) and EC₅₀ values (mg mL⁻¹) obtained for the antioxidant
 358 activity of the aromatic plants (mean ± SD; n=3). In each column different letters mean
 359 significant differences.

Samples	Extraction yield	DPPH scavenging activity	Reducing power	Lipid peroxidation inhibition	
Essential oils extract	<i>Cistus ladanifer</i>	0.63 ± 0.09 d	36.28 ± 0.36 d	4.00 ± 0.01 c	0.12 ± 0.01 d
	<i>Citrus latifolia</i>	1.26 ± 0.16 b	156.92 ± 0.40 b	23.64 ± 0.02 b	4.51 ± 0.03 b
	<i>Cupressus lusitanica</i>	0.80 ± 0.11 c	53.46 ± 0.70 c	4.38 ± 0.02 c	0.75 ± 0.08 c
	<i>Eucalyptus gunnii</i>	5.00 ± 0.89 a	272.93 ± 1.20 a	24.95 ± 0.07 a	5.49 ± 0.04 a
ANOVA	$p \leq 0.05$	$p \leq 0.05$	$p \leq 0.05$	$p \leq 0.05$	
Phenolic extract	<i>Cistus ladanifer</i>	41.24 ± 8.15 a	0.13 ± 0.02 c	0.19 ± 0.01 c	0.12 ± 0.01 c
	<i>Citrus latifolia</i>	15.18 ± 0.20 d	7.53 ± 0.76 a	19.25 ± 0.09 a	0.65 ± 0.02 a
	<i>Cupressus lusitanica</i>	29.84 ± 2.21 c	0.28 ± 0.01 b	0.47 ± 0.00 b	0.39 ± 0.01 b
	<i>Eucalyptus gunnii</i>	38.33 ± 3.12 b	0.10 ± 0.31 c	0.08 ± 0.00 d	0.05 ± 0.00 d
ANOVA	$p \leq 0.05$	$p \leq 0.05$	$p \leq 0.05$	$p \leq 0.05$	




360




361

362 **Figure 1.** Scavenging activity on DPPH radicals (%) of essential oils and phenolic
 363 extracts of aromatic plants. Each value is expressed as mean \pm SD (n=3). Standards:
 364 BHA (96% at 3.6 mg mL⁻¹) and α -tocopherol (95% at 8.6 mg mL⁻¹).

365



366




367


368 **Figure 2.** Reducing power of essential oils and phenolic extracts of aromatic plants.

369 Each value is expressed as mean \pm SD (n=3). Standards: BHA (0.12 at 3.6 mg mL⁻¹)

370 and α -tocopherol (0.13 at 8.6 mg mL⁻¹).




371



372

373 **Figure 3.** Lipid peroxidation inhibition of essential oils and phenolic extracts of
 374 aromatic plants. Each value is expressed as mean \pm SD (n=3). Standard: TBHQ (82.2%
 375 at 2 mg mL⁻¹).



376

377 **Figure 4.** Total phenolic content in the aromatic plants.