



UNIVERSITI PUTRA MALAYSIA
***ANTIOXIDATIVE PROPERTIES OF SELECTED
MALAYSIAN HERBAL PLANTS***

SUMAZIAN BT YUSUF

FBSB 2014 25



**ANTIOXIDATIVE PROPERTIES OF SELECTED
MALAYSIAN HERBAL PLANTS**

SUMAZIAN BT YUSUF

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2014



UPM
UNIVERSITI PUTRA MALAYSIA
BERILMU BERBAKTI

**ANTIOXIDATIVE PROPERTIES OF SELECTED
MALAYSIAN HERBAL PLANTS**

By

SUMAZIAN BT YUSUF

**Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

April 2014

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis dedicated to my parents, Yusuf Isa and Haminah Hamat, my sisters, Suriani Yusuf and Suhazilah Yusuf and my brother, Suhaifa Yusuf, last but not least to my beloved husband Mohd Zulkefly Ramli and all of my family members who have been a great support, motivation and prayers since the beginning of my studies.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Science

ANTIOXIDATIVE PROPERTIES OF SELECTED MALAYSIAN HERBAL PLANTS

By

SUMAZIAN BT YUSUF

April 2014

Chairperson : Professor Maziah Mahmood, PhD

Faculty : Biotechnology and Biomolecular Sciences

Antioxidant activities of nine herbal plants which are commonly used in Malaysia as folk medicines, namely *Barringtonia racemosa* (putat), *Curcuma domestica* (kunyit), *Kaempferia galanga* (cekur), *Hydrocotyle bonariensis* (pegaga embun), *Centella asiatica* (pegaga), *Piper betel* (sireh), *Polygonum minus* (kesum), *Piper sarmentosum* (kadok) and *Cosmos caudatus* (ulam raja) were examined. All of the herbs selected for all readily available for everyday use with a variety of uses well known since time immemorial. The study was conducted to see the appropriate dose to be used for commercial purposes. The compounds in leaf and rhizome were extracted using hot water (100 °C), fresh water and ethanol (80%), separately. Total antioxidant content were measured using ferric reducing power (FRAP), 1,1 diphenyl -2- picrylhydrazyl free radical scavenging assay (DPPH) and β -carotene bleaching assay. Subsequently, the relevant antioxidant compounds; total carotenoids, total ascorbic, total flavonoids and total phenolic. Enzymatic antioxidants include superoxide dismutase (SOD), ascorbate peroxidase (APX), glutathione reductase (GR), catalase (CAT) and peroxidase (POX) activities were determined. Anti-aging enzymes activities; anti-tyrosinase, anti-elastase and anti-hyaluronidase were also determined using spectrophotometry method. For FRAP assay, hot water extract of *C. domestica* (623.87 ± 0.01 mmol Fe²⁺/g DW) showed good activity and fresh water extract of *K. galanga* exhibited a very strong antioxidant properties in DPPH assay with ($83.27 \pm 0.01\%$). *H. bonariensis* leaf extract has been recognized to contain high antioxidant activity with (19.04 ± 0.01 %) using β -carotene bleaching assay. The rhizome extract of *C. domestica* showed the highest total carotenoid content of (0.04 ± 0.01 mg/mg FW) and leaf extract of *P. sarmentosum* indicated (1.72 ± 0.01 mg/g FW) for the highest total of ascorbic acid content. Level of total flavonoids in the herbal plants varied between (0.42 ± 0.00 and 6.28 ± 0.01 mg/g DW) and those of total phenolics between (2.78 ± 0.01 and 12.35 ± 0.01 mg/g DW). For enzymatic antioxidant activities, the results showed that *C. domestica* (0.42 ± 0.01 μ mol/min/g FW), *P. sarmentosum* (0.14 ± 0.01 μ mol/min/g FW), *C. asiatica* (0.76 ± 0.01 μ mol/min/g FW), *H. bonariensis* (0.92 ± 0.01 μ mol/min/g FW) and *P. minus* (0.38 ± 0.01 μ mol/min/g FW) were higher in SOD, APX, GR, CAT and POX activity, respectively. The highest percentage of anti-tyrosinase activity is in fresh water extract of *C. domestica* rhizome (97.86 ± 0.02 %) using L-tyrosine as substrate. *C.*

asiatica leaf exhibited a very good level of elastase and hyaluronidase inhibition with $(84.53 \pm 0.02\%)$ and $(81.01 \pm 0.01\%)$, respectively. The findings showed the potential of nine medicinal plants as powerful antioxidant that can be used in producing herbal based product in cosmetically and pharmaceutical.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

PENILAIAN ANTIOKSIDAN DALAM TUMBUHAN HERBA TERPILIH DI MALAYSIA

Oleh

SUMAZIAN BT YUSUF

April 2014

Pengerusi : Profesor Maziah Mahmood, PhD

Fakulti : Bioteknologi dan Sains Biomolekul

Aktiviti-aktiviti antioksidan di dalam sembilan jenis tumbuhan herba di Malaysia yang popular sebagai ubatan tradisional dikenali sebagai *Barringtonia racemosa* (putat), *Curcuma domestica* (kunyit), *Kaempferia galanga* (cekur), *Hydrocotyle bonariensis* (pegaga embun), *Centella asiatica* (pegaga), *Piper betel* (sireh), *Polygonum minus* (kesum), *Piper sarmentosum* (kadok) dan *Cosmos caudatus* (ulam raja) telah diuji. Kesemua herba yang dipilih kerana semuanya senang didapati untuk kegunaan harian juga dikenali dengan pelbagai kegunaan semenjak dahulu lagi. Kajian dijalankan untuk melihat dos yang sesuai untuk digunakan bagi tujuan komersial. Pengestrakan daun dan rizom ditunjukkan melalui tiga kaedah berlainan iaitu air panas (100 °C), air dan etanolik (80%). Jumlah antioksidan diukur secara penurunan ferik (FRAP), '1,1 diphenyl -2-picrylhydrazyl (DPPH) pengikatan radikal bebas dan penyekatan pelunturan β-karotin. Untuk kandungan antioksidan yang relevan pula termasuk jumlah karotenoid, asid askorbik, flavonoid dan fenolik. Manakala aktiviti antioksidan enzim pula adalah seperti 'Superoxide dismutase (SOD)', 'Ascorbate peroxidase (PRX)', 'Glutathione reductase (GR)', 'Catalase (CAT)' dan 'Peroxidase (POX)'. Di samping itu, jumlah kandungan gula (hidrolisis dan bukan hidrolisis) serta enzim anti-penuaan seperti 'anti-tyrosinase', 'anti-elastase' dan 'anti-hyaluronidase' juga dianalisa menggunakan spektrofotometer. Untuk kaedah FRAP, ekstrak air panas *C. domestica* (623.87 ± 0.00 mmol $^{Fe^{2+}}$ /g berat kering) menunjukkan aktiviti terbaik dan ekstrak air *K. galanga* menunjukkan aktiviti antioksidan paling kuat untuk kaedah DPPH iaitu ($83.27 \pm 0.01\%$). Manakala *H. bonariensis* didapati mengandungi aktiviti antioksidan paling tinggi iaitu ($19.04 \pm 0.01\%$) melalui pelunturan β-karotin. Ekstrak rizom *C. domestica* menunjukkan jumlah karotenoid tertinggi iaitu sebanyak (0.04 ± 0.01 mg/g berat basah) dan ekstrak daun *P. sarmentosum* menunjukkan (1.72 ± 0.01 mg/g berat basah) untuk kandungan asid askorbik tertinggi. Paras kandungan flavonoid dalam pelbagai tumbuhan herba adalah antara (0.42 ± 0.01 dan 6.28 ± 0.01 mg/g berat kering) manakala jumlah fenolik pula adalah antara (2.78 ± 0.01 dan 12.35 ± 0.01 mg/g berat kering). Aktiviti-aktiviti antioksidan enzim telah didapati didalam *C. domestica* (0.42 ± 0.01 μmol/min/g berat basah), *P. sarmentosum* (0.14 ± 0.01 μmol/min/g berat

basah), *C. asiatica* (0.76 ± 0.01 $\mu\text{mol}/\text{min}/\text{g}$ berat basah), *H. bonariensis* (0.92 ± 0.01 $\mu\text{mol}/\text{min}/\text{g}$ berat) dan *P. minus* (0.38 ± 0.01 $\mu\text{mol}/\text{min}/\text{g}$ berat basah) masing-masing tertinggi didalam SOD, APX, GR, CAT dan POX. Peratusan aktiviti anti-tyrosinase yang tertinggi adalah ($97.86 \pm 0.02\%$) didalam ekstrak air rizom *C. domestica* dimana L-tyrosine digunakan sebagai substrat. Ekstrak-ekstrak daun *C. asiatica* yang berbeza menunjukkan perencatan enzim elastase dan hyaluronidase yang sangat baik iaitu masing-masing dengan ($84.53 \pm 0.02\%$) dan ($81.01 \pm 0.01\%$). Penemuan ini telah menunjukkan potensi sembilan ekstrak herba sebagai antioksidan yang berkuasa dan berguna dalam penghasilan produk-produk yang berasaskan herba di dalam industri kosmetik dan farmaseutikal.



ACKNOWLEDGEMENTS

Alhamdulillah, praise be to Allah for His blessing for me to finish up this thesis. This task would not be complete and possible without help, co-operation and encouragement of a large number of peoples in my life. Special thanks to my supervisor, Prof Dr. Maziah Mahmood and my co supervisor, Dr Syahida Ahmad for their guidance, encouragement, help and patience throughout the whole project.

My graceful thank to all my lab mates for their help, comments, critics and recommendation. Special thanks also to Mohd Hakiman for the help on editing my thesis. I also would like to thank to Universiti Putra Malaysia for Grant Research Fellowship (GRF) for two years of financial support. Lastly, I would like to express my appreciation to my family for their encouragement, understanding, advice and support.



I certify that a Thesis Examination Committee has met on 9 April 2014 to conduct the final examination of Sumazian bt Yusuf on her thesis entitled "Antioxidative Properties of Selected Malaysian Herbal Plants" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Noor Azmi Shaharuddin, PhD

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Nor' Aini bt. Abdul Rahman, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Noorjahan Banu binti Mohammed Alitheen, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Aziz bin Ahmad, PhD

Associate Professor

Universiti Malaysia Terengganu

Malaysia

(External Examiner)



NORITAH OMAR, PhD

Associate Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 23 June 2014



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Maziah Mahmood, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Syahida Ahmad, PhD

Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 23 July 2014

Declaration by Graduate Student

I hereby confirm that:

- This thesis is my original work;
- Quotations, illustrations and citations have been duly referenced;
- This thesis has not been submitted previously or concurrently for other degree at any other institutions;
- Intellectual property from the thesis and copyright of thesis are fully-owned by University Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- Written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is publish (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matrix No: _____

Declaration by Members of Supervisory Committee

This is to confirm that:

- The research conducted and the writing of this thesis was under our supervision;
- Supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: **Maziah Mahmood, PhD**

Signature: _____
Name of
Member of
Supervisory
Committee: **Syahida Ahmad, PhD**

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF APPENDICES	xv
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Medicinal Herbs	4
2.1.1 <i>Barringtonia racemosa</i>	5
2.1.2 <i>Curcuma domestica</i>	5
2.1.3 <i>Kaempferia galangal</i>	6
2.1.4 <i>Hydrocotyle bonariensis</i>	6
2.1.5 <i>Centella asiatica</i>	7
2.1.6 <i>Piper betel</i>	7
2.1.7 <i>Piper sarmentosum</i>	8
2.1.8 <i>Polygonum minus</i>	8
2.1.9 <i>Cosmos caudatus</i>	9
2.2 Plant Antioxidants	11
2.3 Functions of Antioxidants	12
2.4 Types of Antioxidants	12
2.4.1 β -carotene	12
2.4.2 Carotenoid	13
2.4.3 Ascorbic acid	13
2.4.4 Flavonoid	14
2.4.5 Phenolic	15
2.5 Enzymatic Antioxidants	16
2.5.1 Superoxide Dismutase (SOD)	16
2.5.2 Ascorbate Peroxidase (APX)	17
2.5.3 Glutathione Reductase (GR)	17
2.5.4 Catalase (CAT)	18
2.5.5 Peroxidase (POX)	19
2.6 Enzymes Inhibitors	20
2.6.1 Tyrosinase Inhibition	20
2.6.1.1 Melanogenesis	22
2.6.2 Hyaluronidase Inhibition	24

2.6.3 Elastase Inhibition	26
3 MATERIALS AND METHODS	
3.1 Chemicals and reagents	28
3.2 Plant materials	28
3.3 Total Antioxidants	35
3.3.1 Extractions of Ferric reducing antioxidant potential power (FRAP) and 1,1 diphenyl-2- picrylhydrazyl (DPPH)	35
3.3.1.1 Ferric reducing antioxidant potential power (FRAP)	35
3.3.1.2 1,1 diphenyl-2-picrylhydrazyl(DPPH)	35
3.3.2 β -carotene Content	36
3.3.2.1 Extraction of sample	36
3.3.2.2 β -carotene Bleaching assay	36
3.4 Quantification of Relevant Antioxidant Compounds	37
3.4.1 Total Carotenoid Content	37
3.4.2 Total Ascorbic Acid Content	37
3.4.3 Total flavonoid Content	37
3.4.4 Total Phenolic Content	38
3.5 Determination of Enzymatic Antioxidants	38
3.5.1 Preparations of Enzyme Extracts	38
3.5.1.1 Superoxide Dismutase Activity	38
3.5.1.2 Ascorbate Peroxidase Activity	39
3.5.1.3 Glutathione Reductase Activity	39
3.5.1.4 Catalase Activity	39
3.5.1.5 Peroxidase activity	39
3.6 Enzymes Inhibition	40
3.6.1 Tyrosinase Inhibition	40
3.6.2 Extraction for Hyaluronidase and Elastase Inhibition	40
3.6.2.1 Hyaluronidase inhibition	41
3.6.2.2 Elastase inhibition	41
3.7 Statistical analysis	41
4 RESULTS AND DISCUSSION	42
4.1 Total Antioxidant Activity	42
4.1.1 Ferric Reducing Antioxidant Potential Assay (FRAP)	42
4.1.2 Total Antioxidant Content using 1,1-diphenyl-2-Picrylhydrazyl (DPPH) Free Radical Scavenging Assay	45
4.1.3 β -carotene Bleaching Assay	48
4.2 Relevant Antioxidant Compounds	50
4.2.1 Total Carotenoid Content	50
4.2.2 Total Ascorbic Acid Content	52
4.2.3 Total Flavonoid Content	54
4.2.4 Total Phenolic Content	57
4.3 Enzymatic Antioxidant	59
4.3.1 Superoxide Dismutase Activity	59
4.3.2 Ascorbate Peroxidase Activity	61
4.3.3 Glutathione Reductase Activity	63

4.3.4 Catalase Activity	65
4.3.5 Peroxidase Activity	67
4.4 Enzyme Inhibitors	69
4.4.1 Tyrosinase Inhibition Using L-tyrosine As Substrate	69
4.4.2 Tyrosinase Inhibition Using L-Dopa As Substrate	71
4.4.3 Hyaluronidase Inhibition	73
4.4.4 Elastase Inhibition	75
4.5 Correlation of between antioxidant, compounds and enzyme activities	78
5 CONCLUSION	
5.1 Conclusion	81
5.2 Future Research	82
REFERENCES	83
APPENDICES	109
BIODATA OF STUDENT	116
LIST OF PUBLICATION	117

LIST OF TABLES

Table		Page
2.1	Secondary metabolites produced in selected Malaysia herbal plants in this study	10
2.2	List of plant based tyrosinase inhibitors that available in market	21
2.3	List of chemical agents that influence in cosmetics and pharmaceuticals as whitening agents	23
2.4	List of plant based hyaluronidase inhibitors that available in market	25
2.5	List of plant based elastase inhibitors that available in market	27

LIST OF FIGURES

Figure		Page
3.1	Malaysian herbs used in this study	29
4.1	Total Antioxidant Content Using Ferric Reducing Potential assay (FRAP)	44
4.2	Total Antioxidant Content Using 1,1-diphenyl-2-picrylhydrazyl (DPPH) Free Radical Scavenging Assay	47
4.3	Total Antioxidant Content Using β -carotene Bleaching Assay	49
4.4	Total Carotenoid Content	51
4.5	Total Ascorbic Acid Content	53
4.6	Total Flavonoid Content	56
4.7	Total Phenolic Content	58
4.8	Superoxide Dismutase Activity	60
4.9	Ascorbate Peroxidase Activity	62
4.10	Gluthathione Reductase Activity	64
4.11	Catalase Activity	66
4.12	Peroxidase Activity	68
4.13	Tyrosinase Inhibition (using L-tyrosine as substrate)	70
4.14	Tyrosinase Inhibition (using L-dopa as substrate)	72
4.15	Hyaluronidase Inhibition	74
4.16	Elastase Inhibition	77

LIST OF APPENDICES

Appendix		Page
A	Standard graph FRAP value	109
B	Standard graph of total flavonoid	110
C	Standard graph of total phenolic acid	111
D	Standard graph of carotenoid	112
E	Standard graph of ascorbic acid	113
F	Standard graph of protein	114
G	Pearson's correlations between total antioxidant, compounds and enzyme activities of selected Malaysian herbal plants	115

LIST OF ABBREVIATIONS

Abs	Absorbance
AlCl ₃	Aluminium chloride
APX	Ascorbate peroxidase
BHA	Butylated hydroxyanisole
BHT	Butylated hydrotoluene
BSA	Bovine Serum Albumin
CAT	Catalase
Cu	Copper
Cd	cadmium
°C	Degree Celcius
DCPIP	2,6-dichloroindophenol
DPPH	2, 2-diphenyl-1-picrylhydrazyl
DNA	Deoxyribonucleic acid
DNS	3,5-dinitrosalicylic acid
DNTB	5, 5-dithio-bis-(2-nitrobenzoic acid)
DW	Dry weight
EDTA	Ethylenediaminetetraacetic acid
<i>et al.,</i>	<i>at alli</i> and other people
Fe	Ferum
Fe ²⁺	Ferrous
FeCl ₃	Ferum chloride
FRAP	Ferric reducing antioxidant power

FW	Fresh weight
G	Gram
GAE	Gallic acid
GlcNAc	N-acetyl-D-glucosamine
GlcUAc	d-glucuronic acid
GR	Glutathione reductase
GSH	Glutathione disulphide
GSSG	Glutathione sulfhydryl
HCl	Hydrochloric acid
Hr	Hour
H ₂ O	Water
H ₂ O ₂	Hydrogen peroxide
KCN	Potassium cyanide
KOH	Potassium hydroxide
Mg	Milligram
Mg/L	Milligram per liter
Min	Minute
ml	Milliliter
mM	Millimolar
Mn	Manganese
μg	Microgram
μl	Microliter
μmol	Micromol
Na ₂ CO ₃	Sodium carbonate
NADPH	Glutamate synthase

NaNO ₂	Sodium nitrate
NaOH	Sodium hydroxide
NBT	Nitroblue tetrazolium salt
Ni	Nickel
nm	Nanometer
O ₂ ⁻	Superoxide
Pb	Plumbum
POX	Peroxidase
PVPP	Polyvinylpolypyrrolidone
%	Percentage
rpm	Rotation per minute
ROS	Reactive oxygen species
SD	Standard deviation
SOD	Superoxide dismutase
TPTZ	Tripyridyltriazine
UV	Ultraviolet
v/v	volume per volume
w/v	weight per volume
Zn	Zinc

CHAPTER 1

INTRODUCTION

Medicinal herbs have long been used as natural source of medicines. Since a long time ago in the past era, herbs were used in traditional therapy, cosmetics and health supplements. According to WHO report, between the years of 1991 to 2001, Malaysia is among the nine countries in this world that gave big amount of sales in herbal medicine where the market of herbal and natural products in Malaysia was estimated to worth approximately RM 10 billion and the rate were increasing about 8% per year in 2008 (Effendy *et al.*, 2012).

Plants are sources of antioxidants that counteract reactive oxygen species (ROS) which include activated and/or free radical oxygen compounds, such as singlet oxygen, superoxide anion radical, hydroxyl radical and hydrogen peroxide (Huda-Faujan *et al.*, 2007). These species can attack tissues in the dermis or epidermis which cause skin aging and DNA damage (Shon *et al.*, 2003). Therefore, antioxidants from plant origins are increasing in demand in the market as source of natural antioxidants since artificial antioxidants such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), and tertiary butylhydroquinone (TBHQ) which are widely used in the industry are toxic and carcinogenic.

In Malaysia, 20% (3,000 species) of the *angiosperma* and *gymnosperma* plant species have been used for traditional treatments for ages and was reported to have medicinal benefits for a long time ago (Zainol *et al.*, 2003). Since herbs have high commercial value, improvement of its quality is important due to its medicinal properties. It was reported that 10-15% of the known higher plant species had been investigated for their important bioactive compounds (Zarate *et al.*, 2001). Due to the potential of plants for medicinal and cosmetically purposes, the biochemical compounds need to be investigated using various techniques because there is lack of information on its biochemical constituents that has been reported. Plant antioxidants are believed to play a role in delaying aging process, because of the several compounds such as flavonoids that contribute to antioxidative properties (Harbone and Williams, 2000). This compound was commonly being used as skin improvement products and their applications in cosmetic industries have also been established. Vitamin C and β -carotene also contribute to antioxidant properties (Chanwitheesuk *et al.*, 2005; Lisiewska *et al.*, 2006). In addition, the compounds can be produced at reasonable low costs, especially when the right natural sources have been identified, for example, anti-wrinkle products which contain elastase and hyaluronidase inhibitors have been sold in the market.

During aging process, the elasticity of the skin decreases because of elastase enzyme that leads to sagging decreases. At the same time, hyaluronic acid also decreases, causing the skin to become dry and wrinkle. The best way to maintain youthful skin is to degrade the enzymes that can cause aging. Tyrosinase is an enzyme inhibitor that can inhibit melanisation process of human skin (Lim et al., 2009). There are a few numbers of synthetic and natural tyrosinases that have been determined but some activities are not significant or suitable which do not comply with the safety regulations for food and cosmetic additives. Therefore, there is a need to find and determine new tyrosinase inhibitors with high activities and decrease the side effects at the same time (Yan *et al.*, 2009). Herbal products are known to have interesting benefits, and they are largely unexplored sources of new and potential drugs. It is for this reason that the emphasis has been given on the potential of herbal medicines for new skincare cosmetics development (Kiken and Cohen, 2002). There are certain plant metabolites which are believed to have the capability to inhibit tyrosinase, hyaluronidase and elastase enzymes such as whitening cream and lotion from *Areca cathecu* or locally known as 'pinang' (Binic *et al.*, 2013).

There are many potential herbs that can be commercialized in Malaysia and most of them are subjected to extensive research. In this study, nine species of medicinal herbs were determined the total antioxidants and anti-aging inhibitors activities. They were *Barringtonia racemosa* (Lecythidaceae) or (putat), *Curcuma domestica* (Zingiberaceae) or (kunyit), *Kaempferia galanga* (Zingiberaceae) or (cekur), *Hydrocotyle bonariensis* (Apiaceae) or (pegaga embun), *Centella asiatica* (Apiaceae) or (pegaga), *Piper betel* (Piperaceae) or (sireh), *Piper sarmentosum* (Piperaceae) or (kadok), *Polygonum minus* (Polygonaceae) or (kesum) and *Cosmos caudatus* (Compositae) or (ulam raja). The selection of plant samples for phytochemical screening is generally based on their own special characteristics for traditional anti-aging treatments. All these nine plants also can be found easily in market and popular among Malaysian peoples. This study may provide information regarding the ranges of antioxidative activity and general antioxidant compounds of the selected herbal plants toward the standardization. The informations in this study will be useful in preparing the herbal formulations for health supplements.

The biochemical properties of selected herbs were carried out by analysing the important biochemical compounds which include total antioxidant activities using different method such as Ferric Reducing Antioxidant Power (FRAP), Free Radical Scavenging Assay (DPPH), total flavonoids and phenolic. Besides that, β -carotene, carotenoids and ascorbic acid also determined. In addition, there are enzyme activities such as superoxide dismutase, ascorbate peroxidase, glutathione reductase, catalase, peroxidase and enzyme inhibitors which include tyrosinase, hyaluronidase and elastase activity.

The objectives are:

1. To evaluate the total antioxidant contents of selected Malaysian herbs such as FRAP, DPPH and β -carotene.
2. To examine the content of antioxidant compound: total carotenoid, total ascorbic acid, total flavonoids and total phenolic.
3. To determine activity of the enzymatic antioxidants such as superoxide dismutase, ascorbate peroxidase, glutathione reductase, catalase and peroxidase.
4. To evaluate the tyrosinase, hyaluronidase and elastase enzyme inhibitors in the selected Malaysian herbs.

REFERENCES

- Aebi, H. 1984. Catalase *in vitro*. *Methods in Enzymology* 105: 121-126.
- Aherne, S.A. and O'Brien, N.M. 2002. Dietary flavonols: Chemistry, food content and metabolism. *Nutrition* 18: 75-81.
- Aida Azlina, A., Farihah, H.S., Qodriyah, H.M.S. and Nur Azlina, M.F. 2009. Effects of *Piper sarmentosum* water extract on 11- β hydroxysteroid dehydrogenase type 1 bioactivity in ovariectomy-induced obese rat. *International Journal of Pharmacology* 5: 362-369.
- Aizawa, K. and Inakuma, T. 2007. Quantitation of carotenoids in commonly consumed vegetables in Japan. *Food Science Technology Research* 13: 247-252.
- Ajani, E.O., Salako, A.A., Sharlie, P.D., Akinleye, W.A., Adeoye, A.O., Salau, B.A. and Adebawo, O.O. 2009. Chemopreventive and remediation effect of *Hydrocotyl bonariensis* omm. Ex Lam (Apiaceae) leave extract in galactose induced cataract. *Journal of Ethnopharmacology* 123: 134-142.
- Akowuah, G.A., Ismail, Z., Norhayati, I. and Sadikun, A. 2005. The effects of different extraction solvents of varying polarities on polyphenols of *Orthosiphon stamineus* and evaluation of the free radical-scavenging activity. *Food Chemistry* 93: 311-317.
- Al-Azzawie, H.F. and Alhamdani, M.S.S. 2006. Hypoglycemic and antioxidant effect of oleuropein in alloxan-diabetic rabbits. *Life Sciences* 78: 1371-1377.
- Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K. and Walter, P. 2002. The amino acid sequence of bovine liver catalase: A preliminary report. *Archives Biochemistry Biophysics* 131: 653-655.
- Amin, I., Norazaidah, Y. and Emmy Hainida, K.I. 2006. Antioxidant capacity and phenolic content of spinach species (*Amaranthus* sp.). *Food Chemistry* 94: 47-52.
- An, B.J., Kwak, J.H., Park, J.M., Lee, J.Y., Park, T.S., Lee, J.T., Son, J.H., Jo, C. and Byun, M.W. 2005. Inhibition of enzyme activities and the anti-wrinkle effect of polyphenol isolated from the persimmon leaf (*Diospyros kaki* folium) on human skin. *Dermatologic Surgery* 31: 848-854.

- Antonicelli, F., Bellon, G., Debelle, L. and Hornebeck, W. 2007. Elastin-elastases and inflamm-aging. *Current Topics in Developmental Biology* 79: 99-155.
- Arai, Y., Watanabe, S., Kimira, M., Shimoi, K., Mochizuki, R. and Kinae, N. 2000. Dietary intakes of flavonols, flavones and isoflavones by Japanese women and the inverse correlation between quercetin intake and plasma LDL cholesterol concentration. *Journal of Nutrition* 130: 2243-2250.
- Atamna, H. and Boyle, K. 2006. Amyloid- β peptide binds with heme to form a peroxidase: Relationship to the cytopathologies of Alzheimer's disease. *Proceedings of the National Academy of Science* 103: 3381-3386.
- Azizah, O., Amin, I., Nawalyah, A.G. and Ilham, A. 2007. Antioxidant capacity and phenolic content of cocoa beans. *Food Chemistry* 100: 1523-1530.
- Azlim, A.A.A., Ahmed Jalal, K.C., Syed, Z.I., Mustapha S.K., Aisyah, M.R. and Kamarul R.K. 2010. Total phenolic content and primary antioxidant activity of methanolic and ethanolic extracts of aromatic plants' leaves. *Journal of International Food Research* 17: 1077-1084.
- Bahroun, T., Luximon-Ramma, A., Crozier, A. and Aruoma, O.I. 2004. Total phenol, flavonoid, proanthocyanidin and vitamin C levels and antioxidant activities of Mauritian vegetables. *Journal of Science Food and Agriculture* 84: 1553-1561.
- Baurin, N., Arnoult, E., Scior, T., Do, Q.T. and Bernard, P. 2002. Preliminary screening of some tropical plants for anti-tyrosinase activity. *Journal of Ethnopharmacology* 82: 155-158.
- Beauchamp, C. and Fridovich, I. 1981. Superoxide dismutase: Improve assays and an assay applicable to acrylamide gels. *Annals of Biochemistry* 44: 276-287.
- Behbahani, M., Ali, A.M. and Muse, R. 2007. Plant regeneration from leaf explants of *Barringtonia racemosa*. *Journal of Medicinal Plant Research* 5: 103-108.
- Belkhadir, Y., Subramaniam, R. and Dangl, J. 2004. Plant disease resistance protein signaling: NBS-LRR proteins and their partners. *Current Opinion Plant Biology* 7: 391-399.
- Benkovic, V., Nada, O., Horvat, K.E.S., Rami, C., Domagoj, D.C., Ivan, B.C. and Nevenka, K. 2008. Evaluation of the radioprotective effects of propolis and flavonoids in gamma-irradiated mice: The alkaline comet assay study. *Biology Pharmaceutical Bulletin* 31: 167-172.

- Bernard, P. and Berthon, J.Y. 2000. Resveratrol: An original mechanism on tyrosinase inhibition. *International Journal of Cosmetic Sciences* 22: 219-226.
- Bhattacharya, S., Mula, S., Gamre, S., Kamat, J.P., Bandyopadhyay, S.K. and Chattopadhyay, S. 2007. Inhibitory property of *Piper betel* extract against photosensitization-induced damages to lipids and proteins. *Food Chemistry* 100: 1474-1480.
- Binic, I., Lazarevic, V., Ljubenovic, M., Mojsa, J. and Sokolovic, D. 2013. Skin ageing: Natural weapons and strategies. *Evidence-based Complementary and Alternative Medicine* 13: 111-121.
- Blokhina, O.B., Virolainen, E. and Fagerstedt, K.V. 2003. Antioxidants, oxidative damage and oxygen deprivation stress. Review article. *Annals of Botany* 91: 179-194.
- Bonfill, M., Mangas, S., Cusidó, R.M., Osuna, L., Piñol, M.T. and Palazón, J. 2006. Identification of triterpenoid compounds of *Centella asiatica* by thin-layer chromatography and mass spectrophotometry. *Biomedical Chromatography* 20: 151-153.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilising the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254.
- Briganti, S., Camera, E. and Picardo, M. 2003. Chemical and instrumental approaches to treat hyperpigmentation. *Pigment Cell Research* 16: 101-110.
- Brinkhaus, B., Lindner, M., Schuppan, D. and Hahn, E.G. 2000. Chemical, pharmacological and clinical profile of the East Asian medical plant *Centella asiatica*. *Phytomedicine* 7: 427-428.
- Brunet, J.M.C., Cetkovic, G.S., Djilas, S.M., Tumbas, V.T., Savatovic, T.S.S., Markov, A.I.M.S.L. and Cvetkovic, O.D. 2009. Radical scavenging and antimicrobial of horsetail (*Equisetum arvense* L.) extracts. *International Journal of Food Science Technology* 44: 269-278.
- Bunawan, H., Talip, N. and Noor, N.M. 2011. Foliar anatomy and micromorphology of *Polygonum minus* Huds. and their taxonomic implications. *Australian Journal of Crop Science* 5: 123-127.

- Cai, L. and Cherian, M.G. 2003. Zinc-metallothionein protects from DNA damage induced by radiation better than glutathione and copper or cadmium metallothioneins. *Toxicology Letters* 136: 193-198.
- Cals-Grierson, M.M. and Ormerod, A.D. 2004. Nitric oxide function in the skin. *Nitric Oxide* 10: 179-193.
- Chan, E.W.C., Lim, Y.Y., Wong, L.F., Lianto, F.S., Wong, S.K., Lim, K.K., Joe, C.E. and Lim, T.Y. 2008. Antioxidant and tyrosinase inhibition properties of leaves and rhizomes of ginger species. *Food Chemistry* 109: 477-483.
- Chang, T.S. 2009. An updated review of tyrosinase inhibitors. *International Journal of Molecular Sciences* 10: 2440-2475.
- Chanwitheesuk, A., Teerawutgulrag, A. and Rakariyatham, N. 2005. Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand. *Food Chemistry* 101: 515-523.
- Chelikani, P., Fita, I. and Loewen, P.C. 2004. Diversity of structures and properties among catalases. *Cellular and Molecular Life Sciences* 61: 192-208.
- Chen, S.W., Wang, W.J., Li, W.J., Wang, R., Li, Y.L., Huang, Y.N., and Liang, X. 2006. Anxiolytic-like effect of asiaticoside in mice. *Pharmacology Biochemistry and Behavior* 85: 339-344.
- Chen, Q.X. and Kubo, I. 2002. Kinetics of mushroom tyrosinase inhibition by quercetin. *Journal of Agricultural and Food Chemistry* 50: 4108-4112.
- Cheng, C.L. and Koo, M.W.L. 2000. Effects of *Centella asiatica* on ethanol induced gastric mucosal lesions in rats. *Life Sciences* 67: 2647-2653.
- Chitra, M., Martin, K.P., Sunandakumari, C. and Madhusoodanan, P.V. 2005. Protocol for rapid propagation and to overcome delayed rhizome formation in field established *in vitro* derived plantlets of *Kaempferia galangal* L. *Scientia Horticulture* 104: 113-120.
- Choi, S.S., Han, K.J., Lee, J.K., Lee, H.K., Han, E.J., Kim, D.H. and Suh, H.W. 2001. Antinociceptive mechanisms of orally administered decursinol in the mouse. *Life Sciences* 73: 471-485.
- Corpas, F.J., Barroso, J.B. and del Río, L.A. 2001. Peroxisomes as a source of reactive oxygen species and nitric oxide signal molecules in plant cells. *Trends of Plant Science* 6: 145-50.

- Corpas, F.J., Fernández-Ocaña, A., Carreras, A., Valderrama, R., Luque, F., Esteban, F.J., Rodríguez-Serrano, M., Chaki, M., Pedrajas, J.R., Sandalio, L.M., del Río, L.A. and Barroso, J.B. 2006. The expression of different superoxide dismutase forms is cell-type dependent in olive (*Olea europaea* L.) leaves. *Plant Cell Physiology* 47: 984-94.
- Cousins, M., Adelberg, J., Chen, F. and Rieck, J. 2007. Antioxidant capacity of fresh and dried rhizomes from four clones of turmeric (*Curcuma longa* L.) grown *in vitro*. *Industrial Crops and Products* 25: 129-135.
- Dasgupta, N. and De, B. 2004. Antioxidant activity of *Piper betel* L. leaf extract *in vitro*. *Food Chemistry* 88: 219-224.
- Datta, A., Ghoshdastidar, S. and Singh, M. 2011. Antimicrobial property of *Piper betel* leaf against clinical isolates of bacteria. *International Journal of Pharma Sciences and Research* 2: 104-109.
- Datta, K., Sinha, S. and Chattopadhyay, P. 2000. Reactive oxygen species in health and disease. *The National Medical Journal of India* 13: 304-310.
- Davey, M.W., Van Montagu, M., Inzé, D., Sanmartin, M., Kanellis, A., Smirnoff, N., Benzie, I. J.J., Strain, J.J., Favell, D. and Fletcher, J. 2000. Plant L-ascorbic acid: Chemistry, function, metabolism, bioavailability and effects of processing. *Journal of the Science of Food and Agriculture* 80: 825-860.
- Davis, S.H.R. and Masten, S.J. 1991. Spectrophotometric method for ascorbic acid using dichlorophenolindophenol: Elimination of the interference due to iron. *Analytica Chimica Acta* 248: 225-227.
- Deb, N., Majumdar, P. and Ghosh, A.K. 2013. Pharmacognostic and phytochemical evaluation of the rhizomes of *Curcuma longa* Linn. *Journal of Pharma SciTech* 2:81-86.
- Deng, Z.H., Liao, J., Zhang, J.Y., Hao, X.H., Liang, C., Wang, L.H., Xue, H. Zhang, K. and Yan, G.T. 2013. Localized leptin release may be important mechanism of curcumin action after acute ischemic injuries. *The Journal of Trauma and Acute Surgery* 74: 1044-1051.
- Deraniyagala, S.A., Ratnasooriya, W.D. and Goonasekara, C.L. 2003. Antinociceptive effect and toxicological study of the aqueous bark extract of *Barringtonia racemosa* on rats. *Journal of Ethnopharmacology* 86: 21-26.
- Devasagayan, T.P.A, Tilak, J.C., Bloor, K.K., Ketaki, S.S., Saroj S.G. and Lele, R.D. 2004. Free radicals and antioxidants in human health: Current status and future prospects. *Journal of Association Physicians India* 52: 794-804.

- Dimitrios, B. 2006. Sources of natural phenolic antioxidants. *Trends in Food Science and Technology* 17: 505-512.
- Diretto, G., Welsch, R., Tavazza, R., Mourgues, F., Pizzichini, D., Beyer, P. and Giuliano, G. 2007. Silencing of β -carotene hydroxylase increases total carotenoid and β -carotene levels in potato tubers. *Biomed Central Plant Biology* 7: 11-19.
- Dixit, V., Pandey, V. and Shyam, R. 2001. Differential antioxidative responses to cadmium in roots and leaves of pea (*Pisum sativum* L.). *The Journal of Experimental Botany* 52: 1101-1109.
- Drake, J., Kanski, J., Varadarajan, S., Tsoras, M. and Butterfield, D.A. 2002. Elevation of brain glutathione by gamma-glutamylcysteine ethyl ester protects against peroxynitrite-induced oxidative stress. *Journal of Neuroscience Research* 68: 776-784.
- Ebrahimian, E. and Bybordi, A. 2012. Effect of salinity, salicylic acid, silicium and ascorbic acid on lipid peroxidation, antioxidant enzyme activity and fatty acid content of sunflower. *African Journal of Agricultural Research* 7: 3685-3694.
- Edeoga, H.O., Okwu, O.E. and Mbaebie, B.O. 2005. Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology* 4: 685-688.
- Effendy, N.M., Mohamed, N., Muhammad, N., Mohamad, I.N. and Shuid, A.N. 2012. *Eurycoma longifolia*: Medicinal plant in the prevention and treatment of male osteoporosis due to Androgen deficiency. *Evidence-based Complementary and Alternative Medicine* 12: 1-9.
- Egea, P. Sánchez-Bel, Romojaro, F. and Pretel, M.T. 2010. Six edible wild fruits as potential antioxidant additives or nutritional supplements. *Plant Foods for Human Nutrition* 65: 121-129.
- El-Safory, N.S., Fazary, A.E. and Lee, C.K. 2010. Hyaluronidase, a group of glycosidases: Current and future perspectives. *Carbohydrate Polymers* 81: 165-181.
- Erlund, I. 2004. Review of the flavonoids quercetin, hesperidin, and naringenin: Dietary sources, bioactivities, bioavailability and epidemiology. *Nutrition Research* 24: 851-874.
- Evans, J.A. and Johnson, E.J. 2010. The role of phytonutrients in skin health. *Nutrients* 2: 903-928.

- Facundo, V.A., Silveira, A.S.P. and Morais, S.M. 2005. Constituents of *Piper alatabacum* Trel and *Yuncker* (Piperaceae). *Biochemical Systematics and Ecology* 33: 753-756.
- Farre, G., Sanahuja, G., Naqvi, S., Bai, C., Capell, T., Zhu, C. and Christou, P. 2010. Travel advice on the road to carotenoids in plants. *Plant Science* 179: 28-48.
- Foyer, C.H. and Noctor, G. 2003. Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Plant Physiology* 119: 355-364.
- Franchi, J., Coutadeur, M.C., Marteau, C., Mersel, M. and Kupferberg, A. 2000. Depigmenting effects of calcium D-pantetheine-S-sulfonate on human melanocytes. *Pigment Cell Research* 13: 165-171.
- Frost, G.I. 2007. Recombinant human hyaluronidase (rHuPH20): An enabling platform for subcutaneous drug and fluid administration. *Expert Opinion on Drug Delivery* 4:427-40.
- Fulop, T., Khalil, A. and Larbi, A. 2012. The role of elastin peptides in modulating the immune response in aging and age-related diseases. *Pathology Biology* 60:28-33.
- Gachkar, L., Yadegari, D., Rezaei, M.B., Taghizadeh, M., Astaneh, S.A. and Rasooli, I. 2007. Chemical and biological characteristics of *Cuminum cyminum* and *Rosmarinus officinalis* essential oils. *Food Chemistry* 102: 898-904.
- Gang, C. and Zora, D. 2001. Carotenoids are degraded by free radicals but do not affect lipid peroxidation in unilamellar liposomes under different oxygen tensions. *Federation of European Biochemical Societies Letters* 505: 151-154.
- Ganguly, S., Mula, S., Chattopadhyay, S. and Chatterjee, M. 2007. An ethanol extract of *Piper betle* Linn. mediates its anti-inflammatory activity via down-regulation of nitric oxide. *Journal of Pharmacy Pharmacology* 59: 711-718.
- Gara, L., Paciolla, C., Tullio, M.C., Motto, M. and Arrigoni, O. 2000. Ascorbate dependent hydrogen peroxide detoxification and ascorbate regeneration during germination of a highly productive maize hybrid: evidence of an improved detoxification mechanism against reactive oxygen species. *Journal of Plant Physiology* 109: 7-13.

- Gasowska, B., Kafarski, P., Wojtasek, H. 2004. Interaction of mushroom tyrosinase with aromatic amines, o-diamines and o-aminophenols. *Biochimica et Biophysica Acta* 1673: 170-177.
- Georgé, S., Brat, P., Alter, P. and Amiot M.J. 2005. Rapid determination of polyphenols and vitamin C in plant-derived products. *Journal of Agriculture and Food Chemistry* 53: 1370-1373.
- Gherraf, N., Ladjel, S., Labed, B. and Samir, H. 2011. Evaluation of antioxidant potential of various extracts of *Traganum nudatum*. *Plant Sciences Feed* 1: 155-159.
- Ghomi, J.S., Ghadami, M. and Batooli, H. 2012. Bioactivity of *Eucalyptus oleosa* var. obtuse leaves and flowers. *Digest Journal of Nanomaterials and Biostructures* 7: 657-660.
- Gil, M. Tomás-Barberán, A.T. Hess-Pierce, B. and Kader, A.A. 2002. Antioxidant capacities, phenolic compounds, carotenoids and vitamin C content of nectarine and plum cultivars from California. *Journal of Agriculture Food Chemistry* 50: 4976-4982.
- Girish, K. and Kemparaju, K. 2005. Inhibition of *Naja naja* venom hyaluronidase by plant-derived bioactive components and polysaccharides. *Biochemistry* 70: 1145-1150.
- Gomez, L.D., Noctor, G., Knight M.R. and Foyer, C.H. 2004. Regulation of calcium signalling and gene expression by glutathione. *Journal of Experimental Botany* 55: 1851-1859.
- Gómez, L.G., Rubio-Moraga, A. and Ahrazem, O. 2010. Understanding carotenoid metabolism in Saffron stigmas: Unravelling aroma and colour formation. *Functional Plant Science and Biotechnology* 67: 56-61.
- Gong, Y., Toivonen, P.M.A, Lau, O.L and Wiersma, P.A. 2001. Antioxidant system level in 'Braeburn' apple is related to its browning disorder. *Botany Bulletin of Academia Sinica* 42: 259-264.
- Goodarzi, M.T., Zal, F., Malakooti, M., Safari, M.R., Sadeghian, S., 2006. Inhibitory activity of flavonoids on the lens aldose reductase of healthy and diabetic rats. *Acta Medica Iranica* 44: 34-45.
- Gosline, J., Lillie, M., Carrington, E., Guerette, P., Ortlepp, C. and Savage, K. 2002. Elastic Proteins: Biological roles and mechanical properties. *Philosophical Transactions of the Royal Society London Series B Biological Sciences* 357: 121-132.

- Garcia-Alonso M., de Pascual-Teresa, S., Santos-Buelga, C. and Rivas-Gonzalo, J.C. 2004. Evaluation of the antioxidant properties of fruits. *Food Chemistry* 8: 13-18.
- Graf, B.A., Milbury, P.E. and Blumberg, J.B. 2005. Flavonols, flavones, flavanones, and human health: Epidemiological evidence. *Journal of Medicinal Food* 8: 281-290.
- Gratão, P.L., Monteiro, C.C., Peres, L.E.P. and Azevedo, R.A. 2008. The Isolation of antioxidant enzymes from mature tomato (cv. Micro-Tom) plants. *Horticulture Science* 43: 1608-1610.
- Griffith, O.W. and Meister, A. 1979. Glutathione: Interorgan translocation, turnover and metabolism. *Proceedings of the National Academy of Sciences* 76: 5606-5610.
- Gurib-Fakim, A. 2006. Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine* 27: 1-93.
- Habsah, M., Amran, M., Mackeen, M.M., Lajis, N.H., Kikuzaki, H. and Nakatani, N. 2000. Screening of Zingiberaceae extracts for antimicrobial and antioxidant activities. *Journal of Ethnopharmacology* 72: 403-410.
- Hancock, R.D., McRae, D., Haupt, S. and Viola, R. 2003. Synthesis of L-ascorbic acid in the phloem. Research article. *Biomed Central Plant Biology* 3: 1-13.
- Harborne, J.B. and Williams, C.A. 2000. Advances in flavonoid research since 1992. *Phytochemistry* 55: 481-504.
- Harbone, J.B. 1973. *Phytochemical Methods: A guide to modern techniques of plant analysis*, pp. 33-80. London: Chapman and Hall.
- Harrison, E.H. 2005. Mechanisms of digestion and absorption of dietary vitamin A. *Annual Review of Nutrition* 25: 7-103.
- Havsteen, B.H. 2002. The biochemistry and medical significance of the flavonoids. *Pharmacology and Therapeutics* 96: 67-202.
- Hearing, V.J. 2005. Biogenesis of pigment granules: a sensitive way to regulate melanocyte function. *Journal of Dermatological Science* 37: 3-14.
- Hernández, I., Alegre, L., Van Breusegem, F. and Munné-Bosch S. 2009. How relevant are flavonoids as antioxidants in plants. *Trends in Plant Science* 14: 125-132.

- Hertel, W., Peschel, G., Ozegowski, J.H. and Muller, P.J., 2006. Inhibitory effects of triterpenes and flavonoids on the enzymatic activity of hyaluronic acid splitting enzymes. *Archives of Pharmaceutical Chemistry and Life Sciences* 339: 313-318.
- Hiemstra, P.S., Eisenhauer, P.B., Harwig, S.S. and Barselaar, M.T. 2002. Antimicrobial proteins of murine macrophages. *Infection and Immunity* 61:3038-3046
- Hiraga, S., Sasaki, K., Ito, H., Ohashi, Y. and Matsui, H. 2001. A large family class III plant peroxidases. *Plant and Cell Physiology* 42: 462-468.
- Huang, H.C., Liaw, C.C., Zhang, L.J., Ho, H.H., Yang-Kuo, L.M., Shen, Y.C. and Kuo, Y.H. 2008a. Triterpenoidal saponins from *Hydrocotyle sibthorpioides*. *Phytochemistry* 69: 1603-1697.
- Huang, L., Yagura, T. and Chen, S. 2008b. Sedative activity of hexane extract of *Kaempferia galanga* L. and its active compounds. *Journal of Ethnopharmacology* 120: 123-125.
- Huda-Faujan, N., Noriham, A., Norrakiah, A.S. and Babji, A.S. 2007. Antioxidative activities of water extracts of some Malaysian herbs. *ASEAN Food Journal* 14: 61-68.
- Huda-Faujan, N., Noriham, A., Norrakiah, A.S. and Babji, A.S. 2009. Antioxidant activity of plant methanolic extracts containing phenolic compounds. *African Journal of Biotechnology* 8: 484-489.
- Hussain, K., Ismail, Z., Sadikun, A. and Ibrahim, P. 2009. Antioxidant, anti-TB activities, phenolic and amide contents of standardised extracts of *Piper sarmentosum* Roxb. *Natural Product Research* 23: 238-249.
- Hsu, M.F. and Chiang, B.H. 2009. Stimulating effects of *Bacillus subtilis* natto-fermented *Radix astragali* on hyaluronic acid production in human skin cells. *Journal of Ethnopharmacology* 125:474-481.
- Isabelle, M., Lee, B.L., Lim, M.T., Koh, W.P., Huang, D. and Ong, C.N. 2010. Antioxidant activity and profiles of common vegetables in Singapore. *Food Chemistry* 120: 993-1003.
- Ishii, T., Matsuura, H., Kaya, K. and Vairappan, C.S. 2011. A new bisabolane-type sesquiterpenoid from *Curcuma domestica*. *Biochemical Systematics and Ecology* 39: 864-867.

- Ishita, C., Kaushik, B., Uday, B. and Ranjit, K.B. 2004. Turmeric and curcumin: Biological actions and medicinal applications. *Current Science* 87: 44-53.
- Jayaprakasha, G.K., Jagan, L., Mohan, R. and Sakariah, K.K. 2005. Chemistry and biological activities of *Curcuma longa*. *Trends in Food Science and Technology* 16: 533-548.
- Jayashree, G., Kurup Muraleedhara, G., Sudarshl, S. and Jacob, V.B. 2003. Antioxidant activity of *Centella asiatica* on lymphoma-bearing mice. *Fitoterapia* 74: 431-434.
- Jimenez-Cervantes, C., Martinez-Esparza, M., Perez, C., Daum, N., Solano, F. and Garcia-Borron, J.C. 2001. Inhibition of melanogenesis in response to oxidative stress: Transient down regulation of melanocyte differentiation markers and possible involvement of microphthalmia transcription factor. *Journal of Cell Science* 114: 2335-2344.
- Jones, K., Hughes, J., Hong, M., Jia, Q. and Orndorff, S. 2002. Modulation of melanogenesis by aloesin: A competitive inhibitor of tyrosinase. *Pigment Cell Research* 15: 335-340.
- Jordan, A.M., Khan, T.H., Malkin, H., Osborn, H.M., Photiou, A. and Riley, P.A. 2001. Melanocyte-directed enzyme prodrug therapy (MDEPT): Development of second generation prodrugs for targeted treatment of malignant melanoma. *Bioorganic and Medicinal Chemistry* 9: 1549-1558.
- Kamkaen, N., Mulsri, N. and Treesak, C. 2007. Screening of some tropical vegetables for anti-tyrosinase activity. *Thai Pharmaceutical and Health Science Journal* 2: 15-19.
- Kanjanapothi, D., Panthong, A., Lertprasertsuke, N., Taesotikul, T., Rujjanawate, C., Kaewpinit, D., Sudthayakorn, R. and Choochote, W. 2004. Toxicity of crude rhizome extract of *Kaempferia galanga* L. (Proh Hom). *Journal of Ethnopharmacology* 90: 359-365.
- Kapoor, I.P.S., Singh, B., Singh, G., De Heluani, C.S., De Lampasona, M.P. and Catalan, C.A.N. 2009. Chemistry and in vitro antioxidant activity of volatile oil and oleoresins of black pepper (*Piper nigrum*). *Journal of Agricultural and Food Chemistry* 57: 5358-5364.
- Karthikeyan, M., Jayakumar, V., Radhika, K., Bhaskaran, R., Velazhahan, R. and Alice, D. 2005. Induction of resistance in host against the infection of leaf blight pathogen (*Alternaria palandui*) in onion (*Allium cepa* var aggregatum). *Indian Journal of Biochemistry and Biophysics* 42: 371-377.

- Katalinic, V., Kulisic, T., Milos, M. and Jukic, M. 2006. Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food Chemistry* 94: 550-557.
- Kaur, C. and Kapoor, H.C. 2002. Antioxidant activity and total phenolic content of some Asian vegetables. *International Journal of Food Science and Technology* 37: 153-161.
- Khan, M.R. and Omoloso, A.D. 2002. Antibacterial, antifungal activities of *Barringtonia asiatica*. *Fitoterapia* 73: 255-260.
- Khan, S., Jabbar, A., Hassan, C.M. and Rahid, M.A. 2001. Antibacterial activity of *Barringtonia racemosa*. *Fitoterapia* 72: 162-164.
- Kiken, D.A. and Cohen, D.E. 2002. Contact dermatitis to botanical extracts. *American Journal of Contact Dermatitis* 13: 148-152.
- Kikuzaki, H. and Nakatani, N. 1993. Antioxidant effects of some ginger constituents. *Journal of Food Science* 58: 1407-1410.
- Kim, D.S., Kim, S.Y., Chung, J.H., Kim, K.H., Eun, H.C. and Park, K.C. 2002. Delayed ERK activation by ceramide reduces melanin synthesis in human melanocytes. *Cellular Signalling* 14: 779-785.
- Kim, J.H., Byun, J.C., Bandi, A.K.R., Hyun, C.G. and Lee, N.M. 2009. Compounds with elastase inhibition and free radical scavenging activities from *Callistemon lanceolatus*. *Journal of Medicinal Plants Research* 3: 914-920.
- Kim, K.S., Kim, J.A., Eom, S.Y., Lee, S.H., Min, K.R. and Kim, Y. 2006a. Inhibitory effect of *Piper longuminine* on melanin production in melanoma B16 cell line by down regulation of tyrosinase expression. *Pigment Cell Research* 19: 90-98.
- Kim, M., Meng, X. and Kim, A. 2006b. Developing a long lasting tyrosinase inhibitor from *Morus alba* L. *Cosmetic and Toiletries Magazine* 27: 91-102.
- Kim, S.S., Hyun, C.G., Lee, J., Lim, J., Kim, J.Y. and Park, D. 2007. *In vitro* screening of Jeju medicinal plants for cosmeceutical materials. *Journal of Applied Biology Chemistry* 50: 215-220.
- Knight, T.M. and Miller, T.E. 2004. Local adaptation within a population of *Hydrocotyle bonariensis*. *Evolutionary Ecology Research* 6: 103-114.
- Kole, P.L., Jadhav, H.R., Thakurdesai, P. and Nagappa, A.N. 2005. Cosmetic potential of herbal extracts. *Natural Product Radiance* 4: 315-321.

- Koleva, I.I., Van B.T.A., Linszen, J.P.H., Groot, A. and Evstatieva, L.N. 2002. Screening of plant extracts for antioxidant activity: A comparative study on three testing methods. *Phytochemical Analysis* 13: 8-17.
- Korkmaz, B., Moreau, T. and Gauthier, F. 2008. Neutrophil elastase, proteinase 3 and cathepsin G: Physicochemical properties, activity and physiopathological functions. *Biochimie* 90: 227-242.
- Kong, K.W., Mat Junit, S., Aminudin, N., Ismail, A. and Abdul Aziz, A. 2012. Antioxidant activities and polyphenolics from the shoots of *Barringtonia racemosa* (L.) Spreng in a polar to apolar medium system. *Food Chemistry* 134: 324-332.
- Kumaran, A. and Karunakaran, R.J. 2006. Antioxidant and free radical scavenging activity of an aqueous extract of *Coleus aromaticus*. *Food Chemistry* 97: 109-114.
- Kumar, C.M., Sathisha, U.V., Dharmesh, S., Rao, A.G. and Singh, S.A. 2011. Interaction of sesamol (3,4-methylenedioxyphenol) with tyrosinase and its effect on melanin synthesis. *Biochimie* 93: 562-9
- Kumar, S.R. and Vallikannan, B. 2010. Carotenoid composition and retinol equivalent in plants of nutritional and medicinal importance: Efficacy of β -carotene from *Chenopodium album* in retinol-deficient rats. *Food Chemistry* 119: 1584-1590.
- Laura, O.M. 2013. Evaluation of the economic effects of inward foreign direct investment in Costa Rica's economy. *Global Journal of Logistics and business Management* 1: 1-14.
- La Vecchia, C., Altieri, A. and Tavani, A. 2001. Vegetables, fruit antioxidants and cancer: A review of Italian studies. *European Journal of Nutrition* 40: 261-267.
- Lee, K.K. and Choi, J. 1999. The effects of *Areca catechu* L extracts on anti-aging. *International Journal of Cosmetic Science* 21: 285-294.
- Lee, K.K., Cho, J.J., Park, E.J. and Choi, J.D. 2001. Anti-elastase and anti-hyaluronidase of phenolic substance from *Areca catechu* as a new anti-aging agent. *International Journal of Cosmetic Science* 23: 341-346.
- Lee, S.E., Shin, H.T., Hwang, H.J. and Kim, H.J. 2003. Antioxidant activity of extracts from *Alpinia katsumadai* seed. *Phytotherapy Research* 17: 1041-1047.

- Lee, S.H., Choi, S.Y., Kim, H., Hwang, J.S., Lee, B.G., Gao, J.J. and Kim, S.Y. 2002. Mulberroside F isolated from the leaves of *Morus alba* inhibits melanin biosynthesis. *Biological and Pharmaceutical Bulletin* 25: 1045-1048.
- Lee, T.K. and Vairappan, C.S. 2011. Antioxidant, antibacterial and cytotoxic activities of essential oils and ethanol extracts of selected South East Asian herbs. *Journal of Medicinal Plants Research* 5: 5284-5290.
- Lim, H.K., Moon, J.Y., Kim, H., Cho, M. and Somi, K.C. 2009. Induction of apoptosis in U937 human leukaemia cells by the hexane fraction of an extract of immature *Citrus grandis* Osbeck fruits. *Food Chemistry* 114: 1245-1250.
- Lim, Y.Y. and Murtijaya, J. 2007. Antioxidant properties of *Phyllanthus amarus* extracts affected by different drying methods. *Food Science and Technology* 40: 1664-1669.
- Lin, C.B., Babiarz, L., Liebel, F., Price, E.R., Kizoulis, M., Gendimenico, G.J., Fisher, D.E. and Seiberg, M. 2002a. Modulation of microphthalmia-associated transcription factor gene expression alters skin pigmentation. *Journal of Investigative Dermatology* 119: 1330-1340.
- Lin, C.M., Chen, C.S., Liang, Y.C. and Lin, J.K. 2002b. Molecular modeling of flavonoids that inhibits xanthine oxidase. *Biochemical and Biophysical Research Communications* 294: 167-172.
- Liu, F. and Ng, T.B. 2000. Antioxidative and free radical scavenging activities of selected medicinal herbs. *Life Sciences* 66: 725-735.
- Liu, N., Yin, M.R., Sun, G. and Wong, P. 2005. Genetic manipulation of carotenoid biosynthesis. *Journal of Biochemistry* 116: 980-985.
- Lisiewska, Z., Kmiecik, W. and Korus, A. 2006. Content of vitamin C, carotenoids, chlorophylls and polyphenols in green parts of dill (*Anethum graveolens* L.) depending on plant height. *Journal of Food Composition and Analysis* 19: 134-140.
- Maity, N., Nema, N.K., Abedy, S., Sarkar, B.K. and Mukherjee, P.K. 2011. Exploring *Tagetes erecta* Linn flower for the elastase, hyaluronidase and MMP-1 inhibitory activity. *Journal of Ethnopharmacology* 137:1300-1305.
- Mallavadhani, U.V., Sudhakar, A.V.S., Sathyanarayana, K.V.S., Mahapatra, A., Li, W. and Vanbreeman, R.B. 2006. Chemical and analytical screening of some edible mushrooms. *Food Chemistry* 95: 58-64.

- Maresca, V., Flori, E., Cardinali, G., Briganti, S., Lombardi, D. and Mileo, A.M. 2006. Ferritin light chain down-modulation generates depigmentation in human metastatic melanoma cells by influencing tyrosinase maturation. *Journal of Cellular Physiology* 206: 843-848.
- Mariod, A.A., Mirghani, M.E.S., Abdul, A.B. and Abdelwahab, S.I. 2009. *Detarium microcarpum* Guill and Perr fruit proximate chemical analysis and sensory characteristics of concentrated juice and jam. *African Journal of Biotechnology* 8: 4217-4221.
- Masuda, M., Murata, K., Fukuhama, A., Naruto, S., Fujita, T., Uwaya, A., Isami, F. and Matsuda, H. 2009. Inhibitory effects of constituents of *Morinda citrifolia* seeds on elastase and tyrosinase. *Journal of Natural Medicines* 63: 267-273.
- Matsuda, H., Morikawa, T., Ueda, H. and Yoshikawa, M. 2001. Saponin constituents Gotu Kola 2: Structures of new ursane- and oleanane-type triterpene oligoglycosides, centellasaponin B, C and D, from *Centella asiatica* cultivated in Sri Lanka. *Chemical and Pharmaceutical Bulletin* 49: 1368-1371.
- Matsushita, A., Sasaki, Y., Warashina, T., Miyase, T., Noguchi, H. and Velde, D.V. 2004. Hydrocotylosides I-VII, new oleanane saponins from *Hydrocotyle sibthorpioides*. *Journal of Natural Product* 67: 384-388.
- Middleton, E., Kandaswami, C. and Theoharides, T.C. 2000. The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer. *Pharmacology Reviews* 52: 673-751.
- Miyazawa, M., Oshima, T., Koshio, K., Itsuzaki, Y. and Anzai, J. 2003. Tyrosinase inhibitor from black rice bran. *Journal of Agricultural and Food Chemistry* 51: 6953-6956.
- Mridula, K.R. and Jayachandran, B.K. 2001. Quality of mango-ginger (*Curcuma amada* roxb.) as influenced by mineral nutrition. *Journal of Tropical Agriculture* 39: 182-183.
- Müller, L., Fröhlich, K. and Böhm, V. 2011. Comparative antioxidant activities of carotenoids measured by ferric reducing antioxidant power (FRAP), ABTS bleaching assay (α TEAC), DPPH assay and peroxy radical scavenging assay. *Food Chemistry*, 129: 139-148.
- Nagarathnam, R., Rengasamy, A. and Balasubramanian, R. 2010. Purification and properties of cysteine protease from rhizomes of *Curcuma longa* (Linn.). *Journal of the Science of Food and Agriculture* 90: 97-105.

- Nakano, Y. and Asada, K. 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and Cell Physiology* 22: 867-880.
- Navickiene, H.M.D., Alécio, A.C., Kato, M.J., Bolzani, V.S., Young, M.C.M., Cavaleiro, A.J. and Furlan, M. 2000. Antifungal amides from *Piper hispidum* and *Piper tuberculatum*. *Phytochemistry* 55: 621-626.
- Nayyar, H. and Gupta, D. 2006. Differential sensitivity of C₃ and C₄ plants to water deficit stress: Association with oxidative stress and antioxidants. *Environmental and Experimental Botany* 58: 106-113.
- Necas, J., Bartosikova, L., Brauner, P. and Kolar, J. 2008. Hyaluronic acid (hyaluronan). *Veterinarni Medicina* 53: 397-411.
- Nema, N.K., Maity, N., Sarkar, B. and Mukherjee, P.K. 2011. *Cucumis sativus* fruit potential antioxidant, anti hyaluronidase and anti-elastase agent. *Archives of Dermatological Research* 303: 247-252.
- Noctor, G., Veljovic-Jovanovic, S., Driscoll, S., Novitskaya, L. and Foyer, C.H. 2002. Drought and oxidative load in the leaves of C₃ plants: A predominant role for photorespiration. *Annals of Botany* 89: 841-850.
- No, J.K., Kim, M.S., Kim, Y.J., Bae, S.J., Choi, J.S. and Chung, H.Y. 2004. Inhibition of tyrosinase by protocatechuic aldehyde. *American Journal of Chinese Medicine* 32: 97-103.
- Norhaiza, M., Maziah, M. and Hakimian, M. 2009. Antioxidative properties of leaf of a popular herb, *Labisia pumila*. *Journal of Medicinal Plant Research* 3: 217-223.
- Nosanchuk, J.D. and Casadevall, A. 2003. The contribution of melanin to microbial pathogenesis. *Journal of Cell Microbiology* 5: 203-223.
- Olorunnisola, O.S., Bradley, G. and Afolayan, A.J. 2012. Antioxidant activity of acetone and ethanolic leaves extracts of *Hippobromus pauciflorus* (L.f.) Radlk. *African Journal of Biotechnology* 11: 1206-1213.
- Olsen, D.L., Raub, W., Bradley, C., Johnson, M., Macias, J.L., Love, V. and Markoe, A. 2001. The effect of *Aloe vera* gel/mild soap versus mild soap alone in preventing skin reactions in patients undergoing radiation therapy. *Oncology Nursing Forum* 28: 543-547.

- Oonsivilai, R., Ferruzzi, M.G. and Ningsanond, S. 2008. Antioxidant activity and cytotoxicity of Rang Chuet (*Thunbergia laurifolia* Lindl.) extracts. *Asian Journal of Food and Agro-Industry* 1: 116-128.
- Othman, R., Ibrahim, H., Mohd, M.A., Mustafa, M.R. and Awang, K. 2006. Bioassay guided isolation of a vasorelaxant active compound from *Kaempferia galangal* L. *Phytomedicine* 13: 61-66.
- Paola, M., Braca, A., Pizza, C. and Tomassi, N.D. 2005. Structure antioxidant activity relationships of flavonoids isolated from different plant species. *Food Chemistry* 92: 349-355.
- Park, S.H., Kim, D.S., Kim, W.G., Ryoo, I.J., Lee, D.H. and Huh, C.H. 2004. Terrein: A new melanogenesis inhibitor and its mechanism. *Cellular and Molecular Life Sciences* 61: 2878-2885.
- Parr, A.J and Bolwell, G.P. 2000. Phenols in the plant and in man. The potential for possible nutritional enhancement of the diet by modifying the phenols content or profile. *Journal of the Science of Food and Agriculture* 80: 985-1012.
- Peng, Z.F., Strack, D., Baumert, A., Subramaniam, R., Goh, N.K., Chia, T.F., Tan, S.N. and Chia, L.S. 2003. Antioxidant flavonoids from leaves of *Polygonum hydropiper* L. *Phytochemistry* 62: 219-228.
- Peschel, W., Sánchez-Rabaneda, F., Diekmann, W., Plescher, A., Gartzía, I. and Jiménez, D. 2006. An industrial approach in the search of natural antioxidants from vegetable and fruit wastes. *Food Chemistry* 97: 137-150.
- Piwowarski, J.P., Kiss, A.K. and Wojciechowska, M.K. 2011. Anti hyaluronidase and anti-elastase activity screening of tannin-rich plant materials used in traditional Polish medicine for external treatment of diseases with inflammatory background. *Journal of Ethnopharmacology* 137: 937-941.
- Prakash, K., Prajapati, S., Ahmad, A., Jain, S.K. and Bhakuni, V. 2002. Unique oligomeric intermediates of bovine liver catalase. *Protein Science* 11: 46-57.
- Qader, S.W., Abdulla, M.A., Chua, L.S., Najim, N., Zain, M.M. and Hamdan, S. 2012. A review: Potential bioactive property of *Polygonum minus* Huds (kesum). *Scientific Research and Essay* 7: 90-93.
- Qader, S.W., Abdulla, M.A., Chua, L.S., Najim, N., Zain, M.M. and Hamdan, S. 2011. Antioxidant, total phenolic content and cytotoxicity evaluation of selected Malaysian plants. *Molecules* 16: 3433-3443.

- Raju, M., Varakumar, S., Lakshminarayana, R., Krishnakantha, T. P. and Baskaran, V. 2007. Carotenoid composition and vitamin A activity of medicinally important green leafy vegetables. *Food Chemistry* 101: 1598-1605.
- Rayjalakshmi, P., Venkatalaxmi, K., Venkatalakshamma, K., Jyothsna, Y., Balachandramanidevi, K. and Suneetha, V. 2001. Total carotenoid and β -carotene contents of forest green leafy vegetables consumed by tribals of South India. *Plant Foods for Human Nutrition* 56: 225-238.
- Rao, L.G., Mackinnon, E.S., Josse, R.G., Murray, T.M., Strauss, A. and Rao, A.V. 2007. Lycopene consumption decreases oxidative stress and bone resorption markers in postmenopausal women. *Osteoporosis International* 18: 109-115.
- Rao, N.V. and Kaladhar, D.S.V.G.K. 2014. Antioxidant and antimicrobial activities of rhizome extracts of *Kaempferia galanga*. *World Journal of Pharmacy and Pharmaceutical Sciences* 3: 1180-1189.
- Rasdi, N.H.M., Samah, O.A., Sule, A. and Ahmed, Q.U. 2010. Antimicrobial studies of *Cosmos caudatus* Kunth. (Compositae). *Journal of Medicinal Plants Research* 4: 669-673.
- Rasmy, N.M., Hassan, A.A., Foda, M.I. and El-Moghazy, M.M. 2012. Assessment of the antioxidant activity of sage (*Salvia officinalis* L.) extracts on the shelf life of mayonnaise. *World Journal of Dairy and Food Science* 7: 28-40.
- Rathee, J.S., Patro, B.S., Mula, S., Gamre, S. and Chattopadhyay, S. 2006. Antioxidant activity of *Piper betel* leaf extract and its constituents. *Journal of Agricultural and Food Chemistry* 54: 9046-9054.
- Raychaudhuri, S.S. and Deng, X.W. 2008. The role of superoxide dismutase in combating oxidative stress in higher plants. *The Botanical Review* 66: 89-98.
- Razak, F.A., Othman, R.Y. and Rahim, Z.H.A. 2006. The effect of *Piper betle* and *Psidium guajava* extracts on the cell-surface hydrophobicity of selected early settlers of dental plaque. *Journal of Oral Science* 48:71-75.
- Raven, E.L. 2000. Peroxidase-catalyzed oxidation of ascorbate structural, spectroscopic and mechanistic correlations in ascorbate peroxidase. *Subcellular Biochemistry* 35: 317-349.
- Reddy, A.R., Chiatanya, K.V. and Vivekanandan, M. 2004. Drought induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of Plant Physiology* 161: 1189-1202.

- Rong, X., Peng, G., Suzuki, T., Yang, Q., Yamahara, J. and Li, Y. 2009. A 35-day gavage safety assessment of ginger in rats. *Regular Toxicology and Pharmacology* 54: 118-123.
- Rosati, C., Aquilani, R., Dharmapuri, S., Pallara, P., Marusic, C., Tavazza, R., Bouvier, F., Camara, B. and Giuliano, G. 2000. Metabolic engineering of β -carotene and lycopene content in tomato fruit. *The Plant Journal* 24: 413-419.
- Ross, J.A. and Kasum, C.M. 2002. Dietary flavonoids: Bioavailability metabolic effects, and safety. *Annual Review of Nutrition* 22: 19-34.
- Rosyid, T.A., Karim, R., Mohd Adzahan, N. and Mohd Ghazali, F. 2011. Antibacterial activity of several Malaysian leaves extracts on the spoilage bacteria of yellow alkaline noodles. *African Journal of Medicinal Plants* 5: 898-903.
- Rukachaisirikul, T., Siriwanakit, P., Sukcharoenphol, K., Wongvein, C., Ruttanaweng, P., Wongwattanavuch, P. and Suksamrarn, A. 2004. Chemical constituents and bioactivity of *Piper sarmentosum*. *Journal of Ethnopharmacology* 93: 173-176.
- Sacchetti, G., Maietti, S., Muzzoli, M., Scaglianti, M., Manfredini, S., Radice, M. and Bruni, R. 2005. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chemistry* 91: 621-632.
- Sadhu, S.K., Tamaki, M., Ohtsuki, T., Toume, K., Kovano, T., Kowithayakorn, T. and Ishibashi, M. 2009. Cadinane sesquiterpenes from *Curcuma parviflora*. *Journal of Natural Product* 72: 782-783.
- Saewan, N., Koysomboon, S., Chantrapromma, K. 2011. Anti-tyrosinase and anti-cancer activities of flavonoids from *Blumea balsamifera* DC. *Journal of Medicinal Plants Research* 5: 1018-1025.
- Sahasrabudhe, A. and Deodhar, M. 2010. Anti-hyaluronidase, anti-elastase activity of *Garcinia indica*. *International Journal of Botany* 6: 299-303.
- Saikia, L.R. and Upadhyaya, S. 2011. Antioxidant activity, phenol and flavonoid content of *Asparagus racemosus* Willd. a medicinal plant grown using different organic manures. *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 2: 457-463.

- Samy, J., Sugumaran, M. and Lee, K.L.W. 2005. Herbs of Malaysia: An Introduction to the Medicinal, Culinary, Aromatic and Cosmetic Use of Herbs, pp. 82-83. Kuala Lumpur: Times Editions.
- Sarkar A, Ray D, Shrivastava AN, Sarker S. 2006. Molecular biomarkers: Their significance and application in marine pollution monitoring. *Ecotoxicology and Environmental Safety* 15: 333-340.
- Sarla, S., Prakash, M.A., Apeksha, R. and Subash, C. 2011. Free radical scavenging (DPPH) and Ferric reducing ability (FRAP) of *Aphanamixis polystachya* (Wall) parker. *International Journal of Drug Development and Research* 3: 271-274.
- Seiberg, M., Paine, C., Sharlow, E., Andrade-Gordon, P., Costanzo, M., Eisinger, M. and Shapiro, S.S. 2000. Inhibition of melanosome transfer results in skin lightening. *Journal of Investigative Dermatology* 115: 162-167.
- Seo, S.Y., Sharma, V.K. and Sharma, N. 2003. Mushroom tyrosinase: Recent prospects. *Journal of Agricultural and Food Chemistry* 51: 2837-2853.
- Seyoum, A., Asres, K. and El-Fiky, F.K. 2006. Structure–radical scavenging activity relationships of flavonoids. *Phytochemistry* 67: 2058-2070.
- Shan, B., Yizhong, Z., Sun, C.M. and Corke, H. 2005. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *Journal of Agricultural Food Chemistry* 53: 7749-7759.
- Sharififar, F., Dehghn, N.G. and Mirtajaldini, M. 2009. Major flavonoids with antioxidant activity from *Teucrium polium* L. *Food Chemistry* 112: 885-888.
- Sharma, A., Gautam, S. and Jadhav, S.S. 2000. Spice extracts as dose modifying factors in radiation inactivation of bacteria. *Journal of Agricultural and Food Chemistry* 48: 1340-1344.
- Sharma, R.A., Gescher, A.J. and Steward, W.P. 2005. Curcumin: The story so far. *European Journal of Cancer* 41: 1955-1968.
- Shigeoka, S., Ishikawa, T., Tamoi, M., Migawa, Y., Takeda, T., Yabuta, Y. and Yoshimura, K. 2002. Regulation and function of ascorbate peroxidase isoenzymes. *Journal of Experimental Botany* 53: 1305-1319.
- Shih, C.D., Chen, J.J. and Leet, H.H. 2006. Activation of nitric oxide signaling pathway mediates hypotensive effect of *Muntingia calabura* L. (Tiliaceae) leaf extract. *American Journal of Medicine* 34: 857-872.

- Shimogaki, H., Tanaka, Y., Tamai, H. and Masuda, M. 2000. *In vitro* and *in vivo* evaluation of ellagic acid on melanogenesis inhibition. *International Journal of Cosmetic Science* 22: 291-303.
- Shon, M.Y., Kim, T.H and Sung, N.J. 2003. Antioxidants and free radical scavenging activity of *Phellinus baumii* (*Phellinus* of *Hymenochaetaceae*) extracts. *Food Chemistry* 82: 593-597.
- Shui, G., Leong, L.P. and Wong, S.P. 2005. Rapid screening and characterisation of antioxidants of *Cosmos caudatus* using liquid chromatography coupled with mass spectrometry. *Journal of Chromatography B* 827: 127-138.
- Siedle, B., Hrenn, A. and Merfort, I. 2007. A review on natural compounds as inhibitors of human neutrophil elastase. *Planta Medica* 73:401-420.
- Singh, G., Kawatra, A. and Sehgal, S. 2001. Nutritional composition of selected green leafy vegetables, herbs and carrots. *Plant Foods for Human Nutrition* 56: 359-364.
- Singh, M., Shakya, S., Soni, V.K., Dangi, A., Kumar, N. and Bhattacharya, S.M. 2009. The n-hexane and chloroform fractions of *Piper betle* L., trigger different arms of immune responses in BALB/c mice and exhibit antifilarial activity against human lymphatic filarid *Brugia malayi*. *International Immunopharmacology* 9: 716-728.
- Singleton, V.L. and Rossi, J.A. 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture* 16: 144-158.
- Sinha, S., Saxena, R. and Singh, S. 2005. Chromium induced lipid peroxidation in the plants of *Pistia stratiotes* L. role of antioxidants and antioxidant enzymes. *Journal of Chemosphere* 58: 595-604.
- Smirnoff, N. and Wheeler, G.L. 2000. Ascorbic acid in plants: Biosynthesis and function. *Critical Reviews in Biochemistry and Molecular Biology* 35: 291-314.
- Somogyi-Nelson, M. 1952. Notes on sugar determination. *Journal of Biological Chemistry* 195:19-23.
- Song, F.L., Gan, R.Y., Zhang, Y. Xiao, Q., Kuang, L. and Li, B.L. 2010. Total phenolic contents and antioxidant capacities of selected chinese medicinal plants. *International Journal of Molecular Sciences* 11: 2362-2372.

- Stern, R., Asari, A.A. and Sugahara, K.N. 2006. Hyaluronan fragments: An information- rich system. *European Journal of Cell Biology* 85:699-715.
- Stevens, R.G., Creissen, G.P. and Mullineaux, P.M. 2000a. Characterisation of pea cytosolic glutathione reductase expressed in transgenic tobacco. *Planta* 211: 537-545.
- Stevens, W.D., Ulloa Ulloa, C., Pool, A. and Montiel, O.M. 2000b. In: Flora de Nicaragua, Missouri Botanical Gardens Press: Missouri, USA.
- Subramaniam, V., Adenan, M.I., Ahmad, A.R. and Sahdan, R. 2003. Natural Antioxidants: *Piper sarmentosum* (Kadok) and *Morinda elliptica* (Mengkudu). *Malaysian Journal of Nutrition* 9: 41-51.
- Sugumaran, M. 2002. Comparative biochemistry of Eumelanogenesis and the protective roles of Phenoloxidase and melanin in insects. *Pigment Cell Research* 15: 2-9.
- Sultana, B. and Anwar, F. 2008. Flavonols (kaempferol, quercetin, myricetin) contents of selected fruits, vegetables and medicinal plants. *Food Chemistry* 108: 879-884.
- Sumantran, V.N., Kulkarni, A.A., Harsulkar, A., Wele, A., Koppikar, S.J., Chandwaskar, R., Gaire, V., Dalvi, M. and Wagh, U.V. 2007. Hyaluronidase and collagenase inhibitory activities of the herbal formulation *Triphala guggulu*. *Journal of Bioscience* 32:755-761.
- Sun, J., Chu, Y.F., Wu, X.Z. and Liu, R.H. 2002. Antioxidant and antiproliferative activities of common fruits. *Journal of Agricultural and Food Chemistry* 50: 7449-7454.
- Sun, T. and Ho, C. 2005. Antioxidant activities of buckwheat extracts. *Food Chemistry* 90: 743-749.
- Tabopda, T.K., Mitaine-Offer, A., Miyamoto, T., Ngadjui Tchaleu, B. and Lacaille-Dubois, M. 2012. Triterpenoid saponins from *Hydrocotyle bonariensis*. *Phytochemistry* 73:142-147.
- Tang, G. 2010. Bioconversion of dietary provitamin A carotenoids to vitamin A in humans. *The American Journal of Clinical Nutrition* 91: 1468-1473.
- Tangkanakul, P., Auttaviboonkul, P., Niyomwit, B., Lowvitoon, N., Charoenthamawat, P. and Trakoontivakorn, G. 2009. Antioxidant capacity, total phenolic content and nutritional composition of Asian foods after thermal processing. *International Food Research Journal* 16: 571-580.

- Tewtrakul, S. and Subhadhirasakul, S. 2007. Anti-allergic activity of some selected plants in the Zingiberaceae family. *Journal of Ethnopharmacology* 109: 535-538.
- Tewtrakul, S., Yuenyongsawad, S., Kummee, S. and Atsawajaruwan, L. 2005. Chemical components and biological activities of volatile oil of *Kaempferia galanga* Linn. Songklanakarin. *Journal of Science and Technology* 27: 503-507.
- Thomas, T.J., Panikkar, B., Subramoniam, A., Nair, M.K. and Panikkar, K.R. 2002. Antitumour property and toxicity of *Barringtonia racemosa* Roxb seed extract in mice. *Journal of Ethnopharmacology* 82: 223-227.
- Thring, T.S., Hili, P. and Naughton, D.P. 2009. Anti-collagenase, anti-elastase and anti-oxidant activities of extracts from 21 plants. *Biomed Central Complementary and Alternative Medicine* 9: 27-38.
- Tilak, J.C., Banerjee, M., Mohan, H. and Devasagayam, T.P.A. 2004. Antioxidant availability of turmeric in relation to its medicinal and culinary use. *Phytotherapy Research* 18: 798-804.
- Tuntiwachwuttikul, P. Phansa, P., Pootaeng-On, Y. and Taylor, W.C. 2006. Chemical constituents of the roots of *Piper sarmentosum*. *Chemical and Pharmaceutical Bulletin*, 54:149-151.
- Tuncagil, S., Odaci, D., Yildiz, E., Timur, S. and Toppare, L. 2009. Design of a microbial sensor using conducting polymer of 4-(2,5-di(thiophen-2-yl)-1H-pyrrole-1-yl) benzenamine. *Sensors and Actuators B* 137: 42-47.
- Valko, M., Rhodes, C.J., Moncol, J., Izakovic, M. and Mazur, M. 2006. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-Biological Interactions* 160: 1-40.
- Vanka, A., Tandon, S., Rao, S.R., Udupa, N. and Ramkumar, P. 2001. The effect of indigenous Neem, *Azadirachta indica* mouthwash on *Streptococcus* mutants and lactobacilli growth. *Indian Journal of Dental Research* 12: 133-144.
- Velioglu, Y.S., Mazza, G., Gao, L. and Oomah, B.D. 1998. Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *Journal of Agricultural and Food Chemistry* 46: 4113-4117.
- Vimala, S., Adenan, M.I., Ahmad, A.R. and Shahdan, R. 2003. Nature's choice to wellness: Antioxidant vegetables/ulam. Forest Research Institute Malaysia: Selangor.

- Vozenin-Brotons, M.C., Sivan, C., Gault, N., Renard, C., Geffrotin, C., Delanian, S., Lefaix, J.L. and Martin, M. 2001. Antifibrotic action of Cu/Zn sod is mediated by tgf-beta 1 repression and phenotypic reversion of myofibroblasts. *Free Radical Biology and Medicine* 30: 30-42.
- Wang, K.H., Lin, R.D., Hsu, F.L., Huang, Y.H., Chang, H.C., Huang, C.Y. and Lee, M.H. 2006. Cosmetic applications of selected traditional Chinese herbal medicines. *Journal of Ethnopharmacology* 106: 353-359.
- Wang, L.Y., Zhang, M., Zhang, C.F. and Wang, Z.T. 2008. Alkaloid and sesquiterpenes from the root tuber of *Curcuma longa*. *Biochemical Systematics and Ecology* 36: 476-481.
- Wang, S.Y. and Ballington, J.R. 2007. Free radical scavenging capacity and antioxidant enzyme in deerberry (*Vaccinium stamineum* L.). *Journal of Food Science and Technology* 40: 1352-1361.
- Wasman, S.Q., Mahmood, A.A., Salehuddin, H., Zahra, A.A and Salmah, I. 2010. Cytoprotective activities of *Polygonum minus* aqueous leaf extract on ethanol-induced gastric ulcer in rats. *Journal of Medicinal Plants Research* 4: 2658-2665.
- Waugh, W.H. 2003. Simplified method to assay total plasma peroxidase activity and ferriheme products in sickle cell anemia, with intial results in assessing clinical severity in a trial with citruline therapy. *Journal of Pediatric Hematology and Oncology* 25: 831-834.
- Wijeweera, P., Arnason, J.T., Koszycki, D. and Merali, Z. 2006. Evaluation of anxiolytic properties of Gotukola (*Centella asiatica*) extracts and asiaticoside in rat behavioral models. *Phytomedicine* 13: 668-676.
- Wirotasangthong, M., Inagaki, N., Tanaka, H., Thanakijcharoenpath, W. and Nagai, H. 2008. Inhibitory effects of *Piper betle* on production of allergic mediators by bone marrow-derived mast cells and lung epithelial cells. *International Immunopharmacology* 8: 453-457.
- Wong, C.C., Li, H.B., Cheng, K.W. and Chen, F. 2006. A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay. *Food Chemistry* 97: 705-711.
- Wu, C.Q., Chen, F., Wang, X., Kim, H.J., He, G.Q., Haley-Zitlin, V. and Huang, G. 2006. Antioxidant constituents in feverfew (*Tanacetum parthenium*) extract and their chromatographic quantification. *Food Chemistry* 96: 220-227.

- Yan, Q., Cao, R., Yi, W., Yu, L., Chen, Z., Ma, L. and Song, H. 2009. Synthesis and evaluation of 5-benzylidene (thio) barbiturate-beta-D-glycosides as mushroom tyrosinase inhibitors. *Bioorganic and Medicinal Chemistry Letters* 19:4055-4058.
- Yannarelli, G.G., Fernández-Alvarez, A.J., Santa-Cruz, D.M and Tomaro, M.L. 2007. Glutathione reductase activity and isoforms in leaves and roots of wheat plants subjected to cadmium stress. *Phytochemistry* 68: 505-512.
- Yildirim, A., Oktay, M. and Bıllaloğlu, V. 2001. The antioxidant activity of the leaves of *Cydonia vulgaris*. *Turkey Journal of Medical Sciences* 31: 23-27.
- Zaeoung, S., Plubrukarn, A. and Keawpradub, N. 2005. Cytotoxic and free radical scavenging activities of Zingiberaceous rhizomes. *Songklanakarin Journal of Science and Technology* 27: 799-812.
- Zaidan, M.R.S., Noor Rain, A., Badrul, A.R., Adlin, A., Norazah, A. and Zakiah, I. 2005. *In vitro* screening of five local medicinal plants for antibacterial activity using disc diffusion method. *Tropical Biomedicine* 22: 165-170.
- Zainol, M.K., Abdul-Hamid, A., Yusof, S. and Muse, R. 2003. Antioxidative activity and total phenolic compounds of leaf, root and petiole of four accessions of *Centella asiatica* (L.) urban. *Food Chemistry* 81: 575-581.
- Zainol, N.A., Voo, S.C., Sarmidi, M.R. and Aziz, R.A. 2008. Profiling of *Centella asiatica* (L.) Urban extract. *The Malaysian Journal of Analytical Sciences* 12: 322-327.
- Zárate, R., Dirk, C., Van der Heijden, R. and Verpoorte, R. 2001. Terpenoid indole alkaloid profile changes in *Catharanthus pusillus* during development. *Plant Science* 160: 971-977.
- Zeng, Y.C., Qiu, F., Takahashi, K., Liang, J.M., Qu, G.X. and Yao, X.S. 2007. New sesquiterpenes and calebin derivatives from *Curcuma longa*. *Chemical and Pharmaceutical Bulletin* 55: 940-943.
- Zheng, W. and Wang, S.Y. 2001. Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agricultural and Food Chemistry* 49: 5165-5170.
- Zheng, N., Wang, Z., Chen, F. and Lin, J. 2011. Evaluation to the antioxidant activity of total flavonoids extract from *Syzygium jambos* seeds and optimization by response surface methodology. *African Journal of Pharmacy and Pharmacology* 5: 2411-2419.

- Zishen, J., Mengcheng, T. and Jianming, W. 1999. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry* 64: 555-559.
- Zuurbier, C.J., Eerbeek, O., Goedhart, P.T., Struys, E.A., Verhoeven, N.M., Jakobs, C. and Ince, C. 2004. Inhibition of the pentose phosphate pathway decreases ischemia- reperfusion-induced creatine kinase release in the heart. *Cardiovascular Research* 62: 145-153.
- Zverlov, V.V., Schantz, N. Schmitt-Kopplin, P. and Schwarz, W.H. 2005. Two new major subunits in the cellulosome of *Clostridium thermocellum*: Xyloglucanase Xgh74A and xylanase Xyn10D. *Microbiology* 151: 3395-3401.