

University of Groningen

Unravelling the stellar Initial Mass Function of early-type galaxies with hierarchical Bayesian modelling

Dries, Matthijs

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Dries, M. (2018). Unravelling the stellar Initial Mass Function of early-type galaxies with hierarchical Bayesian modelling [Groningen]: Rijksuniversiteit Groningen

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



**rijksuniversiteit
groningen**

Unravelling the stellar Initial Mass Function of Early-Type Galaxies with hierarchical Bayesian modelling

Proefschrift

ter verkrijging van de graad van doctor aan de
Rijksuniversiteit Groningen
op gezag van de
rector magnificus prof. dr. E. Sterken
en volgens besluit van het College voor Promoties.

De openbare verdediging zal plaatsvinden op
vrijdag 6 april 2018 om 16:15 uur

door

Matthijs Dries

geboren op 10 februari 1986
te Emmen

Promotores

Prof. dr. S. C. Trager

Prof. dr. L. V. E. Koopmans

Beoordelingscommissie

Prof. dr. C. Conroy

Prof. dr. R. F. Peletier

Prof. dr. T. Treu

ISBN: 978-94-034-0465-3 (printed version)

ISBN: 978-94-034-0464-6 (electronic version)

Cover: Artist impression of the elliptical galaxy Messier 49 unraveling into a spectrum that can be used to infer the IMF of the galaxy

Design by: Caroline Boerema

Image credit: NASA and The Hubble Heritage Team (STScI/AURA) for the original image of M49 and the Digitized Sky Survey (DSS), STScI/AURA, Palomar/Caltech and UKSTU/AAO for the background star field

Acknowledgement: The work contained in this thesis was supported by an NWO grant (project number 614.001.208)

Printed by: Gildeprint - Enschede



Contents

1	Introduction	1
1.1	Formation and evolution of galaxies	4
1.2	Elliptical galaxies	6
1.3	Stellar population synthesis	7
1.3.1	Single Stellar Populations	8
1.3.2	Composite Stellar Populations	10
1.3.3	Population Synthesis models	11
1.4	The initial mass function	12
1.5	Bayesian model comparison	14
1.6	This thesis	19
	References	21
2	Bayesian inference of the IMF in Single Stellar Populations	25
2.1	Introduction	27
2.2	Hierarchical Bayesian inference	29
2.2.1	The first level of inference	32
2.2.2	The second level of inference	40
2.3	Stellar templates	41
2.3.1	Isochrones	42
2.3.2	Stellar library	42
2.3.3	Interpolator	44
2.4	Results - mock single stellar populations	48
2.4.1	The regularization scheme	49
2.4.2	First level of inference	49

2.4.3	Second level of inference	66
2.5	Results - model versus model	67
2.5.1	The regularization scheme	67
2.5.2	IMF reconstruction of MILES SSPs	68
2.5.3	Mass fraction of low-mass stars	71
2.6	Summary and discussion	73
Appendix 2.A	Velocity dispersion mock SSPs	76
Appendix 2.B	Results for mock SSPs	77
Appendix 2.C	Results for MILES SSPs	89
References	94
3	Bayesian inference of the IMF in CSPs	97
3.1	Introduction	99
3.2	Model description	101
3.2.1	Hierarchical Bayesian framework	101
3.2.2	New model features	103
3.3	Stellar templates	109
3.3.1	Isochrones	109
3.3.2	Stellar library and interpolator	110
3.3.3	Isochrone binning	110
3.4	Composite stellar population mock spectra	111
3.4.1	SFHs from semi-analytic models	111
3.4.2	CSP mock spectra with a variable IMF	113
3.5	Results	116
3.5.1	The required number of SSPs	116
3.5.2	Reconstructed IMFs	119
3.5.3	Reconstructed spectra	122
3.5.4	Reconstructing the variable IMF	122
3.6	Summary and discussion	128
Appendix 3.A	Binning isochrone stars	132
Appendix 3.B	Systematic uncertainties	134
References	137
4	Hierarchical Bayesian inference of the IMF in ETGs	141
4.1	Introduction	143
4.2	Model description	146
4.2.1	Hierarchical Bayesian framework	147
4.2.2	New features: abundance and error patterns	149

4.3	The MIX stellar population models	151
4.3.1	The X-shooter Spectral Library	152
4.3.2	The MIX library	155
4.3.3	Additional M dwarfs	156
4.3.4	Optimizing the MIX library	157
4.3.5	Stellar templates	160
4.4	The data	162
4.5	Results	166
4.5.1	Results <code>run1</code> and <code>run2</code>	167
4.5.2	Abundance variations	177
4.5.3	Isochrones	182
4.5.4	The regularization scheme	184
4.5.5	The IMF parameterization	185
4.5.6	IMF shape from theoretical models of star formation	189
4.5.7	Mass-to-light ratios	191
4.6	Summary and discussion	193
	Appendix 4.A SSP-based response functions	197
	References	200
5	Conclusions and future outlook	205
5.1	Results, chapter by chapter	206
5.2	Main conclusions	210
5.3	Future outlook	212
	References	215
A	HSPS manual	217
	HSPS manual	217
A.1	Installation	219
A.1.1	Installation stellar templates	219
A.2	Input file	220
A.3	Stellar templates	220
A.4	Setup files	221
A.4.1	Parameterized version of HSPS	221
A.4.2	Full version of HSPS	224
A.4.3	Values file	225
A.5	Running the code	227
A.6	Output files	228

A.7 Running the pipeline	228
A.8 Running a test spectrum	230
References	231
Nederlandse Samenvatting	233
Acknowledgements	243