

This article was downloaded by: [Vrije Universiteit, Library]

On: 26 November 2010

Access details: Access Details: [subscription number 907218003]

Publisher Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Cognitive Behaviour Therapy

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713926011>

Internet-Based and Other Computerized Psychological Treatments for Adult Depression: A Meta-Analysis

Gerhard Andersson^a; Pim Cuijpers^b

^a Department of Behavioural Sciences and Learning, Swedish Institute for Disability Research, Linköping University, Linköping, and Department of Clinical Neuroscience, Psychiatry Section, Karolinska Institutet, Stockholm, Sweden ^b Department of Clinical Psychology and EMGO+ Institute, VU University, Amsterdam, the Netherlands

Online publication date: 15 December 2009

To cite this Article Andersson, Gerhard and Cuijpers, Pim(2009) 'Internet-Based and Other Computerized Psychological Treatments for Adult Depression: A Meta-Analysis', *Cognitive Behaviour Therapy*, 38: 4, 196 – 205

To link to this Article: DOI: 10.1080/16506070903318960

URL: <http://dx.doi.org/10.1080/16506070903318960>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Internet-Based and Other Computerized Psychological Treatments for Adult Depression: A Meta-Analysis

Gerhard Andersson¹ and Pim Cuijpers²

¹*Department of Behavioural Sciences and Learning, Swedish Institute for Disability Research, Linköping University, Linköping, and Department of Clinical Neuroscience, Psychiatry Section, Karolinska Institutet, Stockholm, Sweden;* ²*Department of Clinical Psychology and EMGO+ Institute, VU University, Amsterdam, the Netherlands*

Abstract. Computerized and, more recently, Internet-based treatments for depression have been developed and tested in controlled trials. The aim of this meta-analysis was to summarize the effects of these treatments and investigate characteristics of studies that may be related to the effects. In particular, the authors were interested in the role of personal support when completing a computerized treatment. Following a literature search and coding, the authors included 12 studies, with a total of 2446 participants. Ten of the 12 studies were delivered via the Internet. The mean effect size of the 15 comparisons between Internet-based and other computerized psychological treatments vs. control groups at posttest was $d = 0.41$ (95% confidence interval [CI]: 0.29–0.54). However, this estimate was moderated by a significant difference between supported ($d = 0.61$; 95% CI: 0.45–0.77) and unsupported ($d = 0.25$; 95% CI: 0.14–0.35) treatments. The authors conclude that although more studies are needed, Internet and other computerized treatments hold promise as potentially evidence-based treatments of depression. *Key words:* computerized treatment; depression; Internet-based; role of support

Received 15 August, 2009; Accepted 8 September, 2009

*Correspondence address: Gerhard Andersson, PhD, Department of Behavioural Sciences, Linköping University, SE-581 83 Linköping, Sweden. Tel: +46 13 28 58 40; Fax: +46 13 28 21 45. E-mail: Gerhard.Andersson@liu.se

Adult depression is a costly condition for which numerous different treatments have been developed (Ebmeier, Donaghey, & Steele, 2006). Among the psychological treatment options, several have been found to be effective (Cuijpers, van Straten, Andersson, & van Oppen, 2008), with no or minor differences between the main treatment alternatives. Cognitive behaviour therapy (CBT), however, has been investigated in by far the most trials and in different administration formats such as group (McDermut, Miller, & Brown, 2001) and telephone-assisted (e.g. Simon, Ludman, Tutty, Operskalski, & Von Korff, 2004) treatments. Computer and, more recently, Internet delivery have become increasingly common administration formats for depression treatment in research and slowly

but gradually in clinical settings as well. Computerized psychological treatments can be delivered on devices such as stand-alone or Internet-linked computers, PCs, palmtops, phone-interactive voice response systems, CD-ROMS, DVDs, cell phones, and VR equipment (Marks, Cavanagh, & Gega, 2007). However, during the last 5 years, it is mostly Internet-delivered treatments that have been tested in research.

Internet-delivered treatments of depression can take on different forms. One approach is largely based on bibliotherapy, with mainly text-based materials and guidance by a therapist via e-mail or phone. On the other end of the continuum, we find treatments that are briefer and usually do not target persons with clinical depression. Moreover, these

treatments are commonly not delivered with guidance and can hence be reached by more people at a lower cost. They are, however, probably less effective (Spek, Cuijpers, et al., 2007). Regardless of the role of support, treatment protocols also differ with regard to how often and how many online activities are included and how much feedback is automated.

There are now several reviews and meta-analyses on computerized (e.g. Cuijpers, Marks, et al., 2009; Reger & Gahm, 2009) and Internet-delivered (Barak, Hen, Boniel-Nissim, & Shapira, 2008; Griffiths & Christensen, 2006; Spek, Cuijpers, et al., 2007) treatments. In one previous review, Internet treatments for depression were covered (Andersson, 2006), but this was not a meta-analysis. The other meta-analyses were not specifically aimed at depression and have included only a limited portion of currently available trials. Given the very strong increase in the number of controlled trials on computerized and Internet-delivered treatments of depression in the past few years, we decided to conduct a new meta-analysis in which we focused on depression only. We were interested in investigating the role of support and the overall effects compared with other treatments. This meta-analysis adds to the literature by providing an overview of the field and by contrasting different approaches to computerized treatment.

Method

Identification and selection of studies

We used several methods to identify studies for inclusion. First, we used a database of 1036 studies on the psychological treatment of depression, which includes reports on combined treatments and comparisons with pharmacotherapy. This database has been described in detail elsewhere (Cuijpers, van Straten, Warmerdam, & Andersson, 2008) and has been used in a series of earlier meta-analyses (<http://www.evidencebasedpsychotherapies.org>). The database was developed through a comprehensive literature search (from 1966 to January 2009), in which we examined a total of 9011 abstracts: 1629 from PubMed, 2439 from PsycINFO, 2606 from Embase, and 2337 from the Cochrane Central Register of Controlled Trials. These abstracts

were identified by combining terms indicative of psychological treatment and depression (both MeSH terms and text words). For this database, we also collected the primary studies from 42 meta-analyses of psychological treatment for depression (<http://www.evidencebasedpsychotherapies.org>). For the current study, we examined the full texts of these 1036 studies, then examined the reference lists of earlier reviews and meta-analyses (e.g. Spek, Cuijpers, et al., 2007), and checked the references of the included primary studies.

We included (a) randomized trials (b) in which the effects of an Internet-based or computerized psychological treatment (c) were compared with a (noncomputerized) control or comparison group or a face-to-face psychological treatment (d) in adults (e) with depression (established by diagnostic interview or elevated levels of depressive symptoms based on self-report measures). We also included studies that were aimed at adults with depression and anxiety. We excluded studies on inpatients and those on adolescents or children (<18 years). We did not include component studies (e.g. Christensen, Griffiths, Mackinnon, & Brittliffe, 2006).

Comorbid general medical or psychiatric disorders were not used as an exclusion criterion. No language restrictions were applied.

Meta-analyses

For each comparison between Internet-based or computerized psychological treatment and control conditions, we calculated the effect size indicating the difference between the two conditions at posttest (Cohen's *d* or standardized mean difference). We calculated the effect sizes by subtracting (at posttest) the average score of the treatment group from the average score of the control group and dividing the result by the pooled standard deviations of the two groups. Effect sizes of 0.8 can be assumed to be large, 0.5 moderate and 0.2 small (Cohen, 1988).

In the calculations of effect sizes, we only used those instruments that explicitly measured symptoms of depression. If more than one depression measure was used, the mean of the effect sizes was calculated, so that each study only provided one effect size. We only used the effect sizes indicating the differences between the two types of treatment

at posttest. We decided not to examine the differential effects at follow-up because the number of effect sizes was too low. In addition, the follow-up period differed considerably among these studies.

To calculate pooled mean effect sizes, we used Biostat's computer program *Comprehensive Meta-Analysis* (version 2.2.021). Because we expected considerable heterogeneity among the studies, we decided to calculate mean effect sizes using a random-effects model. In the random-effects model, it is assumed that the included studies are drawn from "populations" of studies that differ from each other systematically (heterogeneity). In this model, the effect sizes resulting from included studies differ not only because of the random error within studies (as in the fixed-effects model) but also because of true variation in effect size from one study to the next.

As a test of homogeneity of effect sizes, we calculated the I^2 statistic, which is an indicator of heterogeneity in percentages. A value of 0% indicates no observed heterogeneity, and larger values show increasing heterogeneity, with 25% as low, 50% as moderate, and 75% as high heterogeneity (Higgins, Thompson, & Deeks, 2003). We also calculated the Q statistic but only report whether this was significant or not.

Subgroup analyses were conducted according to the mixed-effect model. In this model, studies within subgroups are pooled with the random-effects model, whereas tests for significant differences between subgroups are conducted with the fixed-effects model.

Publication bias was tested by inspecting the funnel plot on primary outcome measures and by Duval and Tweedie's (2000) trim and fill procedure, which yields an estimate of the effect size after the publication bias has been taken into account (as implemented in *Comprehensive Meta-Analysis*, version 2.2.021).

Results

Characteristics of included studies

A total of 12 studies, with 2446 participants (1324 in the Internet-based and computerized psychological treatment conditions, 996 in the control conditions, and 126 in the face-to-face comparison conditions) met all inclusion

criteria. Selected characteristics of these studies are presented in Table 1.

Ten studies were aimed at adults in general, and one was aimed at older adults and one at young adults. All but one study recruited participants from the community. Only two studies included participants with depressive disorder diagnosed in a formal diagnostic interview. Six studies used a wait list control group, four a care-as-usual control group, and the remaining two studies another type of control group. The 12 studies included 15 comparisons between an Internet-based or computerized psychological treatment and a control group (three studies included two comparisons). In three studies, Internet-based or computerized psychological treatment was compared with face-to-face psychological treatment. Eleven of the 15 comparisons examined CBT, two problem-solving therapy, and one psychoeducation. Two studies examined computerized psychological treatment and the remaining 10 Internet-based psychological treatment. Five studies were conducted in the United States, four in the Netherlands, and one each in Sweden, United Kingdom, and Australia. In one study only 50% of the face-to-face treatment was replaced by a computerized treatment (Wright et al., 2005). We decided to include this study and examine whether removal of this resulted in changes of the mean effect size (which was not the case; see later discussion).

Internet-based and computerized psychological treatment versus control groups: overall effect size

The mean effect size of the 15 comparisons between Internet-based and computerized psychological treatment vs. control groups at posttest was 0.41 (95% confidence interval [CI]: 0.29–0.54; Table 2). Heterogeneity was moderate to high ($I^2 = 57.49$). The effect sizes and 95% CIs of the individual contrast groups are plotted in Figure 1.

Inspection of the forest plot suggested that two studies were possible outliers (Selmi et al., 1990; Wright et al., 2005). However, after removal of these studies, the effect size remained almost the same ($d = 0.37$; 95% CI: 0.26–0.49) and heterogeneity remained at a moderate level ($I^2 = 51.36$).

Table 1. Selected characteristics of randomized controlled studies examining the effects of computerized and Internet-based psychotherapies for adult depression

Study	Target population	Recruitment	Inclusion	Condition	N	Intervention	Therapist support	Internet/computer	Measure ^a	Country
Andersson et al. (2005)	Adults (≥ 18 yrs)	Community	$p > .55$ for MDD (CIDI-SF) + MADRS-S 15-30	1. iCBT 2. WL/online discussion group 1. iCBT	36 39	5 modules CBT and BA	E-mail	I, 10 wks	BDI; MADRS-S	Sweden
Christensen et al. (2004)	Adults (18-52 yrs)	Posted questionnaire	K-10 ≥ 22	1. iCBT 2. iPE 3. Attention placebo 1. iCBT	182 165 178	1. 5 CBT modules 2. 5 PE modules	Telephone calls by lay interviewers (all conditions) -	I, 6 wks	CES-D	Australia
Clarke et al. (2002)	Adults	Via HMO	Depressed, nondepressed members	1. iCBT 2. CAU 1. iCBT-a	144 155	7 chapters CBT	None	I, NSD	CES-D	United States
Clarke et al. (2005)	Adults	Via HMO	Depressed, nondepressed members	2. CAU 1. iCBT-a	75	7 chapters CBT	None: iCBT-a: postcard reminders; iCBT-b: telephone reminders	I, NSD	CES-D	United States
Clarke et al. (2009)	Young adults (18-24 yrs)	Via HMO	Depressed, nondepressed members	2. iCBT-b 3. CAU 1. iCBT	80 100 83	4 sections CBT	None	I, NSD	PHQ-9	United States
Proudfoot et al. (2004)	Adults (18-75 yrs)	Primary care	GHQ-12 > 4 + CIS-R > 12 (depression or anxiety)	2. CAU 1. cCBT	77 112	8 sessions cCBT	5 min help at beginning, end of each session	I, 8 weeks	BDI	United Kingdom
Ruwaard et al. (2009)	Adults (≥ 18 yrs)	Community	BDI 10-29	2. CAU 1. iCBT	109 36	8 phases CBT, BA	Asynchronous contact	I, 11 wks	BDI; SCL-90-R-D	The Netherlands
Selmi et al. (1990)	Adults	Community	SCL-90-R > 65th percentile + BDI ≥ 16 + major, intermittent, minor depression (RDC-SADS)	2. WL 1. cCBT	18 12	6 sessions CBT	Help at beginning, end of each session	C; 6 wks	BDI; HAM-D; SCL-90-R-D	United States

Table 1. *Continued*

Study	Target population	Recruitment	Inclusion	Condition	N	Intervention	Therapist support	Internet/computer	Measure ^a	Country
Spek, Nyklicek et al. (2007)	Older adults (≥ 50 yrs)	Community	Subthreshold depression (EDS ≥ 12 , no MDD)	2. ftf-CBT	12	8 weekly modules CBT	None	I; 8 wks	BDI	The Netherlands
				3. WL	12					
				1. iCBT	102					
Van Straten et al. (2008)	Adults	Community	Self-defined depression or anxiety	2. gCBT	99	5 weekly modules PST	E-mail	I; 5 wks	CES-D; MDI	The Netherlands
				3. WL	100					
				1. iPST	107					
Warmerdam et al. (2008)	Adults	Community	CES-D ≥ 16	2. WL	106	1. 8 weekly modules CBT; 2.5 weekly modules PST	E-mail	I; group 1, 8 wks; group 2, 5 wks	CES-D	The Netherlands
				1. iCBT	88					
Wright et al. (2005)	Adults (18–65 yrs)	Community	MDD (SCID) + BDI ≥ 14	2. iPST	87	9 sessions CBT	25-min ftf sessions +25-min cCBT	C; 8 wks	BDI; HAM-D	United States
				3. WL	87					
				1. eCBT + ftf	15					
				2. ftf-only CBT	15					
				3. WL	15					

Note. BA, behavioural activation; BDI, Beck Depression Inventory; C, computerized treatment delivery; CAU, care-as-usual; CBT, cognitive behaviour therapy; cCBT, computerized CBT; gCBT, group CBT; CES-D, Center for Epidemiological Studies–Depression scale; CIDI-SF, Composite International Diagnostic Interview Short-Form; EDS, Edinburgh Depression Scale; ftf, face-to-face; GHQ, General Health Questionnaire; HAM-D, Hamilton Depression Rating Scale; HMO, health maintenance organization; I, Internet treatment delivery; iCBT, Internet-based CBT; iPE, Internet-based psychoeducation; iPST, Internet-based problem-solving therapy; K-10, Kessler-10; MADRS-S, Montgomery–Asberg Depression Rating Scale; MDD, major depressive disorder; MDI, Major Depression Inventory; PE, psychoeducation; PHQ-9, Patient Health Questionnaire; PST, problem-solving therapy; RDC, Research Diagnostic Criteria; SADS, Schedule for Affective Disorders; NSD, no standard duration; SCID, Structured Clinical Interview for DSM-IV; SCL-90-R, Symptom Checklist-90-R; SCL-90-R-D, SCL-90-R Depression subscale; WL, wait-list.

^aOnly the instruments that were used to calculate effect sizes are included.

Table 2. Meta-analyses of studies examining the effects of computerized and Internet-based psychological treatments for adult depression

Study	N_{comp}	d	95% CI	Z	I^2 ^a	p ^b
Overall effects						
All studies	15	0.41	0.29–0.54	6.47****	57.49***	
One ES per study (highest)	12	0.45	0.29–0.61	5.39****	65.35***	
One ES per study (lowest)	12	0.43	0.27–0.59	5.21****	64.86***	
Two possible outliers removed ^c	13	0.37	0.26–0.49	6.41****	51.36**	
Subgroup analyses ^d						
Type						.107
Internet-based	12	0.37	0.24–0.49	5.84****	53.83**	
Computerized	3	0.85	0.27–1.43	2.89***	61.18*	
Control group						.038
Care-as-usual	5	0.23	0.06–0.40	2.68***	46.34	
Wait-list	7	0.56	0.37–0.76	5.60****	43.51	
Other	3	0.45	0.21–0.69	3.64****	59.38*	
Psychological treatment						.923
CBT	12	0.42	0.26–0.59	5.09****	64.82***	
Other	3	0.41	0.27–0.56	5.51****	0	
Anxiety allowed						.234
Only depression	13	0.38	0.25–0.51	5.75****	57.26***	
Depression or anxiety	2	0.64	0.24–1.04	3.13***	42.73	
Professional support						.000
Support	8	0.61	0.45–0.77	7.67****	23.74	
No professional support	7	0.25	0.14–0.35	4.75****	10.41	
C or I psychotherapy vs. ftf psychotherapy (all studies)	3	–0.05	–0.29–0.20	–0.36	0	

Note. CBT, cognitive behaviour therapy; C, computerized treatment delivery; CI, confidence interval; ES, effect size; ftf, face to face; I, Internet treatment delivery; N_{comp} , number of comparisons. ^aThe p values indicate whether the Q statistic is significant (the I^2 statistics does not include a test of significance). ^bThe p values indicate whether the difference between the effect sizes in the subgroups is significant. ^cSelmi et al., 1990; Wright et al., 2005. ^dAll subgroup analyses were conducted with mixed-effects analyses. * $p < .10$. ** $p < .05$. *** $p < .01$. **** $p < .001$.

In our analyses, we included three studies in which two psychological treatments were compared with the same control group. This means that multiple comparisons from these three studies were included in the same analysis. These multiple comparisons, however, are not independent of each other, which may have resulted in an artificial reduction of heterogeneity. Therefore, we conducted another meta-analysis, in which we included only one comparison per study (Table 2). From the three studies with multiple comparisons, we first included only the comparison with the largest effect size. As can be seen in Table 2, these analyses did indicate that heterogeneity increased somewhat in some analyses ($I^2 = 65.35$), although the effect size did not differ very much from the overall analyses. Then we repeated these analyses and

included only the smallest effect size of the three studies with multiple comparisons. These analyses also resulted in increased heterogeneity ($I^2 = 64.86$) and a comparable effect size.

Neither the funnel plots nor Duval and Tweedie's trim and fill procedure indicated a significant publication bias.

Subgroup analyses

We conducted several subgroup analyses (Table 2). These included type of psychological treatment (Internet-based vs. computerized), type of control group (care-as-usual, wait-list, other), content of psychological treatment (CBT vs. other), whether the study only included participants with depression or also persons with anxiety, and whether there was professional support during the therapy (yes or no).




Figure 1. Internet-based and other computerized treatments of depression compared to control groups: standardized effect sizes. The letters A and B in the above figure refer to different subsamples within each study.

As can be seen in Table 2, the type of control group was significantly associated with the effect sizes ($p < .05$). Studies with a wait-list control had higher effect sizes than care-as-usual and other control groups. However, heterogeneity remained high in these subgroups. We also found that studies in which no professional support was given had lower effect sizes than those in which support was given ($p < .000$). The heterogeneity in these two groups was low ($I^2 < 25\%$).

Because the study by Christensen, Griffiths, and Jorm (2004) was complex to interpret (there was some support by telephone but no clear help in working through the treatment), we repeated this subgroup analysis without this study. The results were, however, comparable (unguided: $d = 0.18$; 95% CI = 0.05–0.30, $I^2 = 0$; guided: $d = 0.61$, 95% CI = 0.45–0.77, $I^2 = 23.74$; p difference = .000).

Discussion

The aim of this meta-analysis was to summarize the literature on Internet-based and other computerized psychological treatments for adult depression. We found an overall effect size of $d = 0.41$, but that estimate is probably not meaningful because it hides the finding that interventions in which

support is provided to the participant are more effective. Indeed, the computerized interventions with support showed an average between-group effect size of $d = 0.61$, whereas the unsupported treatments had a much smaller effect of $d = 0.25$. This is similar to the findings by Spek, Cuijpers, et al. (2007), who found that interventions without support had an average effect size of $d = 0.24$, whereas Internet interventions with support had a large mean effect size of $d = 1.0$. Although the meta-analyses overlap in terms of studies included, we included more recent studies and also other computerized treatments not covered in the Spek et al. study (e.g. not Internet-delivered). Overall, it appears that computerized treatments with therapist support are much more effective than unsupported treatments, and this has been confirmed in open studies as well (e.g. Christensen, Griffiths, Groves, & Korten, 2006). However, the concept of support is not fully investigated in the literature, and we cannot exclude the possibility that some forms of support can be automated or that other factors, such as having a clear deadline for completion of a treatment with a scheduled follow-up (e.g. a telephone interview), would make less-supported treatments more effective (Nordin, Carlbring, Cuijpers, & Andersson, in press). Clearly, there is a need to investigate

Downloaded By: [Vrije Universiteit, Library] At: 11:10 26 November 2010

the concept of support further and the role of therapist factors in computerized treatments (e.g. Almlöv, Carlbring, Berger, Cuijpers, & Andersson, 2009).

There were few studies available for the contrast between Internet-based or other computerized treatments and face-to-face treatments. Here we found no difference, which is in line with other studies in the field of anxiety disorders (e.g. Carlbring et al., 2005; Kiropoulos et al., 2008). There is a need to further test whether computerized treatment, and Internet-delivered treatment in particular, can be as effective as face-to-face treatments in depression. These studies need to be designed as equivalence studies, and indeed in our meta-analysis the finding of no difference could be regarded as a promising sign of equivalence between the treatment formats. Although the effects of supported computerized treatments in this meta-analysis are in line with those of previous meta-analyses for psychological treatments (e.g. Cuijpers, van Straten, Andersson, et al., 2008), they are somewhat lower. However, the effects are not low when compared with the effects of psychological treatments in primary care (Cuijpers, van Straten, van Schaik, & Andersson, 2009). Several other findings are worth commenting on. We did not find a significant difference between computerized and Internet-delivered treatments, but this could be due to power problems. In line with findings of other studies, the type of control group influenced the effect size estimate, with lower effects when treatment as usual was the comparison group. This was expected because no treatment is worse than some treatment. We did not find any differences between CBT-oriented and other forms of computerized treatments. This comparison was unbalanced, with most of the studies being done from a CBT perspective, but we welcome more studies on computerized treatments from a non-CBT perspective, because it is not settled whether other evidence-based psychological treatments such as interpersonal psychotherapy are possible to transfer to the computer medium or the Internet. We also did not find any effects of comorbid anxiety on the effect size estimate, but this needs to be further explored in future studies, because there are very few studies on computer interventions for

depression in which other conditions than depression have been assessed.

In this meta-analysis, we did not present data on quality assessment. However, all studies were checked using the four basic criteria, as suggested in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins & Green, 2005): allocation to conditions conducted by an independent (third) party, blinding of assessors of outcomes, completeness of follow-up data, and adequacy of random allocation concealment to respondents. Overall, the study quality was not satisfactory, but studies are increasingly following the CONSORT guidelines (e.g. Boutron, Moher, Altman, Schulz, & Ravaud, 2008).

There are several possible challenges for future research. First, Internet-delivered and other computerized psychological interventions are still not supported by a solid database, and many studies have failed to perform a proper diagnostic assessment (Andersson & Cuijpers, 2008). There is a need for accurate diagnostic procedures in future trials. Second, most studies have been done either in an academic setting with participants recruited via advertisement or by an epidemiological screening approach. Only one of the studies in our meta-analysis included patients recruited from primary care. There is a need for effectiveness studies of Internet and other computerized treatments involving patients from psychiatric settings. Third, studies on long-term effects of treatment are largely lacking. There is a 1-year follow-up of the trial by Christensen et al. (2004), which showed some remaining benefits (Mackinnon, Griffiths, & Christensen, 2008), but more research is needed, in particular, because help-seeking and health care use might be affected as an effect of treatment (Christensen, Leach, Barney, Mackinnon, & Griffiths, 2006). Fourth, our meta-analysis did not cover the issue of whom Internet and other computerized treatments are suitable for. For example, one study found an indication that number of previous depression episodes was related to worse treatment outcome (Andersson, Bergström, Holländare, Ekselius, & Carlbring, 2004). It is also important to investigate differential predictors of outcome for different treatment formats (Spek, Nyklicek, Cuijpers, & Pop, 2008).

This meta-analytic review has some limitations. First, we included very heterogeneous treatments and samples. This can be seen as an advantage for a meta-analysis because we then could investigate differences between studies. However, because relatively few studies could be located, we were underpowered to detect effects for some contrasts. A second limitation has to do with the selection of adult samples only. Our impression is that the available studies on adolescents do not alter our main finding that support may be needed to achieve good outcomes (e.g. O’Kearney, Gibson, Christensen, & Griffiths, 2006). A third limitation concerns the methodological quality of the studies that we did not fully report. For example, we did not include analyses on the drop-out rate for different treatments, which is a known problem in Internet interventions.

Despite the limitations of this meta-analysis, we believe there is emerging evidence that Internet and other computerized intervention can be helpful in reducing symptoms of depression.

Acknowledgments

The preparation of this study was supported in part by a grant from the Swedish Research Council.

References

References marked with an asterisk indicate studies included in the meta-analysis.

- Almlöv, J., Carlbring, P., Berger, T., Cuijpers, P., & Andersson, G. (2009). Therapist factors in Internet-delivered CBT for major depressive disorder. *Cognitive Behaviour Therapy, 38*(4), 247–254.
- Andersson, G. (2006). Internet based cognitive behavioral self-help for depression. *Expert Review of Neurotherapeutics, 6*, 1637–1642.
- *Andersson, G., Bergström, J., Holländare, F., Carlbring, P., Kaldo, V., & Ekselius, L. (2005). Internet-based self-help for depression: A randomised controlled trial. *British Journal of Psychiatry, 187*, 456–461.
- Andersson, G., Bergström, J., Holländare, F., Ekselius, L., & Carlbring, P. (2004). Delivering CBT for mild to moderate depression via the Internet. Predicting outcome at 6-months follow-up. *Verhaltenstherapie, 14*, 185–189.
- Andersson, G., & Cuijpers, P. (2008). Pros and cons of online cognitive-behavioural therapy. *British Journal of Psychiatry, 193*, 270–271.

- Barak, A., Hen, L., Boniel-Nissim, M., & Shapira, N. (2008). A comprehensive review and a meta-analysis of the effectiveness of Internet-based psychotherapeutic interventions. *Journal of Technology in Human Services, 26*, 109–160.
- Boutron, I., Moher, D., Altman, D. G., Schulz, K. F., & Ravaud, P. (2008). Extending the CONSORT statement to randomized trials of nonpharmacologic treatment: Explanation and elaboration. *Annals of Internal Medicine, 148*, 295–309.
- Carlbring, P., Nilsson-Ihrfelt, E., Waara, J., Kollenstam, C., Buhrman, M., Kaldo, V., et al. (2005). Treatment of panic disorder: Live therapy vs. self-help via Internet. *Behaviour Research and Therapy, 43*, 1321–1333.
- Christensen, H., Griffiths, K., Groves, C., & Korten, A. (2006). Free range users and one hit wonders: Community users of an Internet-based cognitive behaviour therapy program. *Australian and New Zealand Journal of Psychiatry, 40*, 59–62.
- *Christensen, H., Griffiths, K. M., & Jorm, A. (2004). Delivering interventions for depression by using the Internet: Randomised controlled trial. *British Medical Journal, 328*, 265–268.
- Christensen, H., Griffiths, K.M., Mackinnon, A.J., & Brittliffe, K. (2006). Online randomized trial of brief and full cognitive behaviour therapy for depression. *Psychological Medicine, 36*, 1737–1746.
- Christensen, H., Leach, L. S., Barney, L., Mackinnon, A. J., & Griffiths, K. M. (2006). The effect of web based depression interventions on self reported help seeking: Randomised controlled trial. *BMC Psychiatry, 6*, 13.
- *Clarke, G., Eubanks, D., Reid, E., Kelleher, C., O’Connor, E., DeBar, L. L., et al. (2005). Overcoming depression on the Internet (ODIN) (2): A randomized trial of a self-help depression skills intervention program with reminders. *Journal of Medical Internet Research, 7*, e16.
- *Clarke, G., Kelleher, C., Hornbrook, M., DeBar, L., Dickerson, J., & Gullion, C. (2009). Randomized effectiveness trial of an Internet, pure self-help, cognitive behavioral intervention for depressive symptoms in young adults. *Cognitive Behaviour Therapy, 38*(4), 222–234.
- *Clarke, G., Reid, E., Eubanks, D., O’Connor, E., DeBar, L. L., Kelleher, C., et al. (2002). Overcoming depression on the Internet (ODIN): A randomized controlled trial of an Internet depression skills intervention program. *Journal of Medical Internet Research, 4*, e14.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ: Erlbaum.
- Cuijpers, P., Marks, I., van Straten, A.-M., Cavanagh, K., Gega, L., & Andersson, G. (2009). Computer-aided psychotherapy for anxiety disorders: A meta-analytic review. *Cognitive Behaviour Therapy, 38*, 66–82.
- Cuijpers, P., van Straten, A., Andersson, G., & van Oppen, P. (2008). Psychotherapy for depression in adults: A meta-analysis of comparative

- outcome studies. *Journal of Consulting and Clinical Psychology*, 76, 909–922.
- Cuijpers, P., van Straten, A., van Schaik, A., & Andersson, G. (2009). Psychological treatment of depression in primary care: A meta-analysis. *British Journal of General Practice*, 59, e51–60.
- Cuijpers, P., van Straten, A., Warmerdam, L., & Andersson, G. (2008). Psychological treatment of depression: A meta-analytic database of randomized studies. *BMC Psychiatry*, 8, 36.
- Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56, 455–463.
- Ebmeier, K. P., Donaghey, C., & Steele, J.D. (2006). Recent developments and current controversies in depression. *Lancet*, 367, 153–167.
- Griffiths, K. M., & Christensen, H. (2006). Review of randomised controlled trials of Internet interventions for mental disorders and related conditions. *Clinical Psychologist*, 10, 16–29.
- Higgins, J. P. T., & Green, S. (2005). Cochrane handbook for systematic reviews of interventions 4.2.5. In *The Cochrane Library, Issue 3*. Chichester, UK: Wiley.
- Higgins, J. P., Thompson, S. G., & Deeks, J. J. (2003). Measuring inconsistency in meta-analyses. *British Medical Journal*, 327, 557–560.
- Kiropoulos, L. A., Klein, B., Austin, D. W., Gilson, K., Pier, C., Mitchell, J., et al. (2008). Is Internet-based CBT for panic disorder and agoraphobia as effective as face-to-face CBT? *Journal of Anxiety Disorders*, 22, 1273–1284.
- Mackinnon, A. J., Griffiths, K. M., & Christensen, H. (2008). Comparative randomised trial of on-line cognitive behaviour therapy and an information website for depression: 12 month outcomes. *British Journal of Psychiatry*, 192, 130–134.
- Marks, I. M., Cavanagh, K., & Gega, L. (2007). *Hands-on help*. Maudsley Monograph no. 49. Hove, UK: Psychology Press.
- McDermut, W., Miller, I. W., & Brown, R. A. (2001). The efficacy of group psychotherapy for depression: A meta-analysis and review of the empirical research. *Clinical Psychology: Science and Practice*, 8, 98–116.
- Nordin, S., Carlbring, P., Cuijpers, P., & Andersson, G. (in press). Expanding the limits of bibliotherapy for panic disorder. Randomized trial of self-help without support but with a clear deadline. *Behavior Therapy*.
- O’Kearney, R., Gibson, M., Christensen, H., & Griffiths, K. M. (2006). Effects of a cognitive-behavioural Internet program on depression, vulnerability to depression and stigma in adolescent males: A school-based controlled trial. *Cognitive Behaviour Therapy*, 35, 43–54.
- *Proudfoot, J., Ryden, C., Everitt, B., Shapiro, D. A., Goldberg, D., Mann, A., et al. (2004). Clinical efficacy of computerised cognitive-behavioural therapy for anxiety and depression in primary care: Randomised controlled trial. *British Journal of Psychiatry*, 185, 46–54.
- Reger, M. A., & Gahm, G. A. (2009). A meta-analysis of the effects of Internet- and computer-based cognitive-behavioral treatments for anxiety. *Journal of Clinical Psychology*, 65, 53–75.
- *Ruwaard, J., Schriecken, B., Schrijver, M., Broeksteeg, J., Dekker, J., Vermeulen, H., et al. (2009). Standardized web-based CBT of mild to moderate depression: A randomized controlled trial with a long-term follow-up. *Cognitive Behaviour Therapy*, 38(4), 206–221.
- *Selmi, P. M., Klein, M. H., Greist, J. H., Sorrell, S. P., & Erdman, H. P. (1990). Computer-administrated cognitive behavioral therapy for depression. *American Journal of Psychiatry*, 147, 51–56.
- Simon, G. E., Ludman, E. J., Tutty, S., Operaskalski, B., & Von Korff, M. (2004). Telephone psychotherapy and telephone care management for primary care patients starting antidepressant treatment: A randomized controlled trial. *Journal of the American Medical Association*, 292, 935–942.
- Spek, V., Cuijpers, P., Nyklicek, I., Riper, H., Keyzer, J., & Pop, V. (2007). Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: A meta-analysis. *Psychological Medicine*, 37, 319–328.
- Spek, V., Nyklicek, I., Cuijpers, P., & Pop, V. (2008). Predictors of outcome of group and Internet-based cognitive behavior therapy. *Journal of Affective Disorders*, 105, 137–145.
- *Spek, V., Nyklicek, I., Smits, N., Cuijpers, P., Riper, H., Keyzer, J., et al. (2007). Internet-based cognitive behavioural therapy for sub-threshold depression in people over 50 years old: A randomized controlled clinical trial. *Psychological Medicine*, 37, 1797–1806.
- *van Straten, A., Cuijpers, P., & Smits, N. (2008). Effectiveness of a web-based self-help intervention for symptoms of depression, anxiety, and stress: Randomized controlled trial. *Journal of Medical Internet Research*, 10(1), e7.
- *Warmerdam, L., van Straten, A., Twisk, J., Riper, H., & Cuijpers, P. (2008). Internet-based treatment for adults with depressive symptoms: Randomized controlled trial. *Journal of Medical Internet Research*, 10, e44.
- *Wright, J. H., Wright, A. S., Albano, A. M., Basco, M. R., Goldsmith, L. J., Raffield, T., et al. (2005). Computer-assisted cognitive therapy for depression: Maintaining efficacy while reducing therapist time. *American Journal of Psychiatry*, 162, 1158–1164.