



EuCRUST-07: A new reference model for the European crust

Magdala Tesauro,^{1,2} Mikhail K. Kaban,^{2,3} and Sierd A. P. L. Cloetingh¹

Received 8 October 2007; revised 29 November 2007; accepted 12 December 2007; published 14 March 2008.

[1] We present a new digital model (EuCRUST-07) for the crust of Western and Central Europe and surroundings (35°N–71°N, 25°W–35°E). Available results of seismic reflection, refraction and receiver functions studies are assembled in an integrated model at a uniform grid (15' × 15'). The model consists of three layers: sediments and two layers of the crystalline crust. Besides depth to the boundaries, we provide average *P*-wave velocities in the upper and lower parts of the crystalline crust. The new model demonstrates large differences in the Moho depth compared to previous compilations, over ±10 km in some specific areas (e.g. the Baltic Shield). Furthermore, the velocity structure of the crust is much more heterogeneous than in previous maps. EuCRUST-07 offers a starting point for numerical modeling of deeper structures by allowing correction for crustal effects beforehand and to resolve trade-off with mantle heterogeneities. **Citation:** Tesauro, M., M. K. Kaban, and S. A. P. L. Cloetingh (2008), EuCRUST-07: A new reference model for the European crust, *Geophys. Res. Lett.*, 35, L05313, doi:10.1029/2007GL032244.

1. Introduction and Basic Model Assumptions

[2] The crust is the most heterogeneous layer in the Earth and its impact to the interpretation of deep structures can mask the effect of deep seated heterogeneities. It is, for instance, nearly impossible to separate the crustal and mantle effects in potential field and geothermal modeling without additional data on the crustal structure [e.g., Kaban *et al.*, 2004; Artiemeva, 2006; Artiemeva *et al.*, 2006]. It is still very difficult to minimize the trade-off between the crustal and upper mantle heterogeneities in seismic tomography, which remains the main tool to investigate the structure of the mantle. Martin *et al.* [2005] have reported that teleseismic tomography for SE Romania without applying sophisticated a priori 3-D crustal correction gives results, which are strongly contaminated in the uppermost 100–150 km depth by a significant effect from an incorrect crustal model. Waldhauser *et al.* [2002] have demonstrated that the non-linear inversion of the synthetic residuals without correcting for the 3-D crustal structure erroneously maps the crustal anomalies into the upper mantle. Therefore, reliable models of the upper mantle can be constructed only if the effect of the crust is reduced from the observed fields beforehand. Crustal models primarily based on existing reflection and refraction seismic profiles have been used

for these purposes during the last decade. However, different models of the European crust are still inconsistent in many respects. In particular, differences of the existing Moho maps often reach and even exceed ±15 km (e.g. CRUST5.1 [Mooney *et al.*, 1998], CRUST2.0 [Bassin *et al.*, 2000] and SVEKALAPKO [Kozlovskaya *et al.*, 2004]). These discrepancies result from the coarser resolution of previous compilations (e.g. CRUST 2.0), which neglect small scale features, and differences of the employed data sets, in particular based on non-seismic information. Consequently, the obtained results after corrections for crustal structure are different in many cases. For example, the residual mantle gravity anomalies obtained by different authors may differ up to about 100 mGal [e.g. Yegorova and Starostenko, 1999; Kaban *et al.*, 2004].

[3] The main purpose of this study is to construct a new high-resolution 3-D model (EuCRUST-07) of the European crust, which can be used as a starting point in a wide range of lithosphere and upper mantle studies. EuCRUST-07 is mainly based on existing seismic refraction, reflection and receiver functions studies, most of them carried out within the ILP program (Figure 1). Whenever possible we use detailed local compilations based on seismic data [e.g. Kozlovskaya *et al.*, 2004]. We refrain from using inferences from non-unique gravity field interpretations commonly employed in previous compilations. The study area is limited to 35°N–71°N, 25°W–35°E. EuCRUST-07 consists of three layers: sediments and 2 layers of the crystalline crust, the latter characterized by an average *P*-wave velocity determined from seismic data. Depth to the crystalline basement and Moho are the parameters most reliably determined in all kinds of seismic data. The situation with the inner crustal boundaries is more complicated. At least two layers within the crystalline crust are detected in most seismic sections; therefore, it has been decided to maintain this division in the generalized model. In the areas, where the crystalline crust consists of only one layer, having a constant velocity (e.g. in the Tyrrhenian Sea) or characterized by a gradual change (e.g. in the Western part of the Black Sea), we arbitrarily divide the crust in two layers of equal thickness having average velocities consistent with the seismic data. In the opposite case, several layers are joint to form one equivalent layer, e.g. in the East European Platform (EEP), where the velocity in the upper layer is calculated as a weighted average of upper and middle crust velocities. The final model (Data Set S1 of the auxiliary material)¹ is presented on a uniform 15' × 15' grid.

[4] The employed local Moho maps are shown in Figure 1. When possible, we have verified these compilations and modified them in some details using available seismic profiles (Data Set S2 of the auxiliary material). All

¹Netherlands Research Centre for Integrated Solid Earth Science, Faculty of Earth and Life Sciences, Vrije Universiteit, Amsterdam, Netherlands.

²GeoForschungsZentrum Potsdam, Potsdam, Germany.

³Institute for Physics of the Earth, Moscow, Russia.




Figure 1. Moho depth (km) updated from Ziegler and Dèzes [2006] (34°N – 62°N , 18°W – 25°E) and extended (35°N – 71°N , 25°W – 35°E) including an array of new datasets. Dashed lines show the location of the seismic profiles incorporated.

the maps have been converted to the same resolution $15' \times 15'$. In the regions densely covered by recent seismic data, not included in the existing Moho compilations (e.g. in the Iberian Peninsula, where the results of the most recent seismic experiments are not incorporated in the Ziegler and Dèzes [2006] map, ZDmap) we interpolated the seismic profiles using a standard kriging method (SURFER, Golden Software package) to trace the Moho boundary. We merged all the compilations in a unified model giving a preference to the most robust. Typically we used detailed local studies, which are based on well specified sources. The regional ZDmap has been largely substituted by local compilations. For the part of the oceans not covered by seismic profiles, we assign Moho depths from the global model CRUST2.0 [Bassin et al., 2000], for a part of Norway from the Geothermal Atlas of Europe [Hurtig et al., 1992] and for a part of the EEP and the Black Sea from the compilation of Kaban [2001]. To produce a smooth transition between different maps we left some free space between the compilations. The size of the gaps corresponds to their resolution (from 30 km for the detailed maps to 100 km for the global CRUST2.0). The same kriging technique was used to interpolate the gaps. The velocity distributions within the crys-

talline crust layers are mostly based on the interpolated determinations (wide-angle seismic data), whereas for most of Fennoscandia we used the model of Kozlovskaya et al. [2004]. In several cases when the non-uniform data coverage does not allow for a robust interpolation, we increased the number of data by adding extra points in agreement with the position of local tectonic units with reliable determinations in other parts (e.g. in Norway). In the oceanic domain without seismic data we assigned P -wave velocities (V_p) of 5.5 and 6.75 km/s to the upper and lower crust, correspondingly. The final grid was also obtained using the kriging interpolation.

[5] The depth to basement is determined based on available maps and sedimentary thickness compilations (Figure S1 of the auxiliary material). These compilations were verified and in several areas modified (e.g. in the North German Basin and in the Adriatic Sea) using the seismic profiles collected. We do not specify a velocity structure of the sediments. Due to extremely strong heterogeneity (both lateral and vertical) of this layer it is difficult to integrate relatively sparse published data into a uniform model. On the other hand, the material properties of sediments (e.g. density) are much less related to velocity variations, while seismic tomography results are mostly biased by crystalline




Figure 2. Moho depth (km). Abbreviations are as follows: A, Apennines; AB, Alboran Basin; AP, Adriatic Promontory; BC, Betic Cordillera; BS, Black Sea; CH, Carpathians; CM, Cantabrian Mountain; D, Dinarides; EB, Eoras Bank; EL, ElbeLineament; EEP, East European Platform; FB, Focsani Basin; FI, Faeroe Islands; GB, Gulf of Bothnia; HB, Hatton Bank; IAP, Iberian Abyssal Plain; IS, Iapetus Suture; LVM, Lofoten–Vesterålen margin; MC, Massif Central; NGB, North German Basin; NS, North Sea; OR, OsloRift; P, Pyrenees; PB, Pannonian Basin; TS, Tyrrhenian Sea; TTZ, Tesseyre–Tornquist zone; URG, Upper Rhine Graben; VB, Vøring Basin; VT, Valencia Trough.

crust heterogeneity. Therefore, we leave this problem for future regional and detailed studies.

[6] Below we highlight some principal features of EuCRUST-07 for the main tectonic units of Europe.

2. Crustal Model of Western and Central Europe

[7] The new Moho map and the average velocities in the upper and lower crust are shown in Figures 2, 3, and 4 respectively. In agreement with previous studies systematic differences in the crustal thickness and velocity between Western and Eastern Europe are found. However, we also




Figure 3. Average P -wave velocity in the upper crust (km/s).

observe a very strong heterogeneity of the crustal structure within these domains.

2.1. Fennoscandia

[8] The Moho depression beneath Central and Southern Finland and the Gulf of Bothnia is much more pronounced than in previous compilations (>60 km, 10 km deeper compared to CRUST2.0). This is the result of recent seismic experiments (SVEKALAPKO [Kozolovskaya *et al.*, 2004]). The boundary tends to rise to the west reaching a value of ~ 40 km in Sweden and of about 30–32 km beneath the Oslo rift and the Norwegian coast. Two local maxima of 40–43 km have been recently detected in southern Norway by a receiver functions study [Svenningsen *et al.*, 2007]. Westward to the Norwegian coast the Moho gradually shallows to ~ 20 km, while it deepens again to ~ 25 km at the transition to the oceanic crust in the Lofoten–Vesterålen margin and Vøring Basin, which is characterized by a thick, high velocity (>7.0 km/s) lower crust [e.g., Mjælde *et al.*, 2005]. The maximum velocity observed in this area (over 8.0 km/s) is interpreted as a deep crustal root of partially eclogitized rocks that formed during the Caledonian orogeny [Raum *et al.*, 2006]. Northward, in the Lofoten–Vesterålen margin, a decrease in the thickness of the lower crust is accompanied by a decreasing amount of breakup intrusives and extrusives. The crust in this area experienced only moderate extension, in contrast to the occurrence of major crustal extension in the southern Vøring margin [e.g. Tsikalas *et al.*, 2005].

2.2. EEP and Tesseyre-Tornquist Zone (TTZ)

[9] The Moho discontinuity is relatively uniform within the considered part of the EEP (40 to 50 km) and the crust is characterized by high P -wave velocities (~ 7.0 km/s) in the lower part except for some areas in the southwestern edge, where relatively low velocities (~ 6.75 km/s) are observed. By contrast, the upper crust is quite inhomogeneous, being characterized by alternation of low and high velocity zones [Grad *et al.*, 2006] with an average value of V_p of 6.3 km/s. The Moho depth beneath the TTZ in Central Poland is intermediate between that in the EEP to the east (~ 45 km) and the Palaeozoic Platform to the south-west (30–35 km). The upper crust is characterized by low velocities, reflecting




Figure 4. Average P -wave velocity in the lower crust (km/s).

a presence of metamorphic sediments or volcanic strata with $V_p < 6.0$ km/s, which depth reaches 18–20 km. On the other hand, velocities in the lower crust in this area are very similar to that of the EEP (~ 7.0 km/s) [Guterch *et al.*, 2006]. The crustal structure is more heterogeneous to the west from the TTZ. It is characterized by Variscan crust (e.g. in France) with thickness of about 30–35 km and slow V_p in the lower crust (~ 6.4 – 6.8 km/s), orogens with deep Moho (45–50 km) and extensional areas with very shallow Moho (e.g. ~ 10 km beneath the Tyrrhenian Sea) and low lower crust velocity (~ 6.0 km/s).

2.3. England and Atlantic Margin

[10] Beneath England and Ireland the Moho depth is between 30 and 35 km [Kelly *et al.*, 2007] and the velocity in the upper crust is between 6.0 and 6.4 km/s, while it increases in the lower crust from 6.7 to 7.0 km/s south of the Iapetus Suture Zone [Landes *et al.*, 2005]. The Moho shallows southwest of the Irish coast from 27 to less than 20 km over a horizontal distance of ~ 100 km, while the velocity decreases both in the upper and in the lower crust to 6.0 and 6.5 km/s, correspondingly. West of the Irish coast the continental crust thickness varies between 12 and 25 km [Kelly *et al.*, 2007], on account of the different amount of stretching to which it was subjected during the opening of the Atlantic Ocean [e.g. O'Reilly *et al.*, 1995]. The difference with CRUST2.0 is up to 15 km in this area. High velocities in the lower crust (~ 7.15 km/s) are observed beneath the Edoras and the Hatton Bank, presumably due to mantle underplating. The Moho deepens to 37–38 km beneath the Faeroe Islands and to 40–42 km approximately 100 km west of them, while the velocity in the lower crust is about 7.0 km/s. These high crustal thickness values (~ 10 km larger than in CRUST2.0) are a consequence of the Faeroe Islands location above the hottest core of the Icelandic mantle plume at the time of continental breakup [Richardson *et al.*, 1998].

2.4. North German Basin

[11] The Elbe Lineament, like the TTZ, represents another important geological boundary, where strong changes in the crustal thickness and composition are observed. The Moho beneath the North German Basin shallows from 35–34 km to 28–25 km from north to south of the Elbe Lineament, while V_p in the upper and lower crust decreases from 6.1 to 5.9 km/s and from 7.0 to 6.3 km/s, respectively. This strong change in the velocity of the lower crust possibly reflects a compositional transition from mafic to granitic rocks [e.g. Scheck *et al.*, 2002]. In the southeastern part of the North Sea similar changes in velocity and crustal thickness are present, suggesting that the WNW–ESE-striking Elbe Lineament continues into the Southern North Sea [e.g. Scheck *et al.*, 2002].

2.5. Carpathians-Pannonian Basin System

[12] The Carpathians, like the Alps, show strong changes in the crustal thickness, being characterized by a Moho depth of over 50 km in the Eastern part, which decreases to about 35 km in the Western part [Horváth *et al.*, 2006]. In the Southern Carpathians the Moho is between 37 and 42 km and reaches a maximum of ~ 44 km in the Focsani Basin [Martin *et al.*, 2005]. These recent determinations exceed values from the ZDmap for more than 10 km. The two layers of the crystalline crust areas are characterized by

P -wave velocities of ~ 6.1 and 6.6–6.9 km/s respectively [e.g. Sroda *et al.*, 2006]. The Pannonian Basin, which experienced strong extension, is characterized by a shallow Moho (~ 25 km) [Horváth *et al.*, 2006] and low upper (~ 6.0 km/s) and lower (~ 6.35 km/s) crust velocities.

2.6. Alps

[13] The Alpine region displays one of the main changes with previous compilations. Strong variations of the Moho depth, observed in recent seismic experiments [Finetti, 2005], are reflected in EuCRUST-07. Beneath the Western and Eastern Alps the European Moho plunges southward to ~ 40 and ~ 55 km, respectively. On the Adriatic side the Moho depth reaches 45 km beneath the Eastern Alps, while beneath the Western Alps a fragment of mantle-like material is observed, which is imbricated into the Alpine crust at the Insubric line [Finetti, 2005]. Therefore, the Moho depth is significantly reduced here (~ 20 km compared to the ZDmap) and the lower crust velocity is increased (~ 7.3 km/s). In the other parts of the Alpine chain the velocity of the upper crust shows normal values (6.0–6.1 km/s), while in the lower crust is anomalously low (6.4–6.6 km/s) [Scarascia and Cassinis, 1997].

2.7. Iberian Peninsula

[14] In the Iberian Peninsula the maximum depth of the Moho discontinuity is found beneath the Pyrenees and Cantabrian area (~ 45 km), significantly deeper than in previous maps (e.g. ZDmap), reflecting the subduction of the Iberian plate. In the same areas relative high velocities are found in the upper (6.25 km/s) and lower crust (from 6.7 to 7.20 km/s) [Pedreira *et al.*, 2003]. Strong changes in the Moho depth are also observed in the southeastern part of the Iberian Peninsula, where it shallows abruptly from 38 km beneath the Betics to 15 km beneath the Eastern part of the Alboran Basin [e.g. Fullea *et al.*, 2007]. In the same units a velocity decrease from 6.7 to 6.3 km/s in the lower crust is found. From the Western Iberian coast to the Iberian Abyssal Plain, the Moho shallows from 30 to 13 km over a horizontal distance of 300 km, showing a transition from continental to oceanic crust. In the Abyssal Plain the velocity in the basaltic lavas, representing the upper crust, decreases to 5.0 km/s, while in the highly serpentinized lower crust it increases up to ~ 7.5 km/s [Chian *et al.*, 1999].

3. Conclusions

[15] We have presented a new digital model of the crust for Western and Central Europe and surroundings that can be used as a constraint in modeling lithosphere and upper mantle structures. EuCRUST-07 demonstrates large differences with previous compilations, mostly resulting from including recently acquired seismic data. The discrepancies with the most recent regional Moho map of Ziegler and Dèzes [2006] reach -25 km (rms = 3.4 km), in the Ivrea zone, where the new map shows the updoming of a 10–20 km shallower Adriatic Moho, and $+17$ km in the Eastern Alps. Under the Cantabrian Mountains the new Moho is 8–10 km deeper, reflecting the subduction of the Iberian plate. Differences with the recent global map of Kaban [2001] reach -14 – $+17$ km (rms = 3.9 km). Furthermore, when we average the local variations of EuCRUST-07 to bring it to the resolution of CRUST2.0, the difference remains signif-

icant from -19 km, in the Atlantic margin, to $+11$ km, in the Faeroe Islands and the Baltic Shield (rms = 4 km). The velocity structure of the crust turns out to be rather heterogeneous, even without considering regional variations from Western to Eastern Europe and the Baltic Shield, which are basically shown in previous models (e.g. CRUST2.0). Small scale features (e.g. ultramafic lower crust in the Vøring basin), revealed in EuCRUST-07 but not considered in the global models, also substantially influence the averaged values. The overall range of V_p variations in the upper part of the crystalline crust is 5.0 – 6.7 km/s and 6.0 – 8.4 km/s for the lower crust.

[16] **Acknowledgments.** We would like to thank P. Dèzes and P. Ziegler (University of Basel), R. W. England (University of Leicester), E. Kozlovskaja (University of Oulu), F. Marone (University of California), M. Martin and J. Ritter (University of Karlsruhe), L. Matias (University of Lisboa), F. Sodoudi and R. Kind (GFZ, Potsdam), for providing Moho data. We are grateful to C. Ayala (IGME), O. Bourgeois (Nantes University), P. Ledru (BRGM), T. Dìhel (ETH), L. Lenkey (Eötvös University) and M. Scheck-Wenderoth (GFZ, Potsdam), for providing sedimentary thickness and basement depth compilations. Valuable comments from the Editor Eric Calais and two anonymous reviewers improved the original manuscript. Funds were kindly provided by NWO (Netherlands Organization for Scientific Research) and SRON (Space Research Organization Netherlands).

References

- Artemieva, I. M. (2006), Global $1^\circ \times 1^\circ$ thermal model TC1 for the continental lithosphere: implications for lithosphere secular evolution, *Tectonophysics*, *416*, 245–277.
- Artemieva, I. M., H. Thybo, and M. Kaban (2006), Deep Europe today: Geophysical synthesis of the upper mantle structure and lithospheric processes over 3.5 By, in *European Lithosphere Dynamics*, edited by D. G. Gee and R. A. Stephenson, *Geol. Soc. Mem.*, *32*, 11–41, Geol. Soc., London.
- Bassin, C., G. Laske, and G. Masters (2000), The current limits of resolution for surface wave tomography in North America, *Eos Trans American Geophysical Union*, *81*(48), Fall Meet. Suppl., Abstract F897.
- Chian, D., K. F. Loudon, T. Minshull, and R. Whitmarsh (1999), Deep structure of the ocean-continent transition in the southern Iberian Abyssal Plain from seismic refraction profiles: Ocean Drilling Program (Legs 149 and 173) transect, *J. Geophys. Res.*, *104*(B4), 7443–7462.
- Finetti, I. R. (2005), Depth contour Map of the Moho discontinuity in the Central Mediterranean region from new CROP seismic data, in *CROP Project, Deep Seismic Exploration of the Central Mediterranean and Italy*, edited by I. R. Finetti, pp. 597–606, Elsevier, New York.
- Fullea, J., M. Fernández, H. Zeyen, and J. Vergés (2007), A rapid method to map the crustal and lithospheric thickness using elevation, geoid anomaly and thermal analysis. Application to the Gibraltar Arc System, Atlas Mountains and adjacent zones, *Tectonophysics*, *430*, 97–117.
- Georgiev, G., C. Dabovski, and G. Stanisheva-Vassileva (2002), East Srednogie-Balkan Rift Zone, in *Peri-Tethys Memoir*, vol. 6, *Peri-Tethyan Rift/Wrench Basins and Passive Margins*, *Mem. Mus. Natl. His. Nat.*, vol. 186, edited by P. A. Ziegler et al., pp. 259–293, Mus. du Natl. His. Nat., Paris.
- Grad, M., et al. (2006), Lithospheric structure of the western part of the East European Craton investigated by deep seismic profiles, *Geol. Q.*, *50*(1), 9–22.
- Guterch, A., and M. Grad (2006), Lithospheric structure of the TESZ in Poland based on modern seismic experiments, *Geological Quarterly*, *50*(1), 23–32.
- Horváth, G., F. Bada, P. Szafián, G. Tari, A. Adám, and S. Cloetingh (2006), Formation and deformation of the Pannonian Basin: Constraints from observational data, in *European Lithosphere Dynamics*, *Geol. Soc. Mem.*, vol. 32, edited by D. Gee and R. Stephenson, pp. 191–206, Geol. Soc., London.
- Hurtig, E., V. Cermak, R. Haenel, and V. Zui (1992), *Geothermal Atlas of Europe, International Association for Seismology and Physics of the Earth's Interior*, Hermann Haack Verlagsgesellschaft mbH-Geographisch-Kartographische Anstalt, Gotha, Germany.
- Kaban, M. K. (2001), A gravity model of the north Eurasia crust and upper mantle: 1. Mantle and isostatic residual gravity anomalies, *Russ. J. Earth Sci.*, *3*(2), 143–163.
- Kaban, M. K., O. G. Flovenz, and G. Palmason (2002), Nature of the crust-mantle transition zone and the thermal state of the upper mantle beneath Iceland from gravity modelling, *Geophys. J. Int.*, *149*, 281–299.
- Kaban, M. K., P. Schwintzer, and C. Reigber (2004), A new isostatic model of the lithosphere and gravity field, *J. Geod.*, *78*, 368–385.
- Kelly, A., R. W. England, and P. K. H. Maguire (2007), A crustal seismic velocity model for the UK, Ireland and surrounding seas, *Geophys. J. Int.*, *171*, 1172–1184.
- Kozlovskaya, E., et al. (2004), 3-D density model of the crust of southern and central Finland obtained from joint interpretation of the SVEKALAPKO crustal P-wave velocity models and gravity data, *Geophys. J. Int.*, *158*, 827–848.
- Landes, M., J. R. R. Ritter, P. W. Readman, and B. M. O'Reilly (2005), A review of the Irish crustal structure and signatures from the Caledonian and Variscan Orogenies, *Terra Nova*, *17*, 111–120.
- Lassen, A. (2005), Structure and evolution of the pre-Permian basement in the Danish Area, Ph.D. thesis, 130 pp., Copenhagen Univ., Copenhagen.
- Marone, F., M. van der Mijede, S. vander Lee, and D. Giardini (2003), Joint inversion of local, regional and teleseismic data for crustal thickness in the Eurasia-Africa plate boundary region, *Geophys. J. Int.*, *154*, 499–514.
- Martin, M., et al. (2005), High-resolution teleseismic body-wave tomography beneath SE Romania: part I. Implications for three-dimensional versus one-dimensional crustal correction strategies with a new crustal velocity model, *Geophys. J. Int.*, *162*, 448–460.
- Mjelde, R., T. Raum, B. Myhren, H. Shimamura, Y. Murai, T. Takanami, R. Karpuz, and U. Naess (2005), Continent-ocean transition on the Vøring Plateau, NE Atlantic, derived from densely sampled ocean bottom seismometer data, *J. Geophys. Res.*, *110*, B05101, doi:10.1029/2004JB003026.
- Mooney, W. D., G. Laske, and G. Masters (1998), CRUST 5.1: A global crustal model of 5×5 , *J. Geophys. Res.*, *103*, 727–747.
- O'Reilly, B. M., F. Hauser, A. W. B. J. P. M. Shannon, J. Makris, and U. Vogt (1995), The transition between the Erris and the Rockall basins: new evidence from wide-angle seismic data, *Tectonophysics*, *241*, 143–163.
- Pedreira, D., J. A. Pulgar, J. Gallart, and J. Díaz (2003), Seismic evidence of Alpine crustal thickening and wedging from the western Pyrenees to the Cantabrian Mountains (north Iberia), *J. Geophys. Res.*, *108*(B4), 2204, doi:10.1029/2001JB001667.
- Raum, T., R. Mjelde, H. Shimamura, Y. Murai, E. Bråstein, R. M. Karpuz, K. Kravik, and H. J. Kolstø (2006), Crustal structure and evolution of the southern Vøring Basin and Vøring Transform Margin, NE Atlantic, *Tectonophysics*, *415*, 167–202.
- Richardson, K. R., J. R. Smallwood, R. S. White, D. B. Snyder, and P. K. H. Maguire (1998), Crustal structure beneath the Faeroe Islands and the Faeroe-Iceland Ridge, *Tectonophysics*, *300*, 159–180.
- Scarcasia, S., and R. Cassinis (1997), Crustal structures in the central-eastern Alpine sector: a revision of the available DSS data, *Tectonophysics*, *271*, 157–188.
- Scheck, M., U. Bayer, V. Otto, J. Lamarche, D. Banka, and T. Pharaoh (2002), The Elbe Fault System in North Central Europe—a basement controlled zone of crustal weakness, *Tectonophysics*, *360*, 281–299.
- Sodoudi, F., R. Kind, D. Hatzfeld, K. Priestley, W. Hanka, K. Wylegalla, G. Stavrakakis, A. Vafidis, H.-P. Harjes, and M. Bohnhoff (2006), Lithospheric structure of the Aegean obtained from P and S receiver functions, *J. Geophys. Res.*, *111*, B12307, doi:10.1029/2005JB003932.
- Šroda, P., W. Czuba, M. Grad, A. Guterch, A. K. Tokarski, T. Janik, M. Rauch, G. R. Keller, E. Hegedus, J. Vozar, and CELEBRATION 2000 Working Group (2006), Crustal and upper mantle structure of the Western Carpathians from CELEBRATION 2000 profiles CEL01 and CEL04: seismic models and geological implications, *Geophys. J. Int.*, *167*, 737–760.
- Svenningsen, L., et al. (2007), Crustal root beneath the highlands of Southern Norway resolved by teleseismic receiver functions, *Geophys. J. Int.*, *170*(3), 1129–1138.
- Tsikalas, F., T. O. Eldholm, and J. I. Faleide (2005), Crustal structure of the Lofoten–Vesterålen continental margin, off Norway, *Tectonophysics*, *404*, 151–174.
- Waldhauser, F., R. Lippitsch, E. Kissling, and J. Ansgore (2002), High-resolution teleseismic tomography of upper mantle structure using an a priori 3D crustal model, *Geophys. J. Int.*, *150*, 141–403.
- Yegorova, T. P., and V. I. Starostenko (1999), Large-scale three-dimensional gravity analysis of the lithosphere below the transition zone from Western Europe to the East European Platform, *Tectonophysics*, *314*, 83–100.
- Ziegler, P. A., P. Dèzes (2006), Crustal evolution of western and central Europe, in *European Lithosphere Dynamics*, edited by D. G. Gee and R. A. Stephenson, *Geol. Soc. Mem.*, vol. 32, pp. 43–56 Geol. Soc., London.

S. A. P. L. Cloetingh and M. Tesauro, Netherlands Research Centre for Integrated Solid Earth Science, Faculty of Earth and Life Sciences, Vrije Universiteit, De Boelelaan, Amsterdam NL-1081 HV, Netherlands. (sierd.cloetingh@falw.vu.nl; magdala.tesauro@falw.vu.nl)

M. K. Kaban, Geoforschungszentrum Potsdam, Telegrafenberg A17, Department 1.3, D-14473 Potsdam, Germany. (Kaban@gfz-potsdam.de)