




Lidar characterizations of atmospheric aerosols and clouds


Richard Anthony Ferrare¹, Chris A Hostetler¹, Johnathan W Hair², Sharon P Burton¹ and NASA LaRC HSRL Team, (1)NASA Langley Research Center, Hampton, VA, United States, (2)NASA Langley, Hampton, VA, United States

NASA Langley Research Center, Hampton VA



CALIPSO observations of smoke over low clouds



CALIPSO clearly demonstrated the necessity of vertically-resolved measurements of clouds and aerosols for climate, weather, and air quality applications.




Elastic Backscatter Lidar Measurements



$$P_a(r) = \frac{C}{r^2} \left[\beta_m(r) + \beta_a(r) \exp \left\{ -2 \int_0^x [\alpha_m(r') + \alpha_a(r')] dr' \right\} \right]$$

2 unknowns



attenuated backscatter



Problem:


- Lidars measure the attenuated backscatter coefficient
- What we actually want is
 - The true aerosol backscatter coefficient
 - And the aerosol extinction coefficient
- To get these, we must somehow correct for extinction as the retrieval proceeds downward through the profile
 - This correction can have significant errors

High Spectral Resolution Lidar (HSRL) Technique



- Tuning laser to an I_2 absorption line blocks particulate backscatter from “Molecular Channel”
- 2 channels enable 2 equations to solve for 2 unknowns: aerosol backscatter and extinction

Why use HSRL for aerosol research?





2 equations

$$P_a(r) = \frac{C}{r^2} [\beta_m(r) + \beta_a(r)] \exp \left\{ -2 \int_0^x [\alpha_m(r') + \alpha_a(r')] dr' \right\}$$


$$P_m(r) = \frac{C}{r^2} [\beta_m(r)] \exp \left\{ -2 \int_0^x [\alpha_m(r') + \alpha_a(r')] dr' \right\}$$

2 unknowns

- **Backscatter coefficient**
 - Direct measure of backscatter, rather than attenuated
 - Accurate at all altitudes; errors do not accumulate with range
- **Independent measure of extinction**
 - No need for inferred/modeled lidar ratio or external constraint
 - Molecular channel also provides direct measure of AOT
- **Highly accurate particulate depolarization**
 - Separating particulate and molecular parts requires accurate backscatter
- **Vertically resolved aerosol type information**
 - Lidar ratio gives the most information about aerosol composition for non-dust aerosol



Why use HSRL for aerosol research?





2 equations

$$P_a(r) = \frac{C}{r^2} [\beta_m(r) + \beta_a(r)] \exp \left\{ -2 \int_0^x [\alpha_m(r') + \alpha_a(r')] dr' \right\}$$

$$P_m(r) = \frac{C}{r^2} [\beta_m(r)] \exp \left\{ -2 \int_0^x [\alpha_m(r') + \alpha_a(r')] dr' \right\}$$

2 unknowns


- **Backscatter coefficient**
 - Direct measure of backscatter, rather than attenuated
 - Accurate at all altitudes; errors do not accumulate with range
- **Independent measure of extinction**
 - No need for inferred/modeled lidar ratio or external constraint
 - Molecular channel also provides direct measure of AOT
- **Highly accurate particulate depolarization**
 - Separating particulate and molecular parts requires accurate backscatter
- **Vertically resolved aerosol type information**
 - Lidar ratio gives the most information about aerosol composition for non-dust aerosol



CALIPSO misses much of the Arctic Aerosol





April 19, 2008 (ARCTAS)




- **CALIPSO sees major aerosol transport events (dust, smoke)**
- **CALIPSO misses background Arctic aerosols below detection threshold**


Aerosol Backscatter



CALIPSO does not detect all radiatively significant aerosol



- CALIPSO aerosol detection sensitivity was compared to ground-based DOE ARM Raman lidars and LaRC airborne HSRL data
- CALIPSO underestimates global-mean AOD, and therefore shortwave aerosol direct radiative effect, by about 22-47% (38-54%) in cloud-free (transparent profiles) (Thorsen and Fu, 2015)
 - Similar sensitivities as other studies (Rogers et al., 2014)
 - Undetected AOT of 0.031 ± 0.052 (Kim et al; 2016)
- CALIPSO's performance is as expected (consistent with ATBD)



Thorsen et al., 2017

HSRL-DIAL group developed and fields three (soon to be four) airborne lidars that employ HSRL technique



HSRL-1
Aerosols, clouds, ocean
2004 – Present




HSRL-2
Aerosols, clouds, ozone,
ocean
2012 - Present



UV DIAL/HSRL
O₃ and aerosols
1983 - Present

Prototype for the
spaceborne lidar on the
ACE Decadal Survey
mission

Coming soon: HALO
Methane, water
vapor, aerosols,
clouds, ocean
2018 -





Technology
development and
demonstration for future
space methane and
water vapor lidar

Comparisons with AERONET provide high confidence in HSRL-2 extinction product






HSRL 0-7 km layer AOD values compare well with column AOD (355 and 532 nm) values from AERONET "DRAGON" stations when HSRL was within 2.5 km of site and 10 minutes from measurement



Airborne NASA LaRC HSRL-2 measuring smoke distribution and properties for model evaluation during ORACLES



- ORACLES Earth Venture Suborbital mission
- Target: Extensive biomass smoke plume over persistent stratus cloud deck off the west coast of Africa
 - Smoke has significant radiative effect: localized absorption and impacts cloud microphysics
- During first ORACLES mission, GEOS-5 model smoke plume was systematically lower than HSRL-2 observations (da Silva - GSFC)

HSRL-2 Aerosol Backscatter (532 nm)



HSRL-2 characterization of smoke layer

- HSRL-2 aerosol backscatter measurements characterize:
 - smoke and cloud layer heights and thicknesses
 - thickness of the aerosol-free layer between the smoke and cloud layers (i.e. the “gap”)
- Much (>40%) of the time, there is no gap







Assessing Aerosol Data Assimilation Products Using HSRL Measurements





Mean Aerosol Extinction Profiles

- HSRL aerosol extinction profiles are being used to assess and improve aerosol data assimilation systems (NASA GEOS-5 and MERRAero, ECMWF/MACC-III)
- With exception of California San Joaquin Valley, MERRAero mean aerosol extinction profiles are in general agreement with HSRL





HSRL technique enables identification of aerosol type and apportioning optical depth by type



Intensive observables:


- Independent of aerosol amount
- Depends only on aerosol type

Burton et al., 2012, AMT


Spectral dependence of particle depolarization differs for smoke and dust




Smoke depolarization decreases with wavelength



Dust depolarization increases with wavelength



Spectral depolarization reveals information about particle size distribution




**3 case studies from HSRL-2 (3 wavelength)
+ 3 from HSRL-1 (2 wavelength)**

Burton et al., ACP, 2015

- Wavelength dependence of particle depolarization reveals information about particle size
- North American dust at the source includes very large particles, monotonically increasing depolarization
- Transported Saharan dust cases peak at mid-wavelength, largest particles were lost during transport
- Non-spherical smoke particles (coated soot aggregates) have decreasing wavelength dependence, smaller particles
- 355 nm particulate depolarization alone (ATLID) not sufficient for separating dust and smoke

Multiwavelength HSRL-2 retrievals characterize aerosol concentration and size during ORACLES



Multiwavelength lidar retrieval algorithms (Müller et al, 1999; Veselovskii et al. 2002; etc)

lidar measurements

particle size distribution

INVERSION


$$\beta(\lambda) = \int K_{\beta}(r, m, \lambda) v(r) dr$$

$$\alpha(\lambda) = \int K_{\alpha}(r, m, \lambda) v(r) dr$$


size, refractive index, wavelength

$3\beta+2\alpha$ (i.e. 3 backscatter + 2 extinction) considered the minimum information content necessary for microphysical retrievals (Bockmann et al, 2005)

Multiwavelength HSRL-2 retrievals characterize aerosol concentration and size during ORACLES






- Microphysical parameters retrieved and archived:
 - Concentration (fine and total) (number, surface, volume)
 - Effective radius (fine and total)
 - Small mode fraction



Output
(Preliminary)

Multiwavelength HSRL aerosol retrieval profiles compare well to coincident airborne in situ measurements



- Relevance of particle size:
 - Improve parameterizations in aerosol transport models
 - Modeling direct radiative effects
 - Indirect effect on cloud radiative properties and precipitation (CCN)

Lidar microphysical retrievals of effective radius and concentrations compare well to airborne in situ measurements

- Relevance of particle concentration:
 - Indirect effects (CCN)
 - Air quality ($\text{PM}_{2.5}$)



Pursuing combined lidar+polarimeter retrievals

- Limits in information content for lidar-only microphysical retrieval means absorption is retrieved less well than size distribution parameters
- Synergistic combination of active (lidar) and passive (polarimeter) measurements will optimize information content on vertical profile of absorption properties

Lidar

- vertically resolved measurements
- multi-wavelength backscatter and extinction coefficients
- good accuracy for size distribution
- less accuracy for absorption

Polarimeter




- multiwavelength, multiangle
- polarized radiances
- good accuracy for absorption
(e.g. ± 0.02 on SSA)
- limited information on vertical profile

Lidar + Polarimeter

- vertically resolved profiles of effective radius, concentrations and complex refractive index

- Ongoing project by Xu Liu et al (NASA Langley) to combine HSRL-2 (lidar) and RSP (polarimeter) measurements in advanced Optimal Estimation retrieval

The unexpected application: CALIPSO ocean measurements



- Particulate Organic Carbon (POC) retrieved from CALIOP spaceborne lidar compared favorably to the MODIS product
- CALIOP retrievals pioneered by Yongxiang Hu
- CALIOP retrievals are vertically-integrated (i.e., not vertically-resolved) due to coarse vertical resolution


From supplementary material for: Behrenfeld et al., Space-based lidar measurements of global ocean carbon stocks, GRL, 2013.

Data in each panel are climatological annual averages for the 2006 to 2012 period. Data are binned to 2° latitude by 2° longitude pixels.

HSRL-1 Measurements of Atmospheric and Oceanic Particulates




Aerosol Backscatter ($\text{Mm}^{-1}\text{sr}^{-1}$)(532nm)
0.01 0.1 1 10




Dense smoke layers make passive 'ocean color' retrievals impossible

HSRL retrieval immune to overlying aerosol 'interference'. Provides depth-resolved profile



Why do we care about depth-resolved profiles?



Net Primary Productivity (NPP)

- NPP = rate at which CO_2 is converted to biomass via photosynthesis
- Phytoplankton fix as much carbon as all the world's terrestrial vegetation – an critical component of the carbon cycle
- Phytoplankton are the base of the food web

Problem: ocean color measurements are sensitive to only the first few meters of the column

- Assumption of vertical homogeneity required to estimate ecosystem properties from ocean color data
- But, this assumption is often violated
- Results in large errors in fundamental quantities such as NPP

At left are data from the SABOR mission off East Coast US (2014) demonstrating differences between surface-weighted and depth resolved profiles using *in situ* and HSRL lidar data.

- Ocean-color-style estimates off by as much as 54%
- Errors can be much larger in other parts of the ocean



Modified from Schullien et al., **Vertically- resolved phytoplankton carbon and net primary production from a High Spectral Resolution Lidar**, submitted March 2017.



High vertical resolution also greatly facilitates water cloud extinction from LaRC airborne HSRL-1

- Fast, linear detector provides high vertical resolution (1.25 m) cloud extinction profile
- Cloud top height measured with high accuracy
- Depolarization provides multiple scattering fraction
- HSRL provides accurate measure of optical depth (i.e. attenuation) to cloud top


Example from HSRL-1 data from NAAMES mission



Cloud Drop Number Concentration (N_d) from Combined HSRL and Polarimeter Cloud Measurements




- CDNC derived from CALIPSO-MODIS-CERES (Hu et al., 2007)
 - CALIPSO cloud extinction retrievals
 - CERES-MODIS effective radius
- Advantage of HSRL, Polarimeter retrievals
 - High resolution HSRL (active) - provides more accurate cloud extinction profile
 - RSP (passive) - provides more accurate cloud drop effective radius, variance



$$N_d = \frac{\text{Extinction}}{\text{Scattering Cross Section}}$$


From HSRL extinction retrieval with multiple scattering correction

From RSP size distributions




MESCAL:

Monitoring the Evolving State of Clouds and Aerosol Layers



- Began as a French cloud-aerosol focused lidar mission concept championed by Dr. Helene Chepfer (LMD/IPSL, Université Pierre et Marie Curie)
- Evolved into a joint mission study between CNES and NASA Langley
- Science requirements include
 - Continuation of CALIPSO cloud and aerosol record for radiation budget and cloud-climate feedback studies
 - Aerosol requirements from the ACE Decadal Survey mission (ACE = Aerosols-Cloud-Ecosystems mission)
 - Ocean requirements from ACE
- Langley concept based on CALIPSO and airborne HSRL heritage





Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO)



Objective: global profiling of clouds and aerosols for radiation budget applications

- 11 years of operations in the A-train constellation
- 1800 journal publications

Demonstrated

1. Necessity of vertically-resolved measurements for cloud and aerosol studies
2. That lidar can be reliable in space
3. Spaceborne lidar has the sensitivity to provide valuable ocean measurements




CALIPSO website: www-calipso.larc.nasa.gov


Cloud Drop Size from LaRC airborne HSRL-1

- Cloud lidar ratio (i.e. extinction/backscatter) is function of cloud drop size
- HSRL provides direct measure of cloud lidar ratio
- Initial comparisons of HSRL-1 derived cloud lidar ratio are well correlated with cloud drop effective radius derived from coincident RSP polarized radiance measurements



HSRL-1 and RSP measurements



Theoretical relationship



- Began as a French cloud-aerosol focused lidar mission concept championed by Dr. Helene Chepfer (LMD/IPSL, Université Pierre et Marie Curie)
- Evolved into a joint mission study between CNES and NASA Langley
- Science requirements include
 - Continuation of CALIPSO cloud and aerosol record for radiation budget and cloud-climate feedback studies
 - Aerosol requirements from the ACE Decadal Survey mission (ACE = Aerosols-Cloud-Ecosystems mission)
 - Ocean requirements from ACE
- Langley concept based on CALIPSO and airborne HSRL heritage



MESCAL raises possibilities:

- Reduce AOD bias, even during daytime
- Direct measurements of backscatter and extinction facilitates additional averaging or integration
- Multiwavelength HSRL techniques provide potential for retrieving additional parameters for radiative flux calculations and estimating aerosol forcing

HSRL measurements used to evaluate WRF Chem aerosol simulations



- WRF-Chem aerosol simulations from over California evaluated using HSRL
- HSRL was most valuable instrument to quantify overprediction of aerosols in free troposphere
- Long-range transport of aerosols by global model was too high in free troposphere. This bias led to overpredictions in AOD by factor of two and offsets effects of underprediction of BL aerosols
- Reducing long-range transport greatly improves simulated AOD

