

Test of Lepton Universality in $Y(1S)$ Decays at *BABAR*

P. del Amo Sanchez,¹ J. P. Lees,¹ V. Poireau,¹ E. Prencipe,¹ V. Tisserand,¹ J. Garra Tico,² E. Grauges,² M. Martinelli,^{3,4} A. Palano,^{3,4} M. Pappagallo,^{3,4} G. Eigen,⁵ B. Stugu,⁵ L. Sun,⁵ M. Battaglia,⁶ D. N. Brown,⁶ B. Hooperman,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Lynch,⁶ I. L. Osipenkov,⁶ T. Tanabe,⁶ C. M. Hawkes,⁷ N. Soni,⁷ A. T. Watson,⁷ H. Koch,⁸ T. Schroeder,⁸ D. J. Asgeirsson,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ A. Khan,¹⁰ A. Randle-Conde,¹⁰ V. E. Blinov,¹¹ A. R. Buzykaev,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ K. Yu. Todyshev,¹¹ A. N. Yushkov,¹¹ M. Bondioli,¹² S. Curry,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² E. C. Martin,¹² D. P. Stoker,¹² H. Atmacan,¹³ J. W. Gary,¹³ F. Liu,¹³ O. Long,¹³ G. M. Vitug,¹³ Z. Yasin,¹³ V. Sharma,¹⁴ C. Campagnari,¹⁵ T. M. Hong,¹⁵ D. Kovalskyi,¹⁵ J. D. Richman,¹⁵ A. M. Eisner,¹⁶ C. A. Heusch,¹⁶ J. Kroseberg,¹⁶ W. S. Lockman,¹⁶ A. J. Martinez,¹⁶ T. Schalk,¹⁶ B. A. Schumm,¹⁶ A. Seiden,¹⁶ L. O. Winstrom,¹⁶ C. H. Cheng,¹⁷ D. A. Doll,¹⁷ B. Echenard,¹⁷ D. G. Hitlin,¹⁷ P. Ongmongkolkul,¹⁷ F. C. Porter,¹⁷ A. Y. Rakitin,¹⁷ R. Andreassen,¹⁸ M. S. Dubrovin,¹⁸ G. Mancinelli,¹⁸ B. T. Meadows,¹⁸ M. D. Sokoloff,¹⁸ P. C. Bloom,¹⁹ W. T. Ford,¹⁹ A. Gaz,¹⁹ J. F. Hirschauer,¹⁹ M. Nagel,¹⁹ U. Nauenberg,¹⁹ J. G. Smith,¹⁹ S. R. Wagner,¹⁹ R. Ayad,^{20,*} W. H. Toki,²⁰ A. Hauke,²¹ H. Jasper,²¹ T. M. Karbach,²¹ J. Merkel,²¹ A. Petzold,²¹ B. Spaan,²¹ K. Wacker,²¹ M. J. Kobel,²² K. R. Schubert,²² R. Schwierz,²² D. Bernard,²³ M. Verderi,²³ P. J. Clark,²⁴ S. Playfer,²⁴ J. E. Watson,²⁴ M. Andreotti,^{25,26} D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,^{25,26} A. Cecchi,^{25,26} G. Cibinetto,^{25,26} E. Fioravanti,^{25,26} P. Franchini,^{25,26} E. Luppi,^{25,26} M. Munerato,^{25,26} M. Negrini,^{25,26} A. Petrella,^{25,26} L. Piemontese,²⁵ R. Baldini-Ferroli,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ M. Nicolaci,²⁷ S. Pacetti,²⁷ P. Patteri,²⁷ I. M. Peruzzi,^{27,†} M. Piccolo,²⁷ M. Rama,²⁷ A. Zallo,²⁷ R. Contri,^{28,29} E. Guido,^{28,29} M. Lo Vetere,^{28,29} M. R. Monge,^{28,29} S. Passaggio,²⁸ C. Patrignani,^{28,29} E. Robutti,²⁸ S. Tosi,^{28,29} B. Bhuyan,³⁰ M. Morii,³¹ A. Adametz,³² J. Marks,³² S. Schenk,³² U. Uwer,³² F. U. Bernlochner,³³ H. M. Lacker,³³ T. Lueck,³³ A. Volk,³³ P. D. Dauncey,³⁴ M. Tibbetts,³⁴ P. K. Behera,³⁵ U. Mallik,³⁵ C. Chen,³⁶ J. Cochran,³⁶ H. B. Crawley,³⁶ L. Dong,³⁶ W. T. Meyer,³⁶ S. Prell,³⁶ E. I. Rosenberg,³⁶ A. E. Rubin,³⁶ Y. Y. Gao,³⁷ A. V. Gritsan,³⁷ Z. J. Guo,³⁷ N. Arnaud,³⁸ M. Davier,³⁸ D. Derkach,³⁸ J. Firmino da Costa,³⁸ G. Grosdidier,³⁸ F. Le Diberder,³⁸ A. M. Lutz,³⁸ B. Malaescu,³⁸ A. Perez,³⁸ P. Roudeau,³⁸ M. H. Schune,³⁸ J. Serrano,³⁸ V. Sordini,^{38,‡} A. Stocchi,³⁸ L. Wang,³⁸ G. Wormser,³⁸ D. J. Lange,³⁹ D. M. Wright,³⁹ I. Bingham,⁴⁰ J. P. Burke,⁴⁰ C. A. Chavez,⁴⁰ J. P. Coleman,⁴⁰ J. R. Fry,⁴⁰ E. Gabathuler,⁴⁰ R. Gamet,⁴⁰ D. E. Hutchcroft,⁴⁰ D. J. Payne,⁴⁰ C. Touramanis,⁴⁰ A. J. Bevan,⁴¹ F. Di Lodovico,⁴¹ R. Sacco,⁴¹ M. Sigamani,⁴¹ G. Cowan,⁴² S. Paramesvaran,⁴² A. C. Wren,⁴² D. N. Brown,⁴³ C. L. Davis,⁴³ A. G. Denig,⁴⁴ M. Fritsch,⁴⁴ W. Gradl,⁴⁴ A. Hafner,⁴⁴ K. E. Alwyn,⁴⁵ D. Bailey,⁴⁵ R. J. Barlow,⁴⁵ G. Jackson,⁴⁵ G. D. Lafferty,⁴⁵ T. J. West,⁴⁵ J. Anderson,⁴⁶ R. Cenci,⁴⁶ A. Jawahery,⁴⁶ D. A. Roberts,⁴⁶ G. Simi,⁴⁶ J. M. Tuggle,⁴⁶ C. Dallapiccola,⁴⁷ E. Salvati,⁴⁷ R. Cowan,⁴⁸ D. Dujmic,⁴⁸ P. H. Fisher,⁴⁸ G. Sciolla,⁴⁸ R. K. Yamamoto,⁴⁸ M. Zhao,⁴⁸ P. M. Patel,⁴⁹ S. H. Robertson,⁴⁹ M. Schram,⁴⁹ P. Biassoni,^{50,51} A. Lazzaro,^{50,51} V. Lombardo,⁵⁰ F. Palombo,^{50,51} S. Stracka,^{50,51} L. Cremaldi,⁵² R. Godang,^{52,§} R. Kroeger,⁵² P. Sonnek,⁵² D. J. Summers,⁵² H. W. Zhao,⁵² X. Nguyen,⁵³ M. Simard,⁵³ P. Taras,⁵³ G. De Nardo,^{54,55} D. Monorchio,^{54,55} G. Onorato,^{54,55} C. Sciacca,^{54,55} G. Raven,⁵⁶ H. L. Snoek,⁵⁶ C. P. Jessop,⁵⁷ K. J. Knoepfel,⁵⁷ J. M. LoSecco,⁵⁷ W. F. Wang,⁵⁷ L. A. Corwin,⁵⁸ K. Honscheid,⁵⁸ R. Kass,⁵⁸ J. P. Morris,⁵⁸ A. M. Rahimi,⁵⁸ N. L. Blount,⁵⁹ J. Brau,⁵⁹ R. Frey,⁵⁹ O. Igonkina,⁵⁹ J. A. Kolb,⁵⁹ R. Rahmat,⁵⁹ N. B. Sinev,⁵⁹ D. Strom,⁵⁹ J. Strube,⁵⁹ E. Torrence,⁵⁹ G. Castelli,^{60,61} E. Feltresi,^{60,61} N. Gagliardi,^{60,61} M. Margoni,^{60,61} M. Morandin,⁶⁰ M. Posocco,⁶⁰ M. Rotondo,⁶⁰ F. Simonetto,^{60,61} R. Stroili,^{60,61} E. Ben-Haim,⁶² G. R. Bonneaud,⁶² H. Briand,⁶² J. Chauveau,⁶² O. Hamon,⁶² Ph. Leruste,⁶² G. Marchiori,⁶² J. Ocariz,⁶² J. Prendki,⁶² S. Sitt,⁶² M. Biasini,^{63,64} E. Manoni,^{63,64} C. Angelini,^{65,66} G. Batignani,^{65,66} S. Bettarini,^{65,66} G. Calderini,^{65,66,||} M. Carpinelli,^{65,66,||} A. Cervelli,^{65,66} F. Forti,^{65,66} M. A. Giorgi,^{65,66} A. Lusiani,^{65,67} N. Neri,^{65,66} E. Paoloni,^{65,66} G. Rizzo,^{65,66} J. J. Walsh,⁶⁵ D. Lopes Pegna,⁶⁸ C. Lu,⁶⁸ J. Olsen,⁶⁸ A. J. S. Smith,⁶⁸ A. V. Telnov,⁶⁸ F. Anulli,⁶⁹ E. Baracchini,^{69,70} G. Cavoto,⁶⁹ R. Faccini,^{69,70} F. Ferrarotto,⁶⁹ F. Ferroni,^{69,70} M. Gaspero,^{69,70} L. Li Gioi,⁶⁹ M. A. Mazzoni,⁶⁹ G. Piredda,⁶⁹ F. Renga,^{69,70} M. Ebert,⁷¹ T. Hartmann,⁷¹ T. Leddig,⁷¹ H. Schröder,⁷¹ R. Waldi,⁷¹ T. Adye,⁷² B. Franek,⁷² E. O. Olaiya,⁷² F. F. Wilson,⁷² S. Emery,⁷³ G. Hamel de Monchenault,⁷³ G. Vasseur,⁷³ Ch. Yèche,⁷³ M. Zito,⁷³ M. T. Allen,⁷⁴ D. Aston,⁷⁴ D. J. Bard,⁷⁴ R. Bartoldus,⁷⁴ J. F. Benitez,⁷⁴ C. Cartaro,⁷⁴ M. R. Convery,⁷⁴ J. Dorfan,⁷⁴ G. P. Dubois-Felsmann,⁷⁴ W. Dunwoodie,⁷⁴ R. C. Field,⁷⁴ M. Franco Sevilla,⁷⁴ B. G. Fulsom,⁷⁴ A. M. Gabareen,⁷⁴ M. T. Graham,⁷⁴ P. Grenier,⁷⁴ C. Hast,⁷⁴ W. R. Innes,⁷⁴ M. H. Kelsey,⁷⁴ H. Kim,⁷⁴ P. Kim,⁷⁴ M. L. Kocian,⁷⁴ D. W. G. S. Leith,⁷⁴ S. Li,⁷⁴ B. Lindquist,⁷⁴ S. Luitz,⁷⁴ V. Luth,⁷⁴ H. L. Lynch,⁷⁴ D. B. MacFarlane,⁷⁴ H. Marsiske,⁷⁴ D. R. Muller,⁷⁴ H. Neal,⁷⁴ S. Nelson,⁷⁴ C. P. O'Grady,⁷⁴ I. Ofte,⁷⁴ M. Perl,⁷⁴ B. N. Ratcliff,⁷⁴ A. Roodman,⁷⁴ A. A. Salnikov,⁷⁴ V. Santoro,⁷⁴ R. H. Schindler,⁷⁴ J. Schwiening,⁷⁴

A. Snyder,⁷⁴ D. Su,⁷⁴ M. K. Sullivan,⁷⁴ K. Suzuki,⁷⁴ J. M. Thompson,⁷⁴ J. Va'vra,⁷⁴ A. P. Wagner,⁷⁴ M. Weaver,⁷⁴ C. A. West,⁷⁴ W. J. Wisniewski,⁷⁴ M. Wittgen,⁷⁴ D. H. Wright,⁷⁴ H. W. Wulsin,⁷⁴ A. K. Yarritu,⁷⁴ C. C. Young,⁷⁴ V. Ziegler,⁷⁴ X. R. Chen,⁷⁵ W. Park,⁷⁵ M. V. Purohit,⁷⁵ R. M. White,⁷⁵ J. R. Wilson,⁷⁵ S. J. Sekula,⁷⁶ M. Bellis,⁷⁷ P. R. Burchat,⁷⁷ A. J. Edwards,⁷⁷ T. S. Miyashita,⁷⁷ S. Ahmed,⁷⁸ M. S. Alam,⁷⁸ J. A. Ernst,⁷⁸ B. Pan,⁷⁸ M. A. Saeed,⁷⁸ S. B. Zain,⁷⁸ N. Guttman,⁷⁹ A. Soffer,⁷⁹ P. Lund,⁸⁰ S. M. Spanier,⁸⁰ R. Eckmann,⁸¹ J. L. Ritchie,⁸¹ A. M. Ruland,⁸¹ C. J. Schilling,⁸¹ R. F. Schwitters,⁸¹ B. C. Wray,⁸¹ J. M. Izen,⁸² X. C. Lou,⁸² F. Bianchi,^{83,84} D. Gamba,^{83,84} M. Pelliccioni,^{83,84} M. Bomben,^{85,86} G. Della Ricca,^{85,86} L. Lanceri,^{85,86} L. Vitale,^{85,86} V. Azzolini,⁸⁷ N. Lopez-March,⁸⁷ F. Martinez-Vidal,⁸⁷ D. A. Milanes,⁸⁷ A. Oyanguren,⁸⁷ J. Albert,⁸⁸ Sw. Banerjee,⁸⁸ H. H. F. Choi,⁸⁸ K. Hamano,⁸⁸ G. J. King,⁸⁸ R. Kowalewski,⁸⁸ M. J. Lewczuk,⁸⁸ I. M. Nugent,⁸⁸ J. M. Roney,⁸⁸ R. J. Sobie,⁸⁸ T. J. Gershon,⁸⁹ P. F. Harrison,⁸⁹ J. Ilic,⁸⁹ T. E. Latham,⁸⁹ G. B. Mohanty,⁸⁹ E. M. T. Puccio,⁸⁹ H. R. Band,⁹⁰ X. Chen,⁹⁰ S. Dasu,⁹⁰ K. T. Flood,⁹⁰ Y. Pan,⁹⁰ M. Pierini,⁹⁰ R. Prepost,⁹⁰ C. O. Vuosalo,⁹⁰ and S. L. Wu⁹⁰

(BABAR Collaboration)

¹Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP), Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France

²Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

³INFN Sezione di Bari, I-70126 Bari, Italy

⁴Dipartimento di Fisica, Università di Bari, I-70126 Bari, Italy

⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Riverside, Riverside, California 92521, USA

¹⁴University of California at San Diego, La Jolla, California 92093, USA

¹⁵University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁶University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁷California Institute of Technology, Pasadena, California 91125, USA

¹⁸University of Cincinnati, Cincinnati, Ohio 45221, USA

¹⁹University of Colorado, Boulder, Colorado 80309, USA

²⁰Colorado State University, Fort Collins, Colorado 80523, USA

²¹Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany

²²Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²³Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

²⁴University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²⁵INFN Sezione di Ferrara, I-44100 Ferrara, Italy

²⁶Dipartimento di Fisica, Università di Ferrara, I-44100 Ferrara, Italy

²⁷INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy

²⁸INFN Sezione di Genova, I-16146 Genova, Italy

²⁹Dipartimento di Fisica, Università di Genova, I-16146 Genova, Italy

³⁰Indian Institute of Technology Guwahati, Guwahati, Assam, 781 039, India

³¹Harvard University, Cambridge, Massachusetts 02138, USA

³²Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

³³Humboldt-Universität zu Berlin, Institut für Physik, Newtonstr. 15, D-12489 Berlin, Germany

³⁴Imperial College London, London, SW7 2AZ, United Kingdom

³⁵University of Iowa, Iowa City, Iowa 52242, USA

³⁶Iowa State University, Ames, Iowa 50011-3160, USA

³⁷Johns Hopkins University, Baltimore, Maryland 21218, USA

³⁸Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 Orsay Cedex, France

³⁹Lawrence Livermore National Laboratory, Livermore, California 94550, USA

⁴⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom

⁴¹Queen Mary, University of London, London, E1 4NS, United Kingdom

⁴²University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

- ⁴³University of Louisville, Louisville, Kentucky 40292, USA
- ⁴⁴Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany
- ⁴⁵University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴⁶University of Maryland, College Park, Maryland 20742, USA
- ⁴⁷University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴⁸Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- ⁴⁹McGill University, Montréal, Québec, Canada H3A 2T8
- ⁵⁰INFN Sezione di Milano, I-20133 Milano, Italy
- ⁵¹Dipartimento di Fisica, Università di Milano, I-20133 Milano, Italy
- ⁵²University of Mississippi, University, Mississippi 38677, USA
- ⁵³Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
- ⁵⁴INFN Sezione di Napoli, I-80126 Napoli, Italy
- ⁵⁵Dipartimento di Scienze Fisiche, Università di Napoli Federico II, I-80126 Napoli, Italy
- ⁵⁶NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵⁷University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵⁸Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁹University of Oregon, Eugene, Oregon 97403, USA
- ⁶⁰INFN Sezione di Padova, I-35131 Padova, Italy
- ⁶¹Dipartimento di Fisica, Università di Padova, I-35131 Padova, Italy
- ⁶²Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France
- ⁶³INFN Sezione di Perugia, I-06100 Perugia, Italy
- ⁶⁴Dipartimento di Fisica, Università di Perugia, I-06100 Perugia, Italy
- ⁶⁵INFN Sezione di Pisa, I-56127 Pisa, Italy
- ⁶⁶Dipartimento di Fisica, Università di Pisa, I-56127 Pisa, Italy
- ⁶⁷Scuola Normale Superiore di Pisa, I-56127 Pisa, Italy
- ⁶⁸Princeton University, Princeton, New Jersey 08544, USA
- ⁶⁹INFN Sezione di Roma, I-00185 Roma, Italy
- ⁷⁰Dipartimento di Fisica, Università di Roma La Sapienza, I-00185 Roma, Italy
- ⁷¹Universität Rostock, D-18051 Rostock, Germany
- ⁷²Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁷³CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France
- ⁷⁴SLAC National Accelerator Laboratory, Stanford, California 94309 USA
- ⁷⁵University of South Carolina, Columbia, South Carolina 29208, USA
- ⁷⁶Southern Methodist University, Dallas, Texas 75275, USA
- ⁷⁷Stanford University, Stanford, California 94305-4060, USA
- ⁷⁸State University of New York, Albany, New York 12222, USA
- ⁷⁹Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel
- ⁸⁰University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁸¹University of Texas at Austin, Austin, Texas 78712, USA
- ⁸²University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁸³INFN Sezione di Torino, I-10125 Torino, Italy
- ⁸⁴Dipartimento di Fisica Sperimentale, Università di Torino, I-10125 Torino, Italy
- ⁸⁵INFN Sezione di Trieste, I-34127 Trieste, Italy
- ⁸⁶Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy
- ⁸⁷IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
- ⁸⁸University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁸⁹Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
- ⁹⁰University of Wisconsin, Madison, Wisconsin 53706, USA

(Received 24 February 2010; published 14 May 2010)

The ratio $R_{\tau\mu}(Y(1S)) = \Gamma_{Y(1S) \rightarrow \tau^+ \tau^-} / \Gamma_{Y(1S) \rightarrow \mu^+ \mu^-}$ is measured using a sample of $(121.8 \pm 1.2) \times 10^6 Y(3S)$ events recorded by the BABAR detector. This measurement is intended as a test of lepton universality and as a search for a possible light pseudoscalar Higgs boson. In the standard model (SM) this ratio is expected to be close to 1. Any significant deviations would violate lepton universality and could be introduced by the coupling to a light pseudoscalar Higgs boson. The analysis studies the decays $Y(3S) \rightarrow Y(1S)\pi^+\pi^-$, $Y(1S) \rightarrow l^+l^-$, where $l = \mu, \tau$. The result, $R_{\tau\mu}(Y(1S)) = 1.005 \pm 0.013(\text{stat}) \pm 0.022(\text{syst})$, shows no deviation from the expected SM value, while improving the precision with respect to previous measurements.

In the standard model (SM), the couplings of the gauge bosons to leptons are independent of the lepton flavor. Aside from small lepton-mass effects, the expression for the decay width $\Upsilon(1S) \rightarrow l^+ l^-$ should be identical for all leptons, and given by [1]:

$$\Gamma_{\Upsilon(1S) \rightarrow ll} = 4\alpha^2 Q_b^2 \frac{|R_n(0)|^2}{M_Y^2} \left(1 + 2 \frac{M_l^2}{M_Y^2}\right) \sqrt{1 - 4 \frac{M_l^2}{M_Y^2}}, \quad (1)$$

where α is the electromagnetic fine structure constant, Q_b is the charge of the bottom quark, $R_n(0)$ is the nonrelativistic radial wave function of the bound $b\bar{b}$ state evaluated at the origin, M_Y is the $\Upsilon(1S)$ mass and M_l is the lepton mass. In the SM, one expects the quantity

$$R_{ll'}(\Upsilon(1S)) = \frac{\Gamma_{\Upsilon(1S) \rightarrow ll}}{\Gamma_{\Upsilon(1S) \rightarrow l'l'}} \quad (2)$$

with $l, l' = e, \mu, \tau$ and $l' \neq l$, to be close to one. In particular, the value for $R_{\tau\mu}(\Upsilon(1S))$ is predicted to be ~ 0.992 [2].

In the next-to-minimal extension of the SM [3], deviations of $R_{ll'}$ from the SM expectation may arise due to a light CP -odd Higgs boson, A^0 . Present data [4] do not exclude the existence of such a boson with a mass below 10 GeV/ c^2 . Among other hypothetical particles, A^0 may mediate the following processes [1]:

$$\Upsilon(1S) \rightarrow A^0 \gamma \rightarrow l^+ l^- \gamma \quad (3)$$

or

$$\Upsilon(1S) \rightarrow \eta_b(1S) \gamma, \quad \eta_b(1S) \rightarrow A^0 \rightarrow l^+ l^-. \quad (4)$$

The latter implies a mixing between A^0 and $\eta_b(1S)$, which is a 1S_0 $b\bar{b}$ state and therefore not expected to decay to a lepton pair to leading order in the SM.

If the photon is energetic enough to be detectable, a monochromatic peak in the photon spectrum recoiling against the lepton pair could be an indication of new physics (NP) [5,6]. Alternatively, if the photon remains undetected, the lepton pair would be ascribed to the $\Upsilon(1S)$ and the proportionality of the coupling of the Higgs to the lepton mass would lead to an apparent violation of lepton universality. This effect should be larger for decays to $\tau^+ \tau^-$ pairs, and enhanced for higher-mass $\Upsilon(nS)$ and $\eta_b(nS)$ resonances. The deviation of $R_{ll'}$ from the expected SM value depends on $X_d = \cos\theta_A \tan\beta$ (where θ_A measures the coupling of the $\Upsilon(1S)$ to the A^0 , and $\tan\beta$ is the ratio of the vacuum expectation values of the two Higgs doublets) and on the mass difference between A^0 and $\eta_b(1S)$. Assuming $X_d = 12$ (a representative value evading present limits [4]), $\Gamma(\eta_b(1S)) = 5$ MeV, and $M_{\eta_b(1S)}$ as measured in [7], the deviation of $R_{\tau\mu}(\Upsilon(1S))$ may be as large as $\sim 4\%$, depending on the A^0 mass [1]. A measurement of this ratio has already been performed, with the result $R_{\tau\mu}(\Upsilon(1S)) = 1.02 \pm 0.02(\text{stat}) \pm 0.05(\text{syst})$ [8].

This Letter focuses on the measurement of $R_{\tau\mu}(\Upsilon(1S))$, in the decays $\Upsilon(3S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ with $\Upsilon(1S) \rightarrow l^+ l^-$ and $l = \mu, \tau$. In this analysis only τ decays to a single charged particle (plus neutrinos) are considered. This choice simplifies the analysis; in particular, it results in final states of exactly four detected particles for both the $\mu^+ \mu^-$ and $\tau^+ \tau^-$ samples. The data collected at the $\Upsilon(3S)$ resonance by the *BABAR* detector at the PEP-II storage rings correspond to 28 fb $^{-1}$. About one tenth of the complete available statistics is used to validate the analysis method and the signal extraction procedure. This validation sample is discarded from the final result in order to avoid any possible bias. A sample of 2.4 fb $^{-1}$ collected about 30 MeV below the $\Upsilon(3S)$ resonance (off-resonance sample) is also used as a background control sample.

The *BABAR* detector is described in detail elsewhere [9,10].

The event selection is optimized using Monte Carlo (MC) simulated events, generated with EVTGEN [11]. GEANT [12] is used to reproduce interactions of particles traversing the *BABAR* detector, taking into account the varying detector conditions and beam backgrounds. Final state radiation effects are simulated using PHOTOS [13].

The selection requires exactly four charged tracks, each with transverse momentum $0.1 < p_T < 10$ GeV/ c , geometrically constrained to come from the same point. The distance of closest approach to the interaction region of each track must be less than 10 cm when projected along the beam axis and less than 1.5 cm in the transverse plane. The ratio of the 2nd to 0th Fox-Wolfram moments (R2) [14] is required to be less than 0.97, and the absolute value of the cosine of the polar angle of the thrust axis [15] to be less than 0.96.

A $\Upsilon(1S) \rightarrow l^+ l^-$ candidate is formed by selecting two oppositely-charged tracks, constrained to come from a common vertex, and it is combined with two other oppositely-charged tracks, assigned the pion mass, to construct a $\Upsilon(3S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ candidate.

Different selection criteria are used for the $\Upsilon(1S) \rightarrow \mu^+ \mu^-$ and the $\Upsilon(1S) \rightarrow \tau^+ \tau^-$ decays, because in the latter the presence of neutrinos in the final state leads to a larger contamination from the background (mainly non-leptonic $\Upsilon(1S)$ decays and $e^+ e^- \rightarrow \tau^+ \tau^-$ events). The $\Upsilon(1S) \rightarrow \mu^+ \mu^-$ candidates (D_μ) are selected by requiring two tracks in the final state identified as muons. This identification is performed by using information from different subdetectors, such as the energy deposited in the electromagnetic calorimeter, the number of hits in the instrumented flux return of the magnet and the number of interaction lengths traversed, combined in a neural-network algorithm. Calculated in the $e^+ e^-$ center-of-mass (c.m.) frame [16], the difference between the initial state energy and the visible final state energy is required to be less than 0.5 GeV, the magnitude of the dipion momentum ($p_{\pi\pi}^*$) less than 0.875 GeV/ c , and the cosine of the

angle between the two lepton candidates less than -0.96 . For the $Y(1S) \rightarrow \tau^+ \tau^-$ candidates (D_τ), tighter selection criteria are applied to reduce background. In these events a large fraction of the energy is not reconstructed, due to the presence of neutrinos; thus the difference between the energy of the initial state and the energy detected in the final state, calculated in the $e^+ e^-$ c.m. frame, is required to exceed 5 GeV. Further requirements are made on the magnitude of the dipion momentum ($p_{\pi\pi}^* < 0.825$ GeV/ c) and on the magnitude of the momentum of each π ($p_\pi^* < 0.725$ GeV/ c). The measured difference in the energy of the $Y(3S)$ and the $Y(1S)$ is restricted to $0.835 < \Delta E^* < 0.925$ GeV. A boosted decision tree [17] is used to further reduce the background, based on several event shape and kinematic variables such as R2 and the energy of the charged tracks reconstructed in the events. The performance of the classifier is assessed using MC simulations and off-resonance data.

Finally, in order to select $Y(3S) \rightarrow Y(1S)\pi^+\pi^-$ candidates, the invariant mass difference $\Delta M = M(Y(3S)) - M(Y(1S))$, calculated with the reconstructed tracks of the final state, is required to be less than 2.5 GeV/ c^2 and the dipion invariant mass ($M_{\pi\pi}$) to be between 0.28 and 0.90 GeV/ c^2 .

For events with multiple candidates, the candidate with the value of ΔM closest to the nominal value [2] is retained as the best one. It has been verified by MC simulations that the selection requirements do not reduce the sensitivity to NP processes. Since the possible NP effects, with the presence of additional photons in the process, should be more evident in $Y(1S) \rightarrow \tau^+ \tau^-$ events, variables that are sensitive to neutral energy are not used in the selection.

The final selection efficiency for the reconstructed decay chains, estimated from a sample of MC-simulated events, are $\epsilon_{\mu\mu} = (44.57 \pm 0.04)\%$ and $\epsilon_{\tau\tau} = (16.77 \pm 0.03)\%$ for the $\mu^+ \mu^-$ and the $\tau^+ \tau^-$ final states, respectively.

An extended unbinned maximum likelihood fit, applied simultaneously to the two disjoint data sets D_μ and D_τ , is used to extract $R_{\tau\mu} = \frac{N_{\text{sig}\tau}}{\epsilon_{\tau\tau}} \cdot \frac{\epsilon_{\mu\mu}}{N_{\text{sig}\mu}}$, where $N_{\text{sig}\mu}$ ($N_{\text{sig}\tau}$) indicates the number of signal events in the D_μ (D_τ) sample. For the D_μ sample, a two-dimensional probability density function (PDF) is used, based on the invariant dimuon mass $M_{\mu^+ \mu^-}$ and $M_{\pi^+ \pi^-}^{\text{reco}}$, the invariant mass of the system recoiling against the pion pair, defined as:

$$M_{\pi^+ \pi^-}^{\text{reco}} = \sqrt{s + M_{\pi\pi}^2 - 2 \cdot \sqrt{s} \cdot E_{\pi\pi}^*}, \quad (5)$$

where \sqrt{s} is the $e^+ e^-$ center-of-mass energy and $E_{\pi\pi}^*$ indicates the $\pi^+ \pi^-$ pair energy. MC simulations are used to check that the two variables are uncorrelated. For the D_τ sample, a one-dimensional PDF is used, based on $M_{\pi^+ \pi^-}^{\text{reco}}$ [Eq. (5)]. The likelihood is written as:

$$\mathcal{L}_{\text{ext}} = \mathcal{L}_{\text{ext}}^\mu \mathcal{L}_{\text{ext}}^\tau, \quad (6)$$

where:

$$\mathcal{L}_{\text{ext}}^i = \frac{e^{-N'_i} (N'_i)^{N_i}}{N'_i!} \prod_{k=1}^{N_i} \mathcal{P}_k^i, \quad (7)$$

with $i = \mu$ or τ and where N_i and N'_i are the sum of the signal and background events, observed and expected, respectively, in each sample. \mathcal{P}_k is the probability to measure a set of physical observables in the k th event, defined as:

$$\begin{aligned} \mathcal{P}_k^\mu &\equiv \frac{N_{\text{sig}\mu}}{N'_\mu} \mathcal{P}_k^\mu(M_{\pi^+ \pi^-}^{\text{reco}}) \mathcal{P}_k^\mu(M_{\mu^+ \mu^-}) \\ &+ \frac{N_{\text{bkg}\mu}}{N'_\mu} \mathcal{P}_k^{\text{bkg}\mu}(M_{\pi^+ \pi^-}^{\text{reco}}) \mathcal{P}_k^{\text{bkg}\mu}(M_{\mu^+ \mu^-}) \end{aligned} \quad (8)$$

and

$$\mathcal{P}_k^\tau \equiv \frac{\epsilon_{\tau\tau}}{\epsilon_{\mu\mu}} \frac{N_{\text{sig}\mu}}{N'_\tau} R_{\tau\mu} \mathcal{P}_k^\tau(M_{\pi^+ \pi^-}^{\text{reco}}) + \frac{N_{\text{bkg}\tau}}{N'_\tau} \mathcal{P}_k^{\text{bkg}\tau}(M_{\pi^+ \pi^-}^{\text{reco}}) \quad (9)$$

where $N_{\text{bkg}\mu}$ ($N_{\text{bkg}\tau}$) indicates the number of background events in the D_μ (D_τ) sample.

The functional forms of the PDFs describing the signal components are modeled from the dedicated subsample consisting of one tenth of the D_μ sample. Both the $M_{\pi^+ \pi^-}^{\text{reco}}$ and the $M_{\mu^+ \mu^-}$ distributions are described by an analytical function approximating a Gaussian distribution function with mean value μ but different left and right widths, $\sigma_{L,R}$, plus asymmetric non-Gaussian tails $\alpha_{L,R}$, defined as:

$$\mathcal{F}(x) = \exp\left\{-\frac{(x - \mu)^2}{2\sigma_{L,R}^2 + \alpha_{L,R}(x - \mu)^2}\right\}. \quad (10)$$

All the parameters (the five parameters describing the $M_{\mu^+ \mu^-}$ distribution, along with the mean values and the widths of both the $M_{\pi^+ \pi^-}^{\text{reco}}$ distributions) are free in the fit, except for $\alpha_{L,R}$ in $M_{\pi^+ \pi^-}^{\text{reco}}$. The off-resonance sample is used to model the background shapes. Constants are chosen for the D_μ sample, and a first order polynomial for the D_τ sample, with all the parameters free in the fit.

The result of the simultaneous fit is $R_{\tau\mu} = 1.006 \pm 0.013$, where the quoted error is statistical only. Figure 1 shows the projections of the fit results for the three variables.

Several systematic errors cancel in the ratio, such as errors on the luminosity, the $Y(3S)$ production cross section, and the $Y(3S) \rightarrow Y(1S)\pi^+\pi^-$ branching fractions, as well as systematic discrepancies between data and simulation in the common event selection and in track reconstruction efficiencies, where a possible dependence on the track energy has been taken into account. The residual systematic uncertainties are related to the differences between data and simulation in the efficiency of event selection, the muon identification, and the trigger and




FIG. 1 (color online). 1D fit projections for $M_{\mu^+\mu^-}$ (top) and for $M_{\pi^+\pi^-}^{\text{reco}}$ (middle) in the D_μ sample, and for $M_{\pi^+\pi^-}^{\text{reco}}$ (bottom) in the D_τ sample. In each plot the dashed line represents the background shape, while the solid line is the sum of signal and background contributions to the fit, and the points are the data.

background digital filters (BGFs) [18]. There is also a systematic uncertainty on the signal and background yields due to the imperfect knowledge of the PDFs used in the fit.

The systematic uncertainty due to the event selection is evaluated by comparing the shape of each variable between data and simulation and estimating the difference in the efficiency. The resulting systematic uncertainty is 1.2%.

The systematic uncertainty related to the difference between data and simulation of muon identification efficiencies, applied only to $Y(1S) \rightarrow \mu^+\mu^-$ events, is estimated by using two independent samples: one where both leptons are required to be identified as muons, and another where exactly one final charged track is a muon. The ratio of the efficiencies for requiring each sample is determined, both on data and on simulation; the ratio of the two results gives an efficiency correction of 1.023 and a related systematic uncertainty of 1.2%.

The systematic uncertainties due to the differences between data and simulation in trigger and BGFs' efficiency are small both in $Y(1S) \rightarrow \mu^+\mu^-$ and in $Y(1S) \rightarrow \tau^+\tau^-$ events, and they cancel partially in the ratio. A correction of 1.020 is needed for the $\epsilon_{\tau\tau}$ efficiency, together with a systematic uncertainty of 0.10% for $Y(1S) \rightarrow \tau^+\tau^-$ events, while a systematic uncertainty of 0.18% is quoted for $Y(1S) \rightarrow \mu^+\mu^-$ events. The impact of the uncertainty in the BGFs' efficiency has been found to be negligible.

The uncertainty due to the imperfect knowledge of the signal and background shapes used in the fit is also estimated. The systematic effect from fixing $\alpha_{L,R}$ in the signal $M_{\pi^+\pi^-}^{\text{reco}}$ PDF is estimated by varying the fixed parameter values by $\pm 1\sigma$ and repeating the fit procedure. Since the correlation between the parameters is found to be negligible, the parameters are varied independently and the deviations from the nominal fit are summed in quadrature, resulting in a total effect of 1.1%. The uncertainty due to the choice of the background PDF shapes is evaluated to be 0.22%, by using alternative parameterizations. In the fit, the same $M_{\pi^+\pi^-}^{\text{reco}}$ functional form is used for both the D_μ and the D_τ sample, ignoring the potential difference in the trigger efficiency. The systematic uncertainty associated with this approximation is evaluated to be 0.6%, by reweighting the parameters for the $M_{\pi^+\pi^-}^{\text{reco}}$ distribution with the parameters obtained from the $\tau^+\tau^-$ data sample, and requiring the magnitude of the momentum of one of the final state charged tracks not to exceed 1 GeV/c.

The $M_{\pi^+\pi^-}^{\text{reco}}$ variable is related only to the $Y(3S) \rightarrow Y(1S)\pi^+\pi^-$ transition and therefore cannot distinguish between $Y(1S) \rightarrow l^+l^-$ events and other $Y(1S)$ decays or the Higgs-mediated events of Eqs. (3) and (4). While this ensures sensitivity to possible NP effects, $Y(1S)$ generic decays could be a relevant source of background in the D_τ sample because the final state is only partially reconstructed. The event selection heavily reduces the yield of the $Y(1S)$ generic decays. It is estimated using a simulated sample of inclusive $Y(1S)$ decays, and is found to be $\sim 0.4\%$ of the $Y(1S) \rightarrow \tau^+\tau^-$ signal yield. Since the hadronic $Y(1S)$ decays are not well measured, the simulation may not be reliable and a systematic uncertainty needs to be considered. A correction factor of 0.996, taking into account this contribution, is applied to the $Y(1S) \rightarrow \tau^+\tau^-$ signal yield, and a systematic uncertainty equal to 0.4% is included as well.

The systematic uncertainty associated with the simulation of the final state radiation by PHOTOS is found to be negligible.

Finally, the finite size of the MC-simulated samples used to determine the efficiencies gives a contribution to the systematic uncertainty less than 0.1% in both the leptonic final states.

The total systematic uncertainty, obtained by summing in quadrature all the contributions, is estimated to be 2.2%.

Including all the systematic corrections, the ratio $R_{\tau\mu}$ is found to be:

$$R_{\tau\mu}(Y(1S)) = 1.005 \pm 0.013(\text{stat}) \pm 0.022(\text{syst}).$$

No significant deviation of the ratio $R_{\tau\mu}$ from the SM expectation is observed. This result improves both the statistical and systematic precision with respect to the previous measurement [8]. According to [1], and assuming values for X_d , $\Gamma(\eta_b(1S))$ and $M_{\eta_b(1S)}$ as previously stated, the present measurement excludes an A^0 with mass lower than 9 GeV/ c^2 at 90% of confidence level.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MES (Russia), MEC (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

*Present address: Temple University, Philadelphia, PA 19122, USA.

†Also at Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

‡Also at Università di Roma La Sapienza, I-00185 Roma, Italy.

§Present address: University of South Alabama, Mobile, AL 36688, USA.

||Also at Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France, USA

¶Also at Università di Sassari, Sassari, Italy.

- [1] M. A. Sanchis-Lozano, *Int. J. Mod. Phys. A* **19**, 2183 (2004); E. Fullana and M. A. Sanchis-Lozano, *Phys. Lett. B* **653**, 67 (2007); F. Domingo *et al.*, *J. High Energy Phys.* 01 (2009) 061.
- [2] C. Amsler *et al.* (Particle Data Group), *Phys. Lett. B* **667**, 1 (2008).
- [3] R. Dermisek and J. F. Gunion, *Phys. Rev. Lett.* **95**, 041801 (2005).
- [4] S. Kraml *et al.*, CERN 2006-009; S. Schael *et al.* (ALEPH, DELPHI, L3 and OPAL Collaborations), *Eur. Phys. J. C* **47**, 547 (2006).
- [5] B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. Lett.* **103**, 081803 (2009).
- [6] B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. Lett.* **103**, 181801 (2009).
- [7] B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. Lett.* **101**, 071801 (2008).
- [8] D. Besson *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **98**, 052002 (2007).
- [9] B. Aubert *et al.* (*BABAR* Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
- [10] W. Menges, in *2006 IEEE Nuclear Science Symposium Conference Record* (IEEE, New York, 2006).
- [11] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [12] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [13] E. Barberio and Z. Was, *Comput. Phys. Commun.* **79**, 291 (1994).
- [14] G. C. Fox and S. Wolfram, *Nucl. Phys. B* **149**, 413 (1979); **157**, 543 (1979).
- [15] S. Brandt *et al.*, *Phys. Lett.* **12**, 57 (1964).
- [16] An asterisk (*) is used to denote variables calculated in the e^+e^- c.m. frame.
- [17] H. J. Yang, B. P. Roe, and J. Zhu, *Nucl. Instrum. Methods Phys. Res., Sect. A* **574**, 342 (2007).
- [18] The background filters are applied to the data in order to reduce the contribution of specific backgrounds. They consist of several requirements on the number, energy, and momentum of the charged tracks and neutral clusters in the event.