

# Anemia Is Associated With Depression in Older Adults: Results From the InCHIANTI Study

Graziano Onder,<sup>1,2</sup> Brenda W. J. H. Penninx,<sup>2</sup> Matteo Cesari,<sup>1,2</sup> Stefania Bandinelli,<sup>3</sup> Fulvio Lauretani,<sup>3</sup>  
Benedetta Bartali,<sup>3</sup> Anna Maria Gori,<sup>4</sup> Marco Pahor,<sup>2</sup> and Luigi Ferrucci<sup>3,5</sup>

<sup>1</sup>Geriatric Department, Catholic University of the Sacred Heart, Rome, Italy.

<sup>2</sup>Sticht Center on Aging, Section on Gerontology and Geriatric Medicine, Wake Forest University School of Medicine, Winston-Salem, North Carolina.

<sup>3</sup>Geriatric Department, "I Fraticini," National Research Institute, Florence, Italy.

<sup>4</sup>Department of Clinical and Surgical Care, Section of Clinical Medicine, Thrombosis Centre, University of Florence, Italy.

<sup>5</sup>Longitudinal Studies Section, Clinical Research Branch, National Institute on Aging, Baltimore, Maryland.

**Background.** Depression is a common disorder among older adults, and it has been associated with adverse outcomes, including increased risk of morbidity and mortality as well as incomplete or delayed recovery from illness and disability. The objective of this study was to examine whether depressive symptoms and anemia are associated among older adults living in the community.

**Methods.** We used data from the "Invecchiare in Chianti" (Aging in the Chianti area, InCHIANTI) study, a prospective population-based study of older people living in the community. Anemia was defined by the World Health Organization (WHO) criteria: hemoglobin concentration below 12 g/dl in women and below 13 g/dl in men. Depressive symptoms were measured by using the Center for Epidemiological Studies Depression Scale (CES-D). Participants with a CES-D score  $\geq 16$  were considered to be depressed.

**Results.** Mean age of the 986 participants was 75 years, and 56% were female; 313 (32%) study participants were depressed. Anemia was recorded in 48 of the 313 (15%) participants with depression and in 53 of the 673 (8%) participants without depression ( $p < .001$ ). After adjusting for potential confounders, depression was associated with a significant higher risk of anemia (odds ratio = 1.93; 95% confidence interval, 1.19–3.13). The risk of anemia progressively and significantly increased with increasing CES-D score (signifying more severe depression). Compared with nondepressed participants (CES-D score  $< 16$ ), the odds ratio for anemia were 1.74, 2.04, and 2.10 for participants with mild (score = 16–20), moderate (score = 21–26), and severe depression (score  $> 26$ ), respectively ( $p$  for linear trend = .01).

**Conclusions.** Depressive symptoms are associated with anemia in a general population of older persons living in the community.

DEPRESSION in the older population is a major health issue, both because of its high prevalence and because of its adverse health consequences. About 12%–20% of community-dwelling older person suffer from symptoms of depression (1,2). Many cross-sectional studies have demonstrated that, compared with nondepressed persons, depressed older persons have poorer health status, higher prevalence of disability, and more severe comorbidity (3,4). In addition, longitudinal studies have shown that depression represents a risk factor for adverse outcomes including increased morbidity, mortality, and incomplete or delayed recovery from illness and disability (5–8).

Anemia is rather common in old age with a prevalence of about 13% in persons aged 70 years or older (9). This condition is assumed to be due to underlying disease such as cancer, renal failure, and infectious disease, and it has been associated with a number of other comorbidities, such as cardiovascular disease (10). This high burden of comorbidity may negatively affect quality of life and lead to the onset of depressive symptoms (11). Moreover, in studies conducted in hemodialysis patients (12,13), it has been shown that anemia directly affects brain function, and it is responsible for the cognitive impairments and symptoms frequently suffered by persons with chronic renal failure (confusion, inability to

concentrate, decreased mental alertness, and impaired memory). Reduced muscle strength and fatigue are commonly associated with anemia and may have detrimental effects on participants' quality of life, therefore facilitating the development of depressive symptoms (14,15). Finally, malnutrition, a common feature in depressed individuals, may be responsible for anemia (10,16). Despite numerous hypothetical pathways suggesting a link between anemia and depression, the association between these two common conditions has rarely been explored in a general population of older adults (17). Therefore, the aim of the present study is to examine whether depressive symptoms are associated with anemia, independent of comorbidity and other potential confounders.

## METHODS

We used data from the "Invecchiare in Chianti" (Aging in the Chianti area, InCHIANTI) study, a population-based study of older people conducted by the Italian National Research Council of Aging (INRCA, Florence, Italy). The InCHIANTI study includes 1156 participants aged 65 years and older who were randomly selected from the residents in two towns of the Chianti geographic area (Greve In Chianti and Bagno a Ripoli, Tuscany, Italy) (18).

### Anemia

Details on blood sampling and determination of biological parameters are reported elsewhere (14,15,18). Anemia was defined by the World Health Organization (WHO) criteria (19): Hemoglobin concentration below 12 g/dl in women and below 13 g/dl in men. A ratio between serum transferrin receptor and log(ferritin) above 1.5 was used to define iron deficiency (20). A cutoff of 350 pg/ml (258 pmol/L) was used to identify vitamin B12 deficiency (21).

### Depression

Depressive symptoms were measured by using the 20-item Center for Epidemiological Studies Depression Scale (CES-D) (22). Participants with a CES-D score  $\geq 16$  were considered to be depressed (23,24). In addition, to explore a potential trend between depressive symptom severity and anemia, we defined four participant groups by CES-D score, based on cutoffs previously reported (25): (1) few or no symptoms of depression (CES-D score  $< 16$ ) ( $n = 673$ ), (2) mild depressive symptoms (score = 16–20) ( $n = 126$ ), (3) moderate depressive symptoms (score = 21–26) ( $n = 96$ ), and (4) severe depression (score  $> 26$ ) ( $n = 91$ ).

### Covariates

A Mini-Mental State Examination (MMSE) score below 18 was used to define cognitive impairment (26,27). The presence of comorbid conditions was determined using an adjudication process that included self-reported history, medical examination data, and medical records information (18). Kidney disease was determined by a serum creatinine level above 1.5 mg/dl in women and above 1.7 mg/dl in men. Based on tertiles of interleukin-6, participants were classified in three groups. Assessment of daily caloric and protein intake was performed based on the questionnaire of the European Prospective Investigation into Cancer and Nutrition (28). Drugs were coded according to the Anatomical Therapeutic and Chemical codes (29).

### Study Sample and Statistical Analyses

From the initial sample of 1156 participants, we excluded persons with missing data on hemoglobin level ( $n = 119$ , 10.3%) or CES-D score ( $n = 76$ , 6.6%). This selection resulted in a final sample size of 986 participants. In comparison to those who were excluded from study participation due to incomplete data ( $n = 170$ ), those included in the study were significantly younger ( $75 \pm 7$  vs  $82 \pm 9$  years,  $p < .001$ ) and tended to be less often female (56% vs 62%,  $p = .18$ ).

Differences between depressed and nondepressed participants in categorical parameters were tested using the chi-square test. Differences between continuous variables were assessed by analysis of variance comparisons for normally distributed parameters; alternatively, the Kruskal–Wallis test was adopted. To establish whether depression represented a risk factor for anemia, a logistic regression model was performed in the 986 participants included in this study. To explore a potential trend between depressive symptom severity and anemia, an additional logistic regression model was conducted, using categorization of the CES-D scores as described above. Variables considered for adjustment were

those associated with depression at  $p \leq .10$  in the univariate analyses. All analyses were performed using SPSS for Windows version 10.0 (Chicago, IL).

## RESULTS

### Participant Characteristics

Mean ( $\pm$  standard deviation) age of the 986 participants entering this study was 75 ( $\pm 7$ ) years, and 56% of the study population was female. The mean ( $\pm$  standard deviation) CES-D score in the total sample was 12.8 ( $\pm 8.7$ ) (range 0–48), and 313 (32%) of the study participants were depressed (CES-D score  $\geq 16$ ). Other characteristics of the study population are summarized in Table 1. Compared with nondepressed participants, those with depression were older, more likely to be female and enrolled in the Bagno a Ripoli site, less likely to be smokers and drinkers, had a lower education, had a higher prevalence of cognitive impairment, and presented a higher number of diseases. In particular, depression was associated with a higher prevalence of coronary artery disease, peripheral artery disease, congestive heart failure, and history of cancer. Finally, participants with depression were more likely using antidepressant medication, nonsteroidal anti-inflammatory drugs, and acetylsalicylic acid.

### Depression and Anemia

Anemia was identified in 101 (10%) study participants: 43 (10%) men and 58 (10%) women. As shown in Table 2, anemia was present in 48/313 (15%) participants with depression and in 53/673 (8%) participants without depression ( $p < .001$ ). After adjusting for potential confounders, depression was associated with a significantly increased risk of anemia (odds ratio [OR] = 1.93; 95% confidence interval [CI], 1.19–3.13). In the fully adjusted model, no significant interaction was observed between depression and sex ( $p = .30$ ) and between depression and age ( $p = .35$ ) on the risk of anemia.

The association between depression and anemia was persistent using a CES-D cutoff of 21 to define depression. Compared with participants scoring  $< 21$ , those with a score of 21 or higher were more likely to present with anemia (anemia among depressed participants: 30/187 [16%] vs anemia among nondepressed participants: 71/799 [9%];  $p = .004$ ), and this association was confirmed after adjusting for potential confounders (OR = 1.78; 95% CI, 1.06–3.00).

As shown in Table 2, depression was still significantly associated with anemia after exclusion of anemia related to iron (transferrin receptor/log[ferritin]  $< 1.5$ ) (OR = 2.11; 95% CI, 1.23–3.60) or vitamin B12 deficiency (OR = 1.96; 95% CI, 1.01–3.80). To explore whether poor health status or cognitive impairment could explain the association between depression and anemia, we repeated analyses after excluding participants with relevant comorbidities (such as congestive heart failure, kidney disease, history of cancer, or cognitive impairment). These analysis also yielded significant associations of depression with anemia.

Figure 1 summarizes the ORs for anemia across different groups, according to CES-D scores. After adjusting for

Table 1. Baseline Characteristics According to Depression Status

Characteristic	No Depression N = 673 (%)	Depression N = 313 (%)	p*
Age			<.001
<80 y	559 (83)	211 (67)	
≥80 y	114 (17)	102 (33)	
Female sex	322 (48)	232 (74)	<.001
Site			.06
Bagno a Ripoli	342 (51)	179 (56)	
Greve in Chianti	331 (49)	134 (43)	
Education			.06
0–5 y	480 (71)	244 (78)	
6–8 y	122 (18)	36 (12)	
9–13 y	47 (7)	20 (6)	
>13 y	23 (3)	13 (4)	
Body mass index			.23
<25 kg/m <sup>2</sup>	171 (25)	95 (30)	
25–29.9 kg/m <sup>2</sup>	340 (51)	173 (46)	
≥30 kg/m <sup>2</sup>	162 (24)	75 (24)	
Smoking			<.001
Never	349 (52)	227 (73)	
Former	241 (36)	56 (18)	
Current	83 (12)	30 (10)	
Alcohol use	443 (66)	171 (55)	.001
Cognitive impairment <sup>†</sup>	15 (2)	14 (5)	.05
No. of diseases, mean ± SD	2.6 ± 1.1	2.7 ± 1.2	.09
Diabetes	71 (10)	31 (10)	.76
Coronary artery disease	56 (8)	37 (12)	.08
Peripheral artery disease	48 (7)	32 (10)	.09
Congestive heart failure	24 (4)	22 (7)	.02
Stroke	38 (6)	23 (7)	.30
Pulmonary disease	45 (7)	20 (6)	.86
History of cancer	36 (5)	26 (8)	.08
Gastric disease	33 (5)	16 (5)	.89
Peptic ulcer	104 (16)	45 (14)	.66
Renal failure <sup>‡</sup>	6 (1)	5 (2)	.33
Albumin (g/dL), mean ± SD	4.2 ± 0.3	4.2 ± 0.3	.13
Energy intake (kcal/kg per day)	28.8 ± 8.4	28.0 ± 8.1	.15
Protein intake (g/kg per day)	1.1 ± 0.3	1.1 ± 0.3	.32
IL-6 (pg/mL) tertiles			.65
<1.03	217 (32)	110 (35)	
1.03–1.84	227 (34)	100 (32)	
≥1.85	228 (34)	101 (33)	
Use of antidepressants	13 (2)	24 (8)	<.001
Use of NSAIDs	122 (18)	90 (29)	<.001
Use of ASA	92 (14)	56 (18)	.08

Notes: Depression was defined as Center for Epidemiological Studies Depression Scale (CES-D) score ≥ 16.

\*p values are based on chi-square test for categorical variables, analysis of variance for continuous variables.

<sup>†</sup>Cognitive impairment was defined by Mini-Mental State Examination score < 18.

<sup>‡</sup>Renal failure was determined by a serum creatinine level above 1.5 mg/dL in women and above 1.7 mg/dL in men.

SD = standard deviation; NSAIDs = nonsteroidal anti-inflammatory drugs; ASA = acetylsalicylic acid.

potential confounders, the risk of anemia progressively and significantly increased as CES-D score increments increased (signifying more severe depression) (*p* for linear trend = .01). Compared with nondepressed participants whose CES-D score was <16, the ORs for anemia were 1.74, 2.04, and 2.10

Table 2. Association Between Depression and Anemia

Patients With	N	Anemia	p	OR (95% CI) for Anemia*
Total sample				
No depression	673	53 (8%)	<.001	1
Depression	313	48 (15%)		1.93 (1.19–3.13)
Sample without congestive heart failure, kidney disease, and history of cancer				
No depression	586	45 (8%)	.001	1
Depression	255	39 (15%)		2.06 (1.21–3.49)
Sample without cognitive impairment <sup>†</sup>				
No depression	658	49 (7%)	.001	1
Depression	299	43 (14%)		1.93 (1.17–3.19)
Sample without iron deficiency anemia <sup>‡</sup>				
No depression	657	37 (6%)	<.001	1
Depression	305	40 (13%)		2.11 (1.23–3.60)
Sample without vitamin B12 deficiency anemia <sup>¶</sup>				
No depression	644	24 (4%)	.003	1
Depression	289	24 (8%)		1.96 (1.01–3.80)

Notes: \*Adjusted for age, sex, site, education, smoking, alcohol use, cognitive impairment, number of diseases, coronary artery disease, peripheral artery disease, congestive heart failure, history of cancer, use of antidepressants, use of nonsteroidal anti-inflammatory drugs, and use of acetylsalicylic acid.

<sup>†</sup>Cognitive impairment was defined by Mini-Mental State Examination score < 18.

<sup>‡</sup>Iron deficiency was defined by (serum transferrin receptor/log[ferritin] < 1.5).

<sup>¶</sup>Vitamin B12 deficiency was defined by vitamin B12 < 350 pg/ml (258 pmol/L).

OR = odds ratio; CI = confidence interval; CES-D = Center for Epidemiological Studies Depression Scale. Depression was defined as CES-D ≥ 16; anemia was defined as Hb < 12 g/dL for women and < 13 g/dL for men.

for participants with mild (score = 16–20), moderate (score = 21–26), and severe depression (score > 26), respectively.

## DISCUSSION

Based on our findings among persons living in the community, depression appears to be associated with anemia, irrespective of comorbidity and other potential confounders. This association increases with severity of depressive symptoms.

How could we explain a link between depression and anemia? It has been hypothesized that psychological distress can activate neuroregulated biological processes. This activation can result in a diminished ability to combat pathological processes, thus favoring the onset of negative outcomes such as anemia. This phenomenon, described by Engel (30) as the “giving-up–given-up complex,” could explain the increased risk of adverse outcomes observed among depressed participants. According to this hypothesis, depression adversely affects cardiac, gastrointestinal, endocrine, neurologic, and immune processes by increasing sympathetic tone and decreasing vagal tone (31). More specifically, the higher risk of anemia that we observed among depressed participants may be due to a number of factors. First, psychological distress has been associated with an increased risk of occult bleeding, which can be responsible for a higher prevalence of anemia among depressed persons (32). Second, the increased sympathetic




Figure 1. Odds ratio for anemia according to Center for Epidemiological Studies-Depression Scale (CES-D) score (CES-D <16 is the reference group). Analysis is adjusted for age, sex, site, education, smoking, alcohol use, cognitive impairment, number of diseases, coronary artery disease, peripheral artery disease, congestive heart failure, history of cancer, use of antidepressants, use of nonsteroidal anti-inflammatory drugs, and use of acetylsalicylic acid.

tone, observed among depressed persons (31), can regulate bone marrow microenvironment, affecting erythropoiesis (33). Third, anemia can represent a consequence of malnutrition among depressed persons (16). However, the fact that daily caloric and protein intake did not differ significantly according to depressive status makes this explanation unlikely.

Another possible explanation for our findings is that depressive symptoms may occur as a consequence of anemia. Anemia may directly affect brain function. Some studies conducted in hemodialysis patients (12,13) have shown that anemia appears to contribute to impaired cognitive function and that persons with chronic renal failure frequently suffer from confusion, inability to concentrate, decreased mental alertness, and impaired memory. In addition, acute isovolemic anemia results in subtle slowing of data-processing ability in humans and degrades memory (12). Previous studies conducted among hemodialysis patients (13,34) have shown that correction of renal anemia with erythropoietin improves quality of life, cognition, and depressive symptoms. This improvement of psychological symptoms seems to be mediated by an increment of oxygen delivery to the brain and particularly to frontotemporal and subcortical brain structures, which are involved in the depression pathway (35,36). Finally, fatigue, which is a common symptom of anemia, can have detrimental effects on a person's quality of life and determine the onset of depressive symptoms (15,16).

An alternative explanation to our findings is that the link between anemia and depression is caused by underlying disease. Although restricted analyses among healthier participants did not change our results, we can not rule out that (unmeasured) diseases affected our results.

The prevalence of depression in our study is higher compared with other observations. A possible explanation for this finding is the low socioeconomic status of this population (as suggested by the low educational status of participants in this study), a factor that has been directly associated with depression (37). However, the prevalence of

depression in this study did not differ substantially from that observed in a general population in Northern Italy (38).

Depressive symptoms have been associated with vitamin B12 deficiency, a factor potentially related to anemia (39). Therefore, we repeated the analysis after excluding cases of anemia related to vitamin B12 deficiency, and achieved results similar to those from the whole sample.

The present study has several limitations. We did not explicitly determine the underlying cause of anemia. Therefore, we may have missed the presence of some other diseases or factors such as liver or pancreatic disease, inflammatory bowel disease, and pernicious anemia, which could have explained some of the findings. In addition, although we adjusted all analyses for several potential confounders, including common chronic conditions, it is possible that anemia is simply an indicator of poor health status, and that its association with depression is illustrative of that. Finally, because of the cross-sectional design of the study, we were not able to identify a causal relationship between anemia and depression.

### Conclusion

Our findings indicate that depression, possibly independent of its underlying cause, is associated with anemia in a general population of older persons living in the community. Future longitudinal studies should confirm and explain the reason for this association.

### ACKNOWLEDGMENTS

The InCHIANTI study was supported as a "targeted project" (ICS 110.1/RS97.71) by the Italian Ministry of Health and in part by the U.S. National Institute on Aging (contracts 263 MD 9164 13 and 263 MD 821336). The work of Drs. Penninx, Cesari, and Pahor is supported by the U.S. National Institutes of Health (Claude Pepper Older Americans Independence Center P30AG021332-01).

Conflict of Interest: Data analyses were in part supported by Ortho Biotech Products, L.P., Bridgewater NJ. Drs. Penninx, Pahor, and Ferrucci have served as consultants for Ortho Biotech Products.

Address correspondence to Graziano Onder, MD, Centro Medicina dell'Invecchiamento, Università Cattolica del Sacro Cuore, Policlinico A. Gemelli, L.go Francesco Vito 1, 00168 Rome, Italy. E-mail: graziano\_onder@rm.unicatt.it

### REFERENCES

- Kessler RC, Berglund P, Demler O, et al. National Comorbidity Survey Replication. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003;289:3095-3105.
- Blazer DG. Depression in late life: review and commentary. *J Gerontol A Biol Sci Med Sci*. 2003;58:249-265.
- Ruo B, Rumsfeld JS, Hlatky MA, Liu H, Browner WS, Whooley MA. Depressive symptoms and health-related quality of life: the Heart and Soul Study. *JAMA*. 2003;290:215-221.
- Onder G, Penninx BW, Landi F, et al. Investigators of the Gruppo Italiano di Farmacoepidemiologia nell'Anziano Study. Depression and adverse drug reactions among hospitalized older adults. *Arch Intern Med*. 2003;163:301-305.
- Rozzini R, Frisoni GB, Sabatini T, Trabucchi M. The association of depression and mortality in elderly persons. *J Gerontol A Biol Sci Med Sci*. 2002;57:M144-M145.
- Penninx BW, Geerlings SW, Deeg DJ, van Eijk JT, van Tilburg W, Beekman AT. Minor and major depression and the risk of death in older persons. *Arch Gen Psychiatry*. 1999;56:889-895.

7. Blazer DG, Hybels CF, Pieper CF. The association of depression and mortality in elderly persons: a case for multiple, independent pathways. *J Gerontol A Biol Sci Med Sci*. 2001;56:M505–M509.
8. Penninx BW, Guralnik JM, Ferrucci L, Simonsick EM, Deeg DJ, Wallace RB. Depressive symptoms and physical decline in community-dwelling older persons. *JAMA*. 1998;279:1720–1726.
9. Salive ME, Comoni-Huntley J, Guralnik JM, et al. Anemia and hemoglobin levels in older persons: relationship with age, gender, and health status. *J Am Geriatr Soc*. 1992;40:489–496.
10. Ania BJ, Suman VJ, Fairbanks VF, Rademacher DM, Melton LJ III. Incidence of anemia in older people: an epidemiologic study in a well defined population. *J Am Geriatr Soc*. 1997;45:825–831.
11. Thomas DR. Anemia and quality of life: unrecognized and undertreated. *J Gerontol A Biol Sci Med Sci*. 2004;59:238–241.
12. Nissenson AR. Epoetin and cognitive function. *Am J Kidney Dis*. 1992;20(Suppl. 1):21–24.
13. Pickett JL, Theberge DC, Brown WS, Schweitzer SU, Nissenson AR. Normalizing hematocrit in dialysis patients improves brain function. *Am J Kidney Dis*. 1999;33:1122–1130.
14. Cesari M, Penninx BW, Lauretani F, et al. Hemoglobin levels and skeletal muscle: results from the InCHIANTI study. *J Gerontol A Biol Sci Med Sci*. 2004;59:249–254.
15. Penninx BW, Pahor M, Cesari M, et al. Anemia is associated with disability and decreased physical performance and muscle strength in the elderly. *J Am Geriatr Soc*. 2004;52:719–724.
16. Mittrache C, Passweg JR, Libura J, et al. Anemia: an indicator for malnutrition in the elderly. *Ann Hematol*. 2001;80:295–298.
17. Katz IR, Beaton-Wimmer P, Parmelee P, Friedman E, Lawton MP. Failure to thrive in the elderly. Exploration of the concept and delineation of psychiatric components. *J Geriatr Psychiatry Neurol*. 1993;6:161–169.
18. Ferrucci L, Bandinelli S, Benvenuti E, et al. Subsystems contributing to the decline in ability to walk: bridging the gap between epidemiology and geriatric practice in the InCHIANTI study. *J Am Geriatr Soc*. 2000;48:1618–1625.
19. World Health Organization (WHO). Nutritional Anaemias. Report of a WHO scientific group. 1968. Geneva, Switzerland: World Health Organization. Technical Report Series No. 405.
20. Punnonen K, Irjala K, Rajamaki A. Serum transferrin receptor and its ratio to serum ferritin in the diagnosis of iron deficiency. *Blood*. 1997;89:1052–1057.
21. Lindenbaum J, Rosenberg IH, Wilson PW, Stabler SP, Allen RH. Prevalence of cobalamin deficiency in the Framingham elderly population. *Am J Clin Nutr*. 1994;60:2–11.
22. Myers JK, Weissman MM. Use of a self-report symptom scale to detect depression in a community sample. *Am J Psychiatry*. 1980;137:1081–1084.
23. Weissman MM, Sholomskas D, Pottenger M, Prusoff BA, Locke BZ. Assessing depressive symptoms in five psychiatric populations: a validation study. *Am J Epidemiol*. 1977;106:203–214.
24. Radloff LS. The CES-D Scale: a self-report depression scale for research in the general population. *Appl Psych Meas*. 1977;1:385–401.
25. Unutzer J, Patrick DL, Marmon T, Simon GE, Katon WJ. Depressive symptoms and mortality in a prospective study of 2,558 older adults. *Am J Geriatr Psychiatry*. 2002;10:521–530.
26. Folstein MF, Folstein SE, McHugh PR. “Mini-mental state”. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12:189–198.
27. Malloy PF, Cummings JL, Coffey CE, et al. Cognitive screening instruments in neuropsychiatry. A report of the Committee on Research of the American Neuropsychiatric Association. *J Neuropsychiatry Clin Neurosci*. 1997;9:189–197.
28. Bartali B, Turrini A, Salvini S, et al. Dietary intake estimated using different methods in two Italian older populations. *Arch Gerontol Geriatr*. 2004;38:51–60.
29. Pahor M, Chrischilles EA, Guralnik JM, Brown SL, Wallace RB, Carbonin P. Drug data coding and analysis in epidemiological studies. *Eur J Epidemiol*. 1994;10:405–411.
30. Engel GL. A life setting conducive to illness. The giving-up–given-up complex. *Ann Intern Med*. 1968;69:355–365.
31. Cacioppo JT, Burleson MH, Poehlmann KM, et al. Autonomic and neuroendocrine responses to mild psychological stressors: effects of chronic stress on older women. *Ann Behav Med*. 2000;22:140–148.
32. Peura DA. Stress-related mucosal damage: an overview. *Am J Med*. 1987;83:3–7.
33. Maestroni GJ. Neurohormones and catecholamines as functional components of the bone marrow microenvironment. *Ann N Y Acad Sci*. 2000;917:29–37.
34. Moreno F, Sanz-Guajardo D, Lopez-Gomez JM, Jofre R, Valderrabano F. Increasing the hematocrit has a beneficial effect on quality of life and is safe in selected hemodialysis patients. Spanish Cooperative Renal Patients Quality of Life Study Group of the Spanish Society of Nephrology. *J Am Soc Nephrol*. 2000;11:335–342.
35. Kusunoki M, Kimura K, Nakamura M, Isaka Y, Yoneda S, Abe H. Effects of hematocrit variations on cerebral blood flow and oxygen transport in ischemic cerebrovascular disease. *J Cereb Blood Flow Metab*. 1981;1:413–417.
36. Oda K, Okubo Y, Ishida R, et al. Regional cerebral blood flow in depressed patients with white matter magnetic resonance hyperintensity. *Biol Psychiatry*. 2003;53:150–156.
37. Lorant V, Deliege D, Eaton W, Robert A, Philippot P, Ansseau M. Socioeconomic inequalities in depression: a meta-analysis. *Am J Epidemiol*. 2003;157:98–112.
38. Minicuci N, Maggi S, Pavan M, Enzi G, Crepaldi G. Prevalence rate and correlates of depressive symptoms in older individuals: the Veneto Study. *J Gerontol A Biol Sci Med Sci*. 2002;57:M155–M161.
39. Penninx BW, Guralnik JM, Ferrucci L, Fried LP, Allen RH, Stabler SP. Vitamin B(12) deficiency and depression in physically disabled older women: epidemiologic evidence from the Women’s Health and Aging Study. *Am J Psychiatry*. 2000;157:715–721.

Received July 1, 2004

Accepted August 4, 2004

Decision Editor: John E. Morley, MB, BCh