

26 freshwater ecotoxicity. The most important source of harmful environmental impacts in
27 all the categories under assessment was the production of the hen feed and, to a lesser
28 extent, the purchase of new laying hens to replace the old ones. On the contrary, water
29 consumption and the employment of chemicals for cleaning barely influenced the
30 impact. One aspect that was noteworthy was the beneficial effect on environmental
31 impact produced by the sale of old laying hens for meat production, especially on the
32 urban land occupation and metal depletion categories. Additionally, the carbon footprint
33 of egg production was calculated and a value of 2.66 kgCO₂eq per dozen eggs was
34 obtained. Environmental improvement actions should be directed mainly towards
35 optimizing the hen feed formulation, not only from an economic perspective, but also
36 considering the environmental aspects involved.

37

38 **Keywords:** LCA, laying hen farm, egg production, environmental impact, carbon
39 footprint.

40

41 **1. INTRODUCTION**

42 The production of eggs worldwide has been increasing during recent decades.
43 According to the FAO, in 2013 the global production of eggs had reached a volume of
44 about 68 million tonnes (FAO, 2016). The European Union produces approximately 7
45 million tonnes of useable eggs per annum. Specifically, France and Spain are the largest
46 egg producers (accounting for approximately a quarter of European production)
47 (MAPAMA, 2017).

48 Food production requires large amounts of energy, which implies several
49 negative environmental impacts such as greenhouse gas (GHG) emissions. In addition,
50 since consumers in developed countries have started to demand high-quality food,

51 produced under more environmentally friendly conditions (González-García et al.,
52 2014), producers confront the contradictory demands of the need to increase food
53 production while having to reduce the ecological impact of intensive production
54 methods (Darnhofer et al., 2016). So, as occurs with other food industries, commercial
55 egg production faces the challenge of producing high quality products in a way that
56 meets consumer expectations, satisfies environmental regulations and maximizes
57 profitability (Freeman et al., 2009). Moreover, egg producing farms are included in the
58 Best Available Techniques (BAT) Reference Document for the Intensive Rearing of
59 Poultry and Pigs contained in the Industrial Emissions Directive (IED, 2010/75/EU)
60 issued by the European IPPC Bureau. Nevertheless, it was not until the 1980s that the
61 environmental impact of intensive livestock farming was considered a problem.
62 Awareness of the implications of farming activities such as the contamination of soil
63 due to excess manure application and its impact on soil and water quality have increased
64 over the years. Hence, the environmental impacts of agriculture and animal production
65 have been increasingly acknowledged (Paolotti et al., 2016).

66 The poultry industry is one of the largest and most developed of the existing
67 industries in the agriculture sector (Ghasempour and Ahmadi, 2016). In Spain in 2017
68 there were 1260 egg producing farms and the average number of hens per egg
69 production facility was 67,700. During the last few years, the tendency in Spain has
70 been to increase the number of hens housed in cages, which now represent 93% of total
71 laying hens (MAPAMA, 2017). Considering all EU countries, in contrast, this
72 percentage is much lower (40%), since free range production facilities are becoming
73 more widespread due to public concern for animal welfare (Leenstra et al., 2014).

74 Life cycle assessment (LCA) is defined as a method for assessing environmental
75 aspects and potential impacts associated with a product (Calderón et al., 2010; Calderón

76 et al., 2017; Iglesias et al., 2012). It has been demonstrated to be a worthwhile tool for
77 quantifying resource use and emissions in a wide range of primary sectors such as meat
78 production (Cederberg, 2014; Velarde et al., 2015) and dairy farms (Hospido et al.,
79 2003) and also in industrial sectors (Tecco et al., 2016; Vázquez-Rowe et al., 2012). In
80 addition, the food system produces a large amount of GHG, specifically 33% of
81 anthropogenic carbon emissions (Zhu et al., 2017). Furthermore, recently, the carbon
82 footprint has been employed as a global measure of the production performance of
83 different foodstuffs (Casolani et al., 2016).

84 There are papers targeting different aspects of the poultry meat chain (Cesari et
85 al., 2017; Da Silva et al., 2014; González-García et al., 2014; Kalhor et al., 2016;
86 Skunca et al., 2015; Wiedemann et al., 2017), but there is an evident lack of studies
87 involving a life-cycle assessment approach for the environmental performance of egg
88 production in egg producing farms. In fact, there are very few published studies
89 regarding egg production (Cederberg et al., 2009; Dekker et al., 2011; Ghasempour and
90 Ahmadi, 2016; Leinonen et al., 2012; Mollenhorst et al., 2006; Pelletier et al., 2013;
91 Pelletier 2017). Thus, the aim of this study was to analyse the environmental
92 performances of egg production in a laying hen farm in Asturias (a region in NW
93 Spain), which has been selected as being representative of intensive European egg
94 production. An LCA has been carried out in order to quantify its environmental impact,
95 and to identify the activities with a major environmental impact, which would permit
96 the establishment of a series of actions aimed at improvement of the situation.
97 Additionally, the carbon footprint of egg production was also calculated.

98

99 **2. MATERIALS AND METHODS**

100

101 **2.1. LCA**

102 **2.1.1 Objectives and functional unit definition**

103 In this study, LCA methodology was used as a tool with the objective of
104 determining the environmental impact of a Spanish-type laying hen farm. The
105 functional unit was the annual egg production in 2015 (1,3344,000 eggs).

106 **2.1.2 System description and boundaries**

107 The laying hen farm involved in this research is situated in northern Spain
108 (Asturias). The facility, which houses 55,000 laying hens, consists of two industrial
109 units of 1540 m² and 1430 m², respectively. One of the units is also used as a storehouse
110 for egg packing materials. In addition, an industrial unit of 500 m² accommodates an
111 egg-sorting room, an office and a toilet. The facilities are not connected to the municipal
112 sewage system, so wastewater is stored and removed by an authorized company.

113 Laying hens employed in this farm are hybrids (Rhode Island Red/Light Sussex
114 cross), medium sized (average weight 2.1 kg), of brown colour with some soft white
115 feathers. Following the ban on conventional cages for laying hens in the EU (Council
116 Directive 1999/74/EC), hens are housed in suspended wire cages placed in four tiers
117 along the length of each industrial unit (Big Dutchman EUROVENT-EV 1250a - EU -
118 60[®]). Sixteen-week-old hens are purchased and they are exploited for 75-80 weeks.
119 After their productive life, exhausted hens are replaced by new laying hens and the old
120 hens are sold for meat production. In 2015 all laying hens were removed and replaced
121 with new ones. Hens are fed with commercial fodder for laying hens (see Table 1 for
122 nutritional data) via automatic feed delivery systems and have continuous access to
123 water supplied from nipple drinkers (6 stainless steel nipples per compartment). Eggs
124 are collected daily by automatic belts, moved to the end of each industrial unit and then

125 to a common egg-sorting room where they are packaged in recycled cardboard boxes
126 and trays.

127 Polypropylene belts beneath the bottom wires collect the manure that is dried by
128 means of an air duct (dry matter content of up to 60%). The dried manure is removed
129 twice a week and loaded directly onto a truck that carries it to a facility which
130 commercialise it as fertilizer.

131 The system considered included the whole life cycle involved in the production
132 of the eggs: transportation, consumption of energy and water, waste management and
133 emissions.

134 **2.1.3. Inventory analysis**

135 Data were mainly collected through personal interviews with farmers.
136 Additionally, some information was obtained from bibliographic sources. Inventory
137 data have been organized into the subsystems shown in Figure 1 and they are
138 summarized in Table 2. The following considerations were taken into account for the
139 inventory analyses:

- 140 • With respect to packaging and fodder, only those elements that exceed
141 5% (w/w) of the total were included, so the polyethylene around the
142 pallets used to transport packaging materials was not considered (<
143 0.05% w/w).
- 144 • Regarding cleaning materials, only bleach was included in the inventory,
145 since it was the main cleaning agent employed (> 90%).
- 146 • Drugs were not taken into consideration since they are only occasionally
147 employed and, in addition, the amounts of these medicinal substances
148 used in the farm were insignificant compared to the total incomes and
149 outcomes.

- 150 • Transport of raw materials (fodder, cleaning products, packaging
151 material), new and old laying hens, wastes and eggs (from the farm to the
152 retail store) was considered. Eggs are transported at room temperature in
153 vans belonging to the farm. Data for these subsystems were included as
154 tkm for external transport and as the diesel consumed by the company
155 vans for internal transport (eggs). Note that the transport of the
156 ingredients for the fodder has not been considered in this study.
- 157 • Wastewater was not generated by cleaning operations in the facilities
158 since they used a dry cleaning technique employing compressed air.
159 Therefore, it was assumed that 10% of consumed water was employed
160 for the cleaning of transport vehicles and human uses (office toilet) and
161 this water was considered as wastewater. It was supposed that the rest of
162 the incoming water was consumed by the hens as drinking water.
- 163 • Emissions were calculated according to the stock sector PRTR emission
164 factors (EPER-Spain): 0.0318 kg NH₃-N, 0.007642 kg N₂O-N and
165 0.08730 kg CH₄, given in all cases per hen and year (MAPAMA, 2017).
- 166 • Exhausted laying hens were sold for slaughtering and the meat is sold for
167 human consumption. Hence, poultry meat is included in the system as an
168 avoided product. The live weight of slaughtered hens was calculated
169 supposing that each hen weighed 2.1 kg and that 96% of replaced hens
170 were sold for slaughter. The remaining 4% were the hens that die and
171 were managed as waste.

172 **2.1.4 Impact assessment**

173 Impact assessment was performed with the LCA software package SimaPro v7,
174 using the ReCiPe Midpoint (H) V1.12 / Europe Recipe H method. This method includes

175 18 impact category indicators (climate change, ozone depletion, terrestrial acidification,
176 freshwater eutrophication, marine eutrophication, human toxicity, photochemical
177 oxidant formation, particulate matter formation, terrestrial ecotoxicity, freshwater
178 ecotoxicity, marine ecotoxicity, ionising radiation, agricultural land occupation, urban
179 land occupation, natural land transformation, water depletion, metal depletion and fossil
180 depletion) to reach wide impact category coverage and follows the latest
181 recommendations of the LCA community (Heinonen et al., 2016). The ReCiPe
182 Endpoint (H) V1.12 / Europe Recipe H method has also been employed with the aim of
183 classifying the damage in only three categories: human health, ecosystems and
184 resources. The ReCiPe method takes its origins from CML and Ecoindicator (Baldini et
185 al., 2017) and it has been applied recently in different LCA studies focused on agro-
186 food industries (Freón et al., 2014; Vázquez-Rowe et al., 2014; Arzoumanidis et al.,
187 2017; Baldini et al., 2017; Khatri and Jain, 2017; Noya et al., 2017). The advantages of
188 this method include (i) the broadest set of midpoint impact categories and (ii) the use of
189 impact mechanisms that have global scope (Santos et al., 2017).

190 Data for the fodder subsystem were obtained from Agri-footprint (maize,
191 soybean, palm oil) and LCA Food (sodium bicarbonate) databases. Data for new laying
192 hens, exhausted laying hens for slaughtering, cleaning elements and water subsystems
193 were obtained from the Agri-footprint database. Data for packaging material, electricity,
194 wastes and emissions to air subsystems were taken from the EcoInvent database.
195 Transport subsystem data were obtained from Agri-footprint (transport by track) and
196 EcoInvent (diesel). Whenever it was possible “Alloc Def”, which follows the
197 attributional approach in which burdens are attributed proportionally to specific
198 processes, was used. Additionally, regions and time span were selected considering the
199 available information regarding the system studied.

200

201 **2.2 Carbon Footprint**

202 The carbon footprint was obtained by employing the Greenhouse Gas Protocol
203 V1.01 / CO₂ eq (kg), again by means of the LCA software package SimaPro v7. This
204 method includes scopes 1 (all direct GHG emissions), 2 (indirect GHG emissions from
205 consumption of purchased electricity, heat or steam) and also 3 (other indirect
206 emissions, such as transport-related activities, waste disposal, etc.). In addition, it is the
207 same method of analysis employed by the Spanish Ministry of Agriculture and Fishing,
208 Food and Environment (MAPAMA, 2017).

209

210 **2.3 Alternative scenarios**

211 Two alternative scenarios to the real analysis (scenario 1) have been considered.
212 As hen feeding turned out to be the most impacting subsystem, in scenario 2, pea was
213 substituted for soybean, whereas, in scenario 3, the analysis examined the replacement
214 of palm oil with cottonseed oil. The criteria followed for the substitution of ingredients
215 have involved using the same mass of products for both scenarios.

216

217 **3. RESULTS AND DISCUSSION**

218

219 **3.1. Impact assessment**

220 Characterization results obtained with the ReCiPe Midpoint method revealed
221 hen feed to be the subsystem with the highest environmental loads in almost all
222 categories considered (Figure 2). Specifically, feed was responsible of more than 55%
223 of impact in all categories evaluated, excepting metal depletion (the contribution of feed
224 here was 11%) and urban land occupation (feed did not contribute to this category).

225 Animal feed was responsible for more than 90 % of impact in terrestrial ecotoxicity and
226 natural land transformation categories. Additionally, it must be mentioned that the feed
227 conversion ratio of the farm under study was $2.8 \text{ kg}_{\text{feed}} / \text{kg}_{\text{egg}}$, a value slightly higher
228 than those found by Dekker et al. (2011) in loose housing systems (2.3-2.6) and in
229 battery cage systems (2.0). It should be taken into account that the non-enriched battery
230 cage systems studied in the previous investigation have been prohibited since 1 January
231 2012 by the Council Directive 1999/74/EC due to animal welfare concerns. On the
232 contrary, the farm studied here fulfils the minimum requirement of 750 cm^2 of cage area
233 per hen fixed by European regulations, so a direct comparison cannot be established.

234 The breeding of new hens that were purchased in 2015 was also an activity with
235 high impact, affecting all categories except for urban land occupation. The categories
236 most affected by this subsystem were particulate matter formation and terrestrial
237 acidification (new hens are responsible for approximately 24% of the total impact in
238 these categories).

239 It is noteworthy that the production of packaging materials was responsible for
240 76 % of the harmful impact in the metal depletion category and almost all the harmful
241 impact in the urban land occupation category. This subsystem also made a percentage
242 contribution higher than 10% to ionising radiation, marine ecotoxicity, human toxicity
243 and ozone depletion.

244 The contribution of transport to ozone depletion, photochemical oxidant
245 formation and fossil depletion categories was 18%, 5% and 4%, respectively. The
246 contribution of waste management to marine ecotoxicity was also noticeable (16%). Gas
247 emissions contributed 9% to terrestrial acidification, 7% to particulate matter formation
248 and 5% to climate change, whereas electricity consumption was responsible for 12%
249 and 7% of ionising radiation and metal depletion, respectively. In this sense, it should

250 be kept in mind that the use of radioactive material within nuclear reactors to generate
251 electricity generates ionising radiation and in Spain nuclear energy is one of the main
252 sources of electricity (MINETAD, 2017).

253 Finally, it is worth noting that some subsystems made a beneficial contribution
254 in some categories. Specifically, the sale of exhausted laying hens for slaughter and also
255 waste management had favourable effects on almost all categories, mainly on urban
256 land occupation and metal depletion. Indeed, in these categories the global impact was
257 seen to be advantageous to the environment. This can easily be explained, since the use
258 of discarded laying hens for meat avoids the need to breed chickens raised with this
259 specific aim, and it is well known that, among all animal products, the largest
260 environmental impacts are usually associated with meat production (Xu and Lan, 2016).
261 Farm processes were identified as the main contributors to the environmental impacts
262 derived from chicken meat production. Specifically, along the production chain, broiler
263 fattening is the phase that has most impact. On the contrary, hatchery, slaughterhouse
264 and packaging have a low impact (Cesari et al., 2017; González-García et al., 2014). In
265 addition, the reason for the beneficial effect of the waste subsystem observed in this
266 work was mainly due to the recycling of waste cardboard, which avoided the
267 consumption of virgin materials.

268 The normalization phase allows the comparison of all environmental impacts
269 using the same scale. According to these outcomes, natural land transformation was the
270 most prominent category, although terrestrial ecotoxicity was also of importance. The
271 rest of the studied categories were less affected in comparison with those mentioned
272 above.

273 As previously commented, the main contributor to natural land transformation
274 was hen feeding, the soybean used as an ingredient in the fodder being responsible for

275 69% of the contribution of hen feeding to this category, whereas palm oil contributed
276 22%. Again, hen feeding was the main subsystem with a harmful impact on terrestrial
277 and freshwater ecotoxicity categories. However, in these cases, palm oil employed as an
278 ingredient in the fodder originated approximately 60% of the impact, followed by
279 soybean (around 30%).

280 In Europe, soy and palm crops are partly or wholly produced overseas. Soy is
281 imported mainly from Brazil and Argentina, where forest areas are being converted into
282 agricultural land. In addition, the production of palm oil also implies land-use changes.
283 Leinonen et al. (2012) reported that the use of soy and palm for feeding laying hens
284 contributes notably to the global warming category, as a result of greenhouse gases
285 being released by changes in land use.

286 It is clear that soybean cultivation is linked to serious environmental problems.
287 In Brazil, soybean production has expanded rapidly in recent decades, whilst in
288 Malaysia oil palm plantations are also expanding, sometimes with the sacrifice of
289 invaluable rain forest (Mattsson et al., 2000). In addition, palm oil production is deeply
290 related with forest transformation and land availability (Uusitalo et al., 2014). Hence, in
291 many regions the production of these crops is a major cause of land use, but in addition,
292 the use of glyphosate in palm oil production in Thailand notably contributes to
293 freshwater ecotoxicity (Saswattecha et al., 2015).

294 The ReCiPe endpoint method allows all the impacts to be grouped into only
295 three categories: human health, ecosystems and resources. As is shown in Figure 3, the
296 ecosystems category was the most affected in the long-term. Again, in this category
297 food was responsible for 81% of impact, whereas 9% was originated by acquisition of
298 new laying hens. With respect to human health, these subsystems were responsible for

299 79% and 13% of impact, respectively. Resources were again affected by food (71%)
300 and the purchase of young chicks (14%), but also by packaging materials (11%).

301 These results are in agreement with those reported regarding the environmental
302 impacts related to egg production in Iran (Ghasempour and Ahmadi, 2016) and also in
303 Canada (Pelletier, 2017). In both cases, the composition and amount of feed consumed
304 in egg producing facilities were found to be the largest contributors to harmful impacts.

305

306 **3.2. Carbon footprint**

307 Results obtained from the Green House Gas Protocol are shown in Figure 4. In
308 the carbon footprint of the egg producing farm analysed here, scope 1 included the
309 direct emissions that correspond to N₂O and CH₄ originated by hen housing and manure
310 storage and also the emission derived from the diesel employed in product
311 commercialization, since commercialization is carried out by the farm. Scope 2 also
312 included emissions derived from electricity use. Considering scopes 1+2, the farm had a
313 carbon footprint of 342 t CO₂ eq during 2015, i.e., 308 g CO₂ eq per dozen eggs. The
314 major factor responsible for this value was direct emissions of N₂O and CH₄, which
315 contributed approximately to the same degree. Scope 3 considers extraction and
316 production of materials, management and treatment of generated wastes by the
317 management company and transport activity carried out by personnel external to the
318 farm itself. Considering only fossil and biogenic carbon, according to the ISO 14067
319 standard, when scope 3 is included a value of 2960 t CO₂ eq was achieved for the year
320 2015. As was expected, hen feeding was again the main contributor to greenhouse gas
321 emission (73%), as was also found by other authors (Cederberg et al., 2009;
322 Ghasempour and Ahmadi, 2016; Pelletier et al., 2013; Xu and Lan, 2016).

323 This value corresponded to a carbon footprint of 2.66 kgCO₂ eq per dozen eggs
324 (approximately 3.4 kgCO₂ eq per kg of eggs). This is a value within the range reported
325 by Nijdam et al. (2012) (1.7-5.5 kg CO₂ eq for 1 kg of eggs) for the egg industry in
326 Canada, England and Wales, but lower than the range found by Mollenhorst et al.
327 (2016) for egg production systems in Netherlands (3.9-4.6 kg CO₂ eq for 1 kg of eggs),
328 Pelletier et al. (2013) for packaged shell eggs in Iowa (4.2-6.0 kg CO₂ eq for 1 kg of
329 eggs) and the value reported by Ghasempour and Ahmadi (2016) for egg production in
330 Iran (4.07 kg CO₂ eq for 1 kg of eggs). The lowest published value for the carbon
331 footprint for eggs at the farm-gate, corresponding to 1.4 kg CO₂ eq per kg egg, was
332 described by Cederberg et al. (2009) for Swedish production systems.

333 These values help to confirm that eggs, together with milk (1-2 kg CO₂ eq kg⁻¹)
334 and chicken meat (2-6 kg CO₂ eq kg⁻¹), turn out to be the animal products that cause
335 least greenhouse emissions, especially if they are compared with foodstuffs such as beef
336 meat (9-129 kg CO₂ eq kg⁻¹), pork meat (4-11 kg CO₂ eq kg⁻¹), lamb meat (10-150 kg
337 CO₂ eq kg⁻¹), cheese (6-22 kg CO₂ eq kg⁻¹) or shellfish (1-86 kg CO₂ eq kg⁻¹) (Del
338 Prado et al., 2013; Ghasempour and Ahmadi, 2016; Nijdam et al., 2012).

339

340 **3.3. Improvement actions**

341 As mentioned above, the activity responsible for most environmental impacts
342 derived from intensive egg production in Spain was found to be the production of the
343 hen feed, which is in accordance with results reported for egg production in the
344 Netherlands, the United Kingdom, Iran and Canada (Dekker et al., 2011; Leinonen et
345 al., 2012; Ghasempour and Ahmadi, 2016; Pelletier, 2017). Similar conclusions were
346 reached for broiler chicken production (González-García et al., 2014; Da Silva et al.,
347 2014).

348 Since soybean and palm oil employed as ingredients in the fodder used in the
349 farm are mainly responsible for the impacts caused by hen feeding, the first
350 environmental improvement that should be tried is the total or partial replacement of
351 these ingredients in the fodder formulations. For instance, pea or bean can be employed
352 instead of soybean, since both these crops have been tested successfully as ingredients
353 in laying hen feed formulation (Koivunen et al., 2014; Koivunen et al., 2015). In the
354 same way, palm oil could be substituted with other crop oils such as cotton, corn, flax,
355 canola, olive or sunflower oils, which are often employed as components of fodders for
356 laying hens (Balevi and Coskun, 2000; Ceylan et al., 2011, Yuan et al., 2014). As an
357 example, two alternative scenarios (2 and 3) have been considered. In scenario 2,
358 soybean was replaced with pea and, in scenario 3, the analysis was carried out
359 substituting cottonseed oil for palm oil. It has been reported that the inclusion of pea in
360 the feed formulation of the laying hens had no effects on production performance or egg
361 quality (Kouvunen et al., 2015), and furthermore, egg yield, egg weight and shell
362 quality was not affected by cotton oil in the feed formulation (Balevi and Coskun,
363 2000). Additionally, when selecting the alternative ingredients, their market prices were
364 checked and found to be similar to those of the original ingredients. Besides, Spain is
365 the second largest European producer of cotton and also peas, exporting to other
366 countries (European Commission - Agricultural and Rural Development, 2017;
367 EUROSTAT, 2017). One third of the Spanish land surface is cultivable; therefore, it is
368 reasonable to expect that these alternative ingredients can be produced *in situ* with
369 lower impacts.

370 According to normalization results, natural land transformation was the most
371 important category, followed by terrestrial ecotoxicity and freshwater ecotoxicity. The
372 pea option (scenario 2) reduced the impact of food on natural land transformation,

373 freshwater ecotoxicity and terrestrial ecotoxicity by 69%, 28% and 30%, respectively.
374 The cottonseed oil alternative (scenario 3) also reduced the impact of food on these
375 three categories, by 22%, 32% and 54%, respectively (Figure 5).

376 Regarding the carbon footprint value, the improvements achieved by employing
377 these variants in fodder formulation would be noticeable too, especially in scenario 3
378 (Figure 6). In fact, the use of cottonseed oil instead of palm oil reduced the impact in the
379 carbon footprint by 22%, giving a carbon footprint of 2.3 kg CO₂ eq per kg of eggs.
380 These attempts to improve the environmental performance of farms through changes in
381 the hen feed formulation were addressed from an environmental point of view.
382 However, economical and nutritional aspects should be taken into account before
383 implementing these modifications in hen fodder formulations. Concerning this matter,
384 De Boer et al. (2014) investigated the replacement of American soybean meal for
385 fattening pigs by other European protein sources. However, only a reduction of 2.5% in
386 the carbon footprint could be achieved without increasing the efficiency of the
387 ingredients production in Europe.

388 Besides, as previously mentioned, the feed conversion ratio is key with respect
389 to the environmental impact of hen feeding. This parameter depends on different
390 factors, such as the kind of housing system utilised. Dekker et al. (2011) reported that
391 feed conversion is higher in loose housing compared with battery cage systems.
392 Therefore, the design of farm facilities is also an important aspect to be borne in mind
393 when attempting to decrease the impact of egg production (without breaching European
394 regulations for animal welfare). Other factors that can contribute to decreasing the feed
395 conversion ratio are changes in feed composition and the genetic selection of the laying
396 hens (Dekker et al. 2011).

397 Another aspect to be considered is the purchase of new chicks for replacement of
398 exhausted laying hens, since it was another cause of high environmental impacts related
399 to egg production. Again, the reformulation of the fodder consumed, in this case by the
400 hens used to breed the new chicks, should be considered. Another possibility might be
401 optimizing the productive life of laying hens from an environmental point of view.
402 Nevertheless, extending the laying cycle is a complex issue, since not only economic
403 aspects, but also the flocks' welfare should be taken into account. Additionally,
404 different factors, including genetics, nutrition and the design of housing systems, should
405 be considered to ensure that adverse effects are avoided (Bain et al., 2016).

406

407 **4. CONCLUSIONS**

408

409 A Spanish egg producing farm was taken as a model of intensive egg production
410 in Europe and a study of the yearly environmental performance of the facility was
411 carried out by LCA and carbon footprint calculation. The global warming potential
412 resulting from the production of 1 kg of shell eggs was 3.5 kgCO₂ eq per kg of eggs, a
413 value in the same order as those reported in other studies. Additionally, natural land
414 transformation, and to a lesser extent, terrestrial ecotoxicity were the most notably
415 affected categories, according to the normalization results. The most important source
416 of environmental impacts in all the categories under assessment was hen feeding
417 (specifically soybean and palm oil cultivation), but also the breeding of young chicks to
418 replace exhausted laying hens. Thus, alternative feed formulations would be an
419 important parameter to take into account in order to lessen the environmental impact.
420 This question should be given serious consideration by the fodder industry and also by
421 the governments that could legislate to limit the amounts of the most harmful

422 components in fodder formulation with respect to environmental impact. An
423 optimisation of the production life of laying hens and actions to decrease the feed
424 conversion ratio could also reduce the environmental impact associated with egg
425 production. These changes should consider not only an environmental perspective, but
426 also productivity and economic aspects.

427

428 **ACKNOWLEDGEMENTS**

429 This study was carried out thanks to funding from the Economy and
430 Employment Office of Principality of Asturias (Spain) through project GRUPIN14-140.
431 “Asturiana de Avicultura, S.L.” egg producing farm (San Cucao, Llanera 33425
432 Asturias) and especially, José Ramón García, is gratefully acknowledged for his kind
433 collaboration in supplying the data employed in this research.

434

435 **REFERENCES**

436

437 Arzoumanidis, I., Salomone, R., Petti, L., Mondello, G., Raggi, A., 2017. Is
438 there a simplified LCA tool suitable for the agri-food industry? An assessment of
439 selected tools. *J. Clean. Prod.* 149, 406-425.

440 Bain, M.M., Nys, Y., Dunn, I.C., 2016. Increasing persistency in lay and
441 stabilising egg quality in longer laying cycles. What are the challenges? *Br. Poult. Sci.*
442 57, 330-338.

443 Baldini, C., Gardoni, D., Guarino, M., 2017. A critical review of the recent
444 evolution of Life Cycle Assessment applied to milk production. *J. Clean. Prod.* 140,
445 421-435.

446 Balevi, T., Coskun, B., 2000. Effects of some dietary oils on performance and
447 fatty acid composition of eggs in layers. *Rev. Med. Vet.* 151, 847-854.

448 Calderón, L.A., Iglesias, L., Laca, A., Herrero, M., Díaz, M., 2010. The utility of
449 Life Cycle Assessment in the ready meal food industry. *Resour. Conserv. Recy.* 54,
450 1196-1207.

451 Calderón, L.A., Iglesias, L., Laca, A., Herrero, M., Díaz, M., 2017.
452 Environmental impact of a traditional cooked dish at four different manufacturing
453 scales: from ready meal industry and catering company to traditional restaurant and
454 homemade. *Int. J. Life Cycle Assess.* (in press).

455 Casolani, N., Pattara, C., Liberatore, L., 2016. Water and Carbon footprint
456 perspective in Italian durum wheat production. *Land Use Policy* 58, 394-402.

457 Cederberg, C., Sonesson, U., Henriksson, M., Sund, V., Davis, J., 2009.
458 Greenhouse gas emissions from Swedish production of meat, milk and eggs 1990 and
459 2005. SIK Report No 793.

460 Cederberg, C., 2014. Environmental impact of meat production. Primary
461 Production/Meat and the Environment. *Encyclopedia of Meat Sciences* (Second
462 Edition). Reference Module in Food Science, 502-507.

463 Cesari, V., Zucali, M., Sandrucci, A., Tamburini, A., Bava, L., Toschi, I., 2017.
464 Environmental impact assessment of an Italian vertically integrated broiler system
465 through a Life Cycle approach. *J. Clean. Prod.* 143, 904-911.

466 Ceylan, N., Ciftci, I., Mizrak, C., Kahraman, Z., Efil, H., 2011. Influence of
467 different dietary oil sources on performance and fatty acid profile of egg yolk in laying
468 hens. *J. Animal Feed Sci.* 20, 71-83.

469 Da Silva, V.P., van der Werf, H.M.G., Soares, S.R., Corson, M.S., 2014.
470 Environmental impacts of French and Brazilian broiler chicken production scenarios:
471 An LCA approach. *J. Environ. Manage.* 133, 222-231.

472 Darnhofer, I., Lamine, C., Strauss, A., Navarrete, M., 2016. The resilience of
473 family farms: Towards a relational approach. *J. Rural Stud.* 44, 111-122.

474 De Boer, H.C, van Krimpen, M.M., Blonk, H., Tyszler, M., 2014. Replacement
475 of soybean meal in compound feed by European protein sources. Effects on carbon
476 footprint. Bron Lelystad: Wageningen UR Livestock Research (Livestock research
477 report 819).

478 Dekker, S.E.M., de Boer, I.J.M., Vermeij, I., Aarnink, A.J.A., Groot Koerkamp,
479 P.W.G., 2011. Ecological and economic evaluation of Dutch egg production systems.
480 *Livest. Sci.* 139, 109-121.

481 Del Prado, A., Mas, K., Pardo, G., Gallejones, P., 2013. Modelling the
482 interactions between C and N farm balances and GHG emissions from confinement
483 dairy farms in northern Spain. *Sci. Total Environ.* 465, 156-165.

484 European Commission – Agriculture and Rural Development, 2017.
485 <https://ec.europa.eu/agriculture/> (accessed 24 November 2017).

486 EUROSTAT, 2017. <http://ec.europa.eu/eurostat/> (accessed 24 November 2017).

487 FAO (Food and Agriculture Organization of the United Nations), 2016.
488 <http://www.fao.org/> (accessed 9 May 2017).

489 Freeman, S.R., Poore, M.H., Middleton, T.F., Ferket, P.R. (2009) Alternative
490 methods for disposal of spent laying hens: Evaluation of the efficacy of grinding,
491 mechanical deboning, and of keratinase in the rendering process. *Bioresour. Technol.*
492 100, 4515-4520.

493 Fréon, P.; Avadí, A., Vinatea Chávez, R.A., Iriarte Ahón, F., 2014. Life cycle
494 assessment of the Peruvian industrial anchoveta fleet: boundary setting in life cycle
495 inventory analyses of complex and plural means of production. *Int. J. Life Cycle*
496 *Assess.* 19, 1068-1086.

497 Ghasempour, A., Ahmadi, E., 2016. Assessment of environment impacts of egg
498 production chain using life cycle assessment. *J. Environ. Manage.* 183, 980-987.

499 González-García, S., Gómez-Fernández, Z., Dias, A.C., Feijoo, G., Moreira,
500 M.T., Arroja, L., 2014. Life Cycle Assessment of broiler chicken production: a
501 Portuguese case study. *J. Clean. Prod.* 74, 125-134.

502 Heinonen, J., Säynäjoki, A., Junnonen, J.M., Pöyry, A., Junnila, S., 2016. Pre-
503 use phase LCA of a multi-story residential building: Can greenhouse gas emissions be
504 used as a more general environmental performance indicator? *Build. Environ.* 95, 116-
505 125.

506 Hospido, A., Moreira, M.T., Feijoo, G., 2003. Simplified life cycle assessment
507 of galician milk production. *Int. Dairy J.* 13, 783-796.

508 Iglesias, L., Laca, A., Herrero, M., Díaz, M., 2012. A life cycle assessment
509 comparison between centralized and decentralized biodiesel production from raw
510 sunflower oil and waste cooking oils. *J. Clean. Prod.* 37, 162-171.

511 Kalhor, T., Rajabipour, A., Akram, A., Sharifi, M., 2016. Environmental impact
512 assessment of chicken meat production using life cycle assessment. *IPA* 3, 262-271.

513 Khatri, P., Jain, S., 2017. Environmental life cycle assessment of edible oils: A
514 review of current knowledge and future research challenges. *J. Clean. Prod.* 152, 63-76.

515 Koivunen, E., Tuunainen, P., Valkonen, E., Rossow, L., Valaja, J., 2014. Use of
516 faba beans (*Vicia faba* L.) in diets of laying hens. *AFSci* 23, 165-172.

517 Koivunen, E., Tuunainen, P., Valkonen, E., Valaja, J., 2015. Use of semi-
518 leafless peas (*Pisum sativum* L) in laying hen diets. *AFSci* 24, 84-91.

519 Leenstra, F., Maurer, V., Galea, F., Bestman, M., Amsler-Kepalaite, Z.,
520 Visscher, J., Vermeij, I., van Krimpen, M., 2014. Laying hen performance in different
521 production systems; why do they differ and how to close the gap? Results of discussions
522 with groups of farmers in The Netherlands, Switzerland and France, benchmarking and
523 model calculations. *Europ. Poult. Sci.* 78, 1-10.

524 Leinonen, I., Williams, A.G., Wiseman, J., Guy, J., Kyriazakis, I., 2012.
525 Predicting the environmental impacts of chicken systems in the United Kingdom
526 through a life cycle assessment: egg production systems. *Poult Sci.* 91, 26-40.

527 MAPAMA (Ministry of Agriculture, Fishing, Food and Environment (Spain)),
528 2017. <http://www.mapama.gob.es/es/> (accessed 17 July 2017).

529 Mattsson, B., Cederberg, C., Blix, L., 2000. Agricultural land use in life cycle
530 assessment (LCA): case studies of three vegetable oil crops. *J. Clean. Prod.* 8, 283-292.

531 MINETAD (Ministry of Energy, Tourism and Digital Agenda (Spain)), 2017.
532 <http://www.minetad.gob.es/es-ES/Paginas/index.aspx> (accessed 17 July 2017).

533 Mollenhorst, H., Berentsen, P., de Boer, I., 2006. On-farm quantification of
534 sustainability indicators: An application to egg production systems. *Br. Poult. Sci.* 47,
535 405-417.

536 Noya I., González-García, S., Berzosa, J., Baucells, F., Feijoo, G., Moreira,
537 M.T., 2017. Environmental and water sustainability of milk production in Northeast
538 Spain. *Sci. Total Environ.* (in press).

539 Nijdam, D., Rood, T., Westhoek, H., 2012. The price of protein: Review of land
540 use and carbon footprints from life cycle assessments of animal food products and their
541 substitutes. *Food Policy* 37, 760-770.

542 Paolotti, L., Boggia, A., Castellini, C., Rocchi, L., Rosati, A., 2016. Combining
543 livestock and tree crops to improve sustainability in agriculture: a case study using the
544 Life Cycle Assessment (LCA) approach. *J. Clean. Prod.* 131, 351-363.

545 Pelletier, N., Ibarburu, M., Xin, H., 2013. A carbon footprint analysis of egg
546 production and processing supply chains in the Midwestern United States. *J. Clean.*
547 *Prod.* 54, 108-114.

548 Pelletier, N., 2017. Life cycle assessment of Canadian egg products, with
549 differentiation by hen housing system type. *J. Clean. Prod.* 152, 167-180.

550 Santos, H.C.M., Jr., Maranduba, H.L., de Almeida Neto, J.A., Rodrigues, L.B.,
551 2017. Life cycle assessment of cheese production process in a small-sized dairy industry
552 in Brazil. *E.S.P.R.* 24, 3470-3482.

553 Saswattecha, K., Kroeze, C., Jawjit, W., Hein, L., 2015. Assessing the
554 environmental impact of palm oil produced in Thailand. *J. Clean. Prod.* 100, 150-169.

555 Skunca, D., Tomasevic, I., Djekic, I., 2015. Environmental Performance of the
556 Poultry Meat Chain - LCA Approach. *Procedia Food Sci.* 5, 258-261.

557 Tecco, N., Baudino, C., Girgenti, V., Peano, C., 2016. Innovation strategies in a
558 fruit growers association impacts assessment by using combined LCA and s-LCA
559 methodologies. *Sci. Total Environ.* 568, 253-262.

560 Vázquez-Rowe, I., Hospido, A., Moreira, M.T., Feijoo, G., 2012. Best practices
561 in life cycle assessment implementation in fisheries. Improving and broadening
562 environmental assessment for seafood production systems. *Trends Food Sci. Technol.*
563 28, 116-131.

564 Vázquez-Rowe, I., Villanueva-Rey, P., Hospido, A., Moreira, M.T., Feijoo, G.,
565 2014. Life cycle assessment of European pilchard (*Sardina pilchardus*) consumption. A
566 case study for Galicia (NW Spain). *Sci. Total Environ.* 475, 48-60.

567 Velarde, A., Fàbrega, E., Blanco-Penedo, I., Dalmau, A., 2015. Animal welfare
568 towards sustainability in pork meat production. *Meat Sci.* 109, 13-17.

569 Wiedemann, S.G., McGahan, E.J., Murphy, C.M., 2017. Resource use and
570 environmental impacts from Australian chicken meat production. *J. Clean. Prod.* 140,
571 675-684.

572 Xu, X., Lan, Y., 2016. A comparative study on carbon footprints between plant-
573 and animal-based foods in China. *J. Clean. Prod.* 112, 2581-2592.

FIGURE CAPTIONS

Figure 1. System boundaries referred to the functional unit expressed per functional unit (FU = 13344000 eggs).


Figure 2. Characterization results obtained using ReCiPe Midpoint.


Figure 3. Normalization results obtained using ReCiPe Endpoint.


Figure 4. Normalization results obtained using Greenhouse Gas Protocol.


Figure 5. Normalization results obtained using ReCiPe Midpoint. Comparison of the most significant categories for three different scenarios: Scenario 1 (real data), Scenario 2 (substituting soybean by pea in hen fodder) and Scenario 3 (substituting palm oil by cottonseed oil in hen fodder).




Figure 6. Normalization results obtained using Greenhouse Gas Protocol. Comparison of the carbon footprint values considering fossil and biogenic carbon for three different scenarios: Scenario 1 (real data), Scenario 2 (substituting soybean by pea in hen fodder) and Scenario 3 (substituting palm oil by cottonseed oil in hen fodder).














Table 1. Nutritional composition of the commercial fodder employed in the facility under study.

Component	% (w/w)
Protein	17.3
Lipids	4.0
Fibre	4.5
Ash	14.5
Lysine	0.85
Methionine	0.40
Calcium	4.1
Sodium	0.16
Phosphorus	0.51

Table 2. Inventory data of the farm, expressed per functional unit (FU 1,334,4000 eggs).

Inputs	
1. New laying hens (units)	55000
2. Water (m ³)	3471
3. Electricity (kWh)	49369
4. Cleaning products (bleach) (t)	0.017
5. Fodder (t)	
a. Maize (50%)	1200
b. Soybean (31%)	744
c. Palm oil (11%)	264
d. Sodium bicarbonate (8%)	192
6. Packaging material (t)	
a. Recycled cardboard	56.70
b. Solid cardboard	30.57
7. Transport	
a. By truck (tkm)	543486.42
b. Diesel (t)	3.0
Outputs	
1. Eggs (units)	13344000
2. Exhausted laying hens for slaughtering (t)	111.3
3. Wastes	
a. Wastewater (to treat) (m ³)	347.1
b. Cardboard (to recycle) (t)	69.6
c. Manure (to be used as fertilizer) (t)	1980
d. Municipal wastes (to landfill) (t)	10.4
e. Dead hens (hazardous waste for incineration) (t)	4.77
4. Emissions to air (t)	
a. CH ₄	4.8
b. N ₂ O-N	0.42
c. NH ₃ -N	1.88