

# Complete Genome Sequences of *Lactobacillus* Phages J-1 and PL-1

Maria Eugenia Dieterle,<sup>a,b</sup> Deborah Jacobs-Sera,<sup>b</sup> Daniel Russell,<sup>b</sup> Graham Hatfull,<sup>b</sup> Mariana Piuri<sup>a</sup>

Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, IQUIBICEN-CONICET, Buenos Aires, Argentina<sup>a</sup>; Department of Biological Sciences and Pittsburgh Bacteriophage Institute, University of Pittsburgh, Pittsburgh, Pennsylvania, USA<sup>b</sup>

***Lactobacillus* phages J-1 and PL-1 were isolated during the 1960s from abnormal fermentations of Yakult. The genomes are almost identical, but PL-1 has a deletion in the genetic switch region and also differs in a gene coding for a putative tail protein.**

Received 24 October 2013 Accepted 7 December 2013 Published 2 January 2014

Citation Dieterle ME, Jacobs-Sera D, Russell D, Hatfull G, Piuri M. 2014. Complete genome sequences of *Lactobacillus* phages J-1 and PL-1. *Genome Announc.* 2(1):e00998-13. doi:10.1128/genomeA.00998-13.

Copyright © 2014 Dieterle et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Mariana Piuri, [mpiuri@qb.fcen.uba.ar](mailto:mpiuri@qb.fcen.uba.ar).

*Lactobacillus* phage J-1 was originally isolated in 1965 from an abnormal fermentation of Yakult, a Japanese beverage fermented from skim milk and *Chlorella* extracts, using the *Lactobacillus casei* Shirota strain as a starter (1). *Lactobacillus* phage PL-1 was isolated 2 years later when a strain resistant to J-1 was used for manufacturing this beverage. J-1 and PL-1 are serologically related (2).

Most of the early work on these phages was done by research groups in Japan, including aspects of phage morphology, phage inactivation, adsorption to cells, DNA injection, and characterization of a lytic enzyme (3–10). Although both phages were isolated in the 1960s and were extensively studied for decades, the genome sequences have not yet been available.

Here, we describe the sequencing and genome annotations of phages J-1 and PL-1. The J-1 virion DNA is 40,931 bp and PL-1 is 38,880 bp in length. The G+C% contents are 44.8 and 44.9, respectively. Both phages have unique ends with 10 bases and single-stranded cohesive 3' extensions (left end, 3'-CGGTCGGCCT).

Using 454 sequencing, phage genomic DNAs were sequenced by the Pittsburgh Bacteriophage Genome Center to a depth of 75-fold coverage for phage J-1 and 45-fold coverage for phage PL-1. Raw reads were assembled using Newbler version 1.1. The assemblies were then quality controlled using Consed version 20. For phage J-1, 14 Sanger reads were required to resolve weak areas in the assembly. No Sanger reads were needed for PL-1. The finished sequences were analyzed and annotated in genome editors, including DNAMaster (<http://cobamide2.bio.pitt.edu>), GBrowse (11), Glimmer (12), GeneMark (13), tRNAscan-SE (14), Aragorn (15), and then were manually curated. Each of the determined open reading frames (ORFs) was functionally annotated using BLASTp (16), CDD (17), and HHpred (18).

An analysis of the J-1 genome reveals 63 potential ORFs, with 57 genes on the positive strand and 6 genes on the negative strand. The PL-1 genome contains 59 putative ORFs; 57 are rightward transcribed, while two are read in the opposite direction.

Interestingly, the PL-1 and J-1 genomes are almost identical. Compared to J-1, PL-1 has a deletion of 1,900 bp comprising 4 genes that belong to the genetic switch region. PL-1 also differs from J-1 in a gene coding for a putative tail protein (gp16). Since

PL-1 was isolated from an abnormal fermentation process using a J-1-resistant strain, gp16 might be involved in host recognition.

The genome organization shares the structure observed in other *Lactobacillus* phages and can be divided into the following modules: packaging, structural proteins, lysis, integration, genetic switch, and replication. An integrase gene (gp24) of the tyrosine recombinase family was found, supporting the temperate origin of these phages.

Regarding the differences found among the J-1 and PL-1 genomes, these data might provide new insights on how phages evolve to counteract bacterial resistance mechanisms.

**Nucleotide sequence accession numbers.** Both genome sequences have been deposited in the GenBank under accession no. [KC171646](https://www.ncbi.nlm.nih.gov/nuccore/KC171646) (J-1) and accession no. [KC171647](https://www.ncbi.nlm.nih.gov/nuccore/KC171647) (PL-1).

## ACKNOWLEDGMENTS

This work was partially supported by ANPCyT grant no. PICT2008-0218 to M.P. M.E.D. is a doctoral fellow of CONICET.

We thank Andrea Quiberoni (Instituto de Lactología Industrial, Argentina) for providing J-1 and PL-1 phages.

## REFERENCES

- Hino M, Ikebe N. 1965. Studies on the lactic acid bacteria employed for beverage production. II. Isolation and some properties of a bacteriophage infected during the fermentation of lactic acid beverage. *J. Agric. Chem. Soc. Jpn.* 39:472–476.
- Watanabe K, Takesue S, Jin-Nai K, Yoshikawa T. 1970. Bacteriophage active against the lactic acid beverage-producing bacterium *Lactobacillus casei*. *Appl. Microbiol.* 20:409–415.
- Kakita Y, Kashige N, Murata K, Kuroiwa A, Funatsu M, Watanabe K. 1995. Inactivation of *Lactobacillus* bacteriophage PL-1 by microwave irradiation. *Microbiol. Immunol.* 39:571–576. <http://dx.doi.org/10.1111/j.1348-0421.1995.tb02244.x>.
- Kashige N, Kakita Y, Nakashima Y, Miake F, Watanabe K. 2001. Mechanism of the photocatalytic inactivation of *Lactobacillus casei* phage PL-1 by Titania thin film. *Curr. Microbiol.* 42:184–189. <http://dx.doi.org/10.1007/s002840010201>.
- Kashige N, Nakashima Y, Miake F, Watanabe K. 2000. Cloning, sequence analysis, and expression of *Lactobacillus casei* phage PL-1 lysis genes. *Arch. Virol.* 145:1521–1534. <http://dx.doi.org/10.1007/s007050070073>.
- Capra ML, Quiberoni A, Reinheimer J. 2006. Phages of *Lactobacillus casei/paracasei*: response to environmental factors and interaction with collection and commercial strains. *J. Appl. Microbiol.* 100:334–342. <http://dx.doi.org/10.1111/j.1365-2672.2005.02767.x>.

7. Nakashima Y, Ikeda H, Kakita Y, Miake F, Watanabe K. 1994. Restriction map of the genomic DNA of *Lactobacillus casei* bacteriophage PL-1 and nucleotide sequence of its cohesive single-stranded ends. *J. Gen. Virol.* 75(Pt 9):2537–2541. <http://dx.doi.org/10.1099/0022-1317-75-9-2537>.
8. Sechaud L, Cluzel PJ, Rousseau M, Baumgartner A, Accolas JP. 1988. Bacteriophages of *Lactobacilli*. *Biochimie* 70:401–410. [http://dx.doi.org/10.1016/0300-9084\(88\)90214-3](http://dx.doi.org/10.1016/0300-9084(88)90214-3).
9. Watanabe K, Kakita Y, Nakashima Y, Sasaki T. 1990. Protoplast transfection of *Lactobacillus casei* by phage PL-1 DNA. *Agric. Biol. Chem.* 54: 937–941. <http://dx.doi.org/10.1271/bbb1961.54.937>.
10. Watanabe K, Ishibashi K, Nakashima Y, Sakurai T. 1984. A phage-resistant mutant of *Lactobacillus casei* which permits phage adsorption but not genome injection. *J. Gen. Virol.* 65:981–986. <http://dx.doi.org/10.1099/0022-1317-65-5-981>.
11. Stein LD, Mungall C, Shu S, Caudy M, Mangone M, Day A, Nickerson E, Stajich JE, Harris TW, Arva A, Lewis S. 2002. The generic genome browser: a building block for a model organism system database. *Genome Res.* 12:1599–1610. <http://dx.doi.org/10.1101/gr.403602>.
12. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with Glimmer. *Nucleic Acids Res.* 27: 4636–4641. <http://dx.doi.org/10.1093/nar/27.23.4636>.
13. Borodovsky M, McIninch J. 1993. Recognition of genes in DNA sequence with ambiguities. *Biosystems* 30:161–171. [http://dx.doi.org/10.1016/0303-2647\(93\)90068-N](http://dx.doi.org/10.1016/0303-2647(93)90068-N).
14. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25: 955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
15. Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res.* 32:11–16. <http://dx.doi.org/10.1093/nar/gkh152>.
16. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic Local Alignment Search Tool. *J. Mol. Biol.* 215:403–410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).
17. Marchler-Bauer A, Bryant SH. 2004. CD-Search: protein domain annotations on the fly. *Nucleic Acids Res.* 32(Web Server issue):W327–W331. <http://dx.doi.org/10.1093/nar/gkh454>.
18. Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein homology detection and structure prediction. *Nucleic Acids Res.* 33:W244–W248. <http://dx.doi.org/10.1093/nar/gki408>.