


Graphical Abstract

Chemical constituents from *Sophora tonkinensis* and their glucose transporter 4 translocation activities

Leave this area blank for abstract info.

Xinzhou Yang, Shihao Deng, Mi Huang, Jialin Wang, Li Chen, Mingrui Xiong, Jie Yang, Sijiang Zheng, Xinhua Ma, Ping Zhao*, Yunjiang Feng*





Chemical constituents from *Sophora tonkinensis* and their glucose transporter 4 translocation activities

Xinzhou Yang^{a,1}, Shihao Deng^{a,1}, Mi Huang^a, Jialin Wang^a, Li Chen^a, Mingrui Xiong^a, Jie Yang^a, Sijiang Zheng^a, Xinhua Ma^a, Ping Zhao^{a,*}, Yunjiang Feng^{b,*}

^aSchool of Pharmaceutical Sciences, South-Central University for Nationalities, Wuhan 430074, China

^bEskitis Institute, School of Natural Sciences, Griffith University, Brisbane, QLD 4111, Australia

ARTICLE INFO

Article history:

Received

Revised

Accepted

Available online

Keywords:

Flavonoids

Polyprenylated flavonoids

Flavone

Antidiabetic

GLUT-4

Traditional Chinese Medicine

ABSTRACT

Bioassay-guided phytochemical investigation of the EtOAc fraction (ST-EtOAc) from the roots of *Sophora tonkinensis* resulted in the isolation of a new compound 1-hydroxy-4-isoprenyl-maackiain (**1**), along with 12 known compounds (**2-13**). The structure of the new compound was established by 1D and 2D NMR, MS data and circular dichroism analysis. Polyprenylated flavonoids **6-9** and **11-13** increased GLUT-4 translocation by the range of 1.35-2.75 folds. Sophoranone (**8**) exerted the strongest activity with 2.75 folds GLUT-4 translocation enhancement at the concentration of 10 μ M. This is the first report of the GLUT-4 translocation activity of the plant *Sophora tonkinensis*.

2009 Elsevier Ltd. All rights reserved.

¹ These authors contributed to this article equally.

* Corresponding author. Tel.: +86 27 67841196; fax: +86 27 67841196 (P.Z.); tel.: +61 7 37356014; fax: +61 7 37356000 (Y.F.).

E-mail: p.zhao@scuec.edu.cn (P. Zhao); y.feng@griffith.edu.au (Y. Feng).

Diabetes Mellitus (DM) is a metabolic disorder with symptoms of high blood sugar and many other complications.¹ 90% patients worldwide have type 2 diabetes (T2DM) which is resulted from a loss of glucose homeostasis or insulin resistance.² The costs to prevent and treat T2DM and its associated complications have become global health and economic burdens.^{3,4} This problem is particularly prevalent in China due to its rapid economic development, improved survival rate from communicable diseases, and the genetic susceptibility.⁵

GLUT-4, one of the 13 sugar transporter proteins, is highly expressed in adipose tissue and skeletal muscle and catalyzes hexose transport across cell membranes.⁶ The protein also mediates glucose removal from the circulation and regulates glucose homeostasis.^{7,8} Given its role in sugar regulation, GLUT-4 has been used as an important target for anti-diabetic drug discovery.

In this study, a cell-based GLUT-4 translocation system was established using L6 recombinant GLUT-4. Confocal imaging technique was used to quantify the translocation activity. The extracts and fractions from 800 traditional Chinese medicines (TCMs) were screened against GLUT4 translocation.⁹⁻¹² One EtOAc fraction (ST-EtOAc) from the roots of *Sophora tonkinensis* exerted a promising stimulatory effect on GLUT-4 translocation. Bioassay-guided phytochemical investigation on the active ST-EtOAc led to the isolation of a new compound, 1-hydroxy-4-isoprenyl-maackiain (**1**), along with 12 known flavonoids.

S. tonkinensis, a traditional medicine used for the treatment of asthma, allergic dermatitis, gastrointestinal hemorrhage, chronic

bronchitis, acute pharyngolaryngeal infections and throat inflammation, was collected from Jingxi County, Guangxi Zhuang Autonomous Region, China, in June, 2012. A specimen (No. SC0060) of this plant has been deposited in the Herbarium of South-Central University for Nationalities, Wuhan, China. The roots of the plant were triturated and then extracted sequentially by maceration with *n*-hexane, EtOAc, methanol successively. The EtOAc fraction was subjected to repeated flash chromatography on silica gel, MCI gel CHP20P and Sephadex LH-20, followed by C₁₈ reverse phase HPLC to afford a new compound, 1-hydroxy-4-isoprenyl-maackiain (**1**, 8 mg), with an isoflavanone conjugate skeleton, as well as 12 known compounds (**2-13**) (Fig 1). The known compounds were identified as trifolirhizin (**2**),¹³ trifolirhizin-6"-monoacetate (**3**),¹³ maackiain (**4**),¹⁴ medicarpin (**5**),¹⁵ 7,4'-dihydroxy-6,8-diprenylflavanone (**6**),¹⁶ glabrol (**7**),¹⁷ sophoranone (**8**),¹⁸ sophoranochromene (**9**),¹⁸ quercetin (**10**),¹⁹ 6,8-diprenylkaempferol (**11**),²⁰ 2-(2',4'-dihydroxyphenyl)-8,8-dimethyl-10-(3-methyl-2-butenyl)-8H-pyrano[2,3-d]chroman-4-one (**12**)²¹ and dehydrolupinifolinol (**13**),²² by the comparison of their spectroscopic data with those previously reported in literature. Herein, we describe the structure elucidation of the new compound and the GLUT-4 translocation activities of compounds **1-13**.

Compound **1** was obtained as an optically active light yellow powder. The molecular formula of C₂₁H₂₀O₆ was established according to the quasi-molecular ion [M - H]⁻ at *m/z* 367.1186 in the HRESIMS, which was in agreement with its ¹³C-NMR data. The UV spectrum showed absorption maximum at 235, 285, 305 nm, typical absorptions of pterocarpanes.¹⁴




Figure 1. Chemical structures of compounds **1-13**.



Table 1. ^1H and ^{13}C NMR data for compound **1** in CD_3OD .

Position	δ_{H} (J in Hz) ^a	δ_{C}	HMBC
1		148.6	
2	6.24, s	100.0	C-1, 3, 4, 4a
3		147.3	
4		108.8	
4a		137.2	
6 α	3.43, d, 10.9 Hz	65.1	C-1, 6a, 6b, 11a
6 β	4.10, dd, 10.9, 4.9 Hz		C-1, 6a, 6b, 11a
6a	3.33, obs	39.6	C-6, 6b, 10a
6b		118.2	
7	6.39, s	92.2	C-6b, 8, 9, 10, 10a
8		141.2	
9		147.1	
10	6.79, s	103.9	C-7, 8, 9, 6a, 6b,
10a		153.8	
11a	5.47, d, 7.0 Hz	77.1	C-4, 6, 6b, 11b
11b		128.2	
OCH ₂ O	5.87, s	100.5	C-8, 9
	5.85, s		
1'	3.65, m	24.5	C-4, 2', 4a, 3
	3.47, m		
2'	5.18, m	122.2	C-4', 5'
3'		130.0	
4'	1.76, s	16.1	C-3', 4', 5'
5'	1.67, s	24.0	C-2', 3', 4'

^a Spectra were recorded for ^1H and ^{13}C NMR data at 30 °C.

The ^1H -NMR spectrum disclosed signals (**Table 1**) for two isolated aromatic protons at δ_{H} 6.79 (1H, s) and 6.39 (1H, s), signals for a -OCHCHCH₂O- moiety [5.47 (1H, d, J = 7.0 Hz), 4.10 (1H, dd, J = 10.9, 4.9 Hz), 3.43 (1H, d, J = 10.9 Hz), 3.33 (1H, obs)], and a dioxygenated methylene at δ_{H} 5.87 (1H, s) and 5.85 (1H, s), suggesting a chemical structure similar to the known compound maackiain.¹⁴ Extra set of signals in **1**, including two methyls at δ_{H} 1.76 (3H, s) and 1.67 (3H, s), an olefin proton at δ_{H} 5.18 (1H, m), and a methylene moiety at δ_{H} 3.65 (1H, m) and δ_{H} 3.47 (1H, m) were assigned to an isoprenyl functionality based on the COSY correlation data from H-1' to H-2', and from H-2' to H-4' and H-5'.


On the basis of above partial structures, the molecular framework was constructed by the HMBC analysis (**Fig. 2**). The characteristic HMBC correlations of the methyl groups at δ_{H} 1.76 and 1.67 to two olefin carbons at δ_{C} 122.2 (C-2') and 130.0 (C-3'), and the olefin proton at δ_{H} 5.18 to the methylene carbon at δ_{C} 24.5 (C-1') confirmed the presence of an isoprenyl group. The HMBC correlations of the olefin proton at δ_{H} 5.18 (H-2') to the aromatic carbon at δ_{C} 108.8 (C-4), and the methylene at δ_{H} 3.65 and 3.47 (H₂-1') to two aromatic carbons at δ_{C} 137.2 (C-4a) and C-4, confirmed that the isoprenyl group was attached at C-4. Moreover, the key HMBC correlations of the isolated aromatic protons at δ_{H} 6.24 to C-4 and C-4a, and two oxygenated aromatic carbons at δ_{C} 148.6 and 147.3 confirmed that the proton at δ_{H} 6.24 located at C-2, and two hydroxyl groups were attached to C-1 and C-3, respectively. Thus, **1** was elucidated as 1-hydroxy-4-isoprenyl-maackiain.²³

**Figure 2.** ^1H - ^1H COSY (←) and key HMBC (→) correlations for **1**.**Figure 3.** The CD spectra of compound **1** (red) and maackiain (blue).

As shown in **Figure 3**, the circular dichroism (CD) spectrum of compound **1** (red line) showed a positive absorption at 308 nm ($\Delta\epsilon$ = + 6.7) and negative absorptions at 235 nm ($\Delta\epsilon$ = - 26.9) and 291 nm ($\Delta\epsilon$ = - 6.9), similar to those of (-)-maackiain (blue line). Based the combined analysis of CD spectra of compound **1** and maackiain. The absolute configuration of **1** was, therefore, determined to be 6a*R*,11a*R*-, the same as that of maackiain.¹⁴

The potential GLUT-4 translocation activity of compounds **1-13** was tested against pIRAP-mOrange cDNAs transfected L6 cells.⁹⁻¹² Insulin (100 nM) was used as the positive control.

Compound **1** showed weak GLUT-4 translocation effect with 0.79-fold enhancement within 30 minutes. Compounds **2-5** and **10** displayed no activity on GLUT-4 translocation (**Fig. 4**). Polyprenylated flavonoids **6-9** and **11-13** exerted moderate to strong activity, increasing GLUT-4 translocation by 1.35-2.75 folds, respectively. Sophoranone (**8**) exerted the strongest activity with 2.75 folds GLUT-4 translocation enhancement (**Fig. 4, 5A**). Dose-response analysis showed that compound **8** increased GLUT4 translocation of L6 cells in a dose-dependent manner at the lower concentrations ($\leq 10 \mu\text{M}$). At higher concentrations ($> 10 \mu\text{M}$), the compound maintained its GLUT-4 translocation activity at around 2.7 folds (**Fig. 5B**). Initial examination of the structure-activity relationship revealed that the active compounds **6-9** and **11-13** all possessed a flavonoid ring system with an aromatic substituent at the C-6 position and an isoprenyl functionality at the C-3 position, suggesting that these structural features may contribute to their GLUT-4 translocation activity.

**Figure 4.** GLUT-4 translocation effects of compounds **1-13**.




Figure 5. (A) Confocal images in L6 cells incubated in the absence (basal) or presence of compound **8** for 30 minutes (L6 cells were infected with pIRAP-mOrange in order to detect externalized IRAP by confocal microscopy). (B) Dose response curve of compounds **8**.

In conclusion, 6a*R*,11a*R*-1-hydroxy-4-prenyl-maackiain (**1**), a new isoprenyl substituted derivative of maackiain was isolated from the roots of *S. tonkinensis*, along with 12 known compounds (**2-13**). Detailed CD spectroscopic data analysis confirmed the absolute stereochemistry of compound **1**. Polyprenylated flavonoids **6-9** and **11-13** increased GLUT-4 translocation by 1.35-2.75 folds, respectively. Polyprenylated flavonoids may be the active principles responsible for GLUT-4 translocation activity of the ST-EtOAc. This is the first report of GLUT-4 translocation activity of the herbal medicine *Sophora tonkinensis*.

Acknowledgments

We acknowledge the financial support from Chinese National Natural Science Foundation grants (No. 81573561 and 31070744), Hubei Provincial Training Programs of Innovation and Entrepreneurship for Undergraduates (SCX15025), and the State Key Laboratory of Drug Research Foundation (SIMM1403KF-07). We thank Dr. Jingquan Yuan from Guangxi Medicinal Herb Garden for sample identification.

Supplementary material

UV, IR, ¹H NMR, ¹³C NMR, COSY, HSQC, HMBC and CD spectra for 6a*R*,11a*R*-1-hydroxy-4-isoprenyl-maackiain (**1**), detailed experimental procedures including general experimental procedure, plant material, extraction, isolation procedure and ¹H NMR for compounds **2-13**, and GLUT-4 translocation assay.

References and notes

1. Zimmet, P.; Alberti, K. G.; Shaw, J. *Nature* **2001**, *414*, 782.

- Whiting, D. R.; Guariguata, L.; Weil, C.; Shaw, J. *Diabetes Res. Clin. Pract.* **2011**, *94*, 311.
- Afroz, A. *Economic Burden of Type 2 Diabetes Mellitus*, LAP Lambert Academic Publishing: Saarbrücken, **2012**.
- Ha, D. T.; Tuan, D. T.; Thu, N. B.; Nhiem, N. X.; Ngoc, T. M.; Yim, N.; Bae, K. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 5556.
- Yang, W.; Lu, J.; Weng, J.; Jia, W.; Ji, L.; Xiao, J.; Shan, Z.; Liu, J.; Tian, H.; Ji, Q.; Zhu, D.; Ge, J.; Lin, L.; Chen, L.; Guo, X.; Zhao, Z.; Li, Q.; Zhou, Z.; Shan, G.; He, J. *N. Engl. J. Med.* **2010**, *362*, 1090.
- Huang, S. H.; Czech, M. P. *Cell Metab.* **2007**, *5*, 237.
- Zhang, H.; Matsuda, H.; Kumahara, A.; Nakamura, S.; Yoshikawa, M. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 4972.
- Bryant, N. J.; Govers, R.; James, D. E. R. *Nature Rev. Mol. Cell Biol.* **2002**, *3*, 267.
- Wang, C.; Yang, J.; Zhao, P.; Zhou, Q.; Mei, Z. N.; Yang, G. Z.; Yang, X. Z.; Feng, Y. J. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 3096.
- Yang, X. Z.; Yang, J.; Xu, C.; Huang, M.; Zhou, Q.; Lv, J. N.; Ma, X. H.; Ke, C. Q.; Ye, Y.; Shu, G. W.; Zhao, P. *J. Ethnopharmacol.* **2015**, *171*, 161.
- Huang, M.; Zhao, P.; Xiong, M. R.; Zhou, Q.; Zheng, S. J.; Ma, X. H.; Xu, C.; Yang, J.; Yang, X. Z.; Zhang, T. C. *J. Ethnopharmacol.* **2016**, *191*, 71.
- Yang, J.; Zhao, P.; Wan, D. R.; Zhou, Q.; Wang, C.; Shu, G. W.; Mei, Z. N.; Yang, X. Z. *Evid. Based Complement. Alternat. Med.* **2014**, *106206*. doi: 10.1155/2014/106206.
- Yang, X. Z.; Yang, J.; Wang, C.; Wan, J. F.; Yuan, J. Q.; Ren, Y. S. *J. Yunnan. Univ. (Natural Sciences)* **2014**, *36*, 267.
- Song, P.; Yang, X. Z.; Yu, J. *Chem. Res. Chin. Univ.* **2010**, *26*, 563.
- Goel, A.; Kumar, A.; Hemberger, Yasmin, A.; Raghuvanshi, A.; Jeet, R.; Tiwari, G.; Knauer, M.; Kureel, J.; Singh, A. K.; Gautam, A.; Trivedi, R.; Singh, D.; Bringmann, G. *Org. Biomol. Chem.* **2012**, *10*, 9583.
- Kazuaki, K.; Katsuo, H.; Kunihiro, S.; Sadakazu, Y. *Chem. Pharm. Bull.* **1973**, *21*, 1777.
- Cho, S.; Park, J. H.; Pae, A. N.; Han, D.; Kim, D.; Cho, N. C.; No, K. T.; Yang, H.; Yoon, M.; Lee, C.; Shimizu, M.; Baek, N. I. *Bioorg. Med. Chem.* **2012**, *20*, 3493.
- Li, X. N.; Lu, Z. Q.; Chen, G. T.; Yan, H. X.; Sha, N.; Guan, S. H.; Yang, M.; Hua, H. M.; Wu, L. J.; Guo, D. A. *Magn. Reson. Chem.* **2008**, *46*, 898.
- Wan, J. F.; Yang, X. Z.; Yuan, J. Q. *J. Yunnan. Univ. (Natural Sciences)* **2011**, *33*, 463.
- Meragelman, K. M.; McKee, T. C.; Boyd, M. R. *J. Nat. Prod.* **2001**, *64*, 546.
- Kyogoku, K.; Hatayama, K.; Yokomori, S. *Chem. Pharm. Bull.* **1973**, *21*, 1192.

22. Sutthivaiyakit, S.; Thongnak, O.; Lhinhatrakool, T.; Yodchun, O.; Srimark, R.; Dowtaisong, P.; Chuankamnerdkarn, M. *J. Nat. Prod.* **2009**, *72*, 1092.

23. 6a*R*,11a*R*-1-hydroxy-4-isoprenyl-maackiain (**1**): yellow powder; UV (CH₃OH) max (log ε) 235 (1.96), 285 (1.02) and 305 (2.08) nm; IR (film) 3432, 2913, 1702, 1616, 1469, 1326, 1126, 1037, 844 cm⁻¹; ¹H and ¹³C NMR data (CD₃OD) see **Table 1**; (-)-HRESIMS m/z 367.1186 [M - H]⁻ (calcd. for C₂₁H₁₉O₆, 367.1186).