

Construction of tilted algebras

Dieter Happel and Claus Michael Ringel

Let A be a finite dimensional algebra. The (finite dimensional) A -module T_A is said to be a tilting module provided it satisfies the following three properties:

(1) There is an exact sequence $0 \rightarrow P''_A \rightarrow P'_A \rightarrow T_A \rightarrow 0$ with P', P'' projective (thus $\text{proj. dim. } (T_A) \leq 1$).

(2) $\text{Ext}^1(T_A, T_A) = 0$.

(3) There is an exact sequence $0 \rightarrow A_A \rightarrow T'_A \rightarrow T''_A \rightarrow 0$ with T', T'' being direct sums of direct summands of T_A (thus, T_A -codim. $(A_A) \leq 1$).

The finite dimensional algebras of the form $\text{End}(T_A)$, where T_A is a tilting module, with A a finite dimensional hereditary algebra, are called the tilted algebras. The main interest in tilted algebras comes from the fact that any finite dimensional algebra with a faithful indecomposable module and with Auslander-Reiten quiver without oriented cycle is a tilted algebra. We will recall some properties of tilted algebras in Section 1, but refer for the proofs to [5].

We will show that any hereditary algebra A has only finitely many tilting modules with basic endomorphism ring of finite representation type (Proposition 2.1). In fact, the proof will provide an inductive procedure for obtaining all these algebras explicitly. The same method also shows that in case A is in addition tame, there are only finitely many basic algebras which are endomorphism rings of tilting modules. Besides those of finite type, one also obtains concealments and domestic regular or coregular enlargements of concealed hereditary algebras (see [6]), the finite type being characterized by the fact that the tilting module has both non-zero preprojective and non-zero preinjective direct summands. The essential properties of tilting modules over tame hereditary algebras without preinjective or preprojective direct summands are collected in Proposition 3.2 and 3.2*. Finally, we want to exhibit some special hereditary algebras A in detail and list corresponding tilting modules, namely we consider hereditary algebras of type A_n, E_6, \tilde{A}_n and \tilde{E}_6 .


1. Report on tilted algebras

Let A be a hereditary algebra. In this case, it is rather easy to check whether a given module is a tilting module. Namely, T_A is a tilting module if and only if $\text{Ext}^1(T_A, T_A) = 0$, and the number of (isomorphism classes of) indecomposable direct summands of T_A is equal to the number of simple A -modules.


Now let T_A be a tilting module. We consider the following two full subcategories: $\mathcal{T}(T_A)$ will denote the full subcategory of all modules generated by T_A . Note that a module X_A is generated by T_A if and only if $\text{Ext}_A^1(T_A, X_A) = 0$. And, $\mathcal{F}(T_A)$ will denote the full subcategory of all modules cogenerated by τT_A , where τ is the Auslander-Reiten translation $\tau = D \text{Tr}$. A module X_A is cogenerated by T_A if and only if $\text{Hom}_A(T_A, X_A) = 0$. Always, the pair $(\mathcal{T}(T_A), \mathcal{F}(T_A))$ forms a torsion theory.

Consider now the tilted algebra $B = \text{End}(T_A)$. In M_B , there is a splitting torsion theory (X, Y) such that X is equivalent, as a category, to $\mathcal{F}(T_A)$, and Y is equivalent to $\mathcal{T}(T_A)$. In fact, the functor $\text{Hom}_A(B T_A, -)$ has as image just Y , and its restrictions to $\mathcal{T}(T_A)$ furnishes an equivalence $\mathcal{T}(T_A) \rightarrow Y$, as Brenner and Butler have shown. Similarly, the functor $\text{Ext}_A^1(B T_A, -)$ has as image just X , and its restriction to $\mathcal{F}(T_A)$ furnishes an equivalence $\mathcal{F}(T_A) \rightarrow X$. Since the torsion theory (X, Y) is splitting, we see that we obtain any indecomposable B -module either in the form $\text{Hom}_A(B T_A, M_A)$ with $M_A \in \mathcal{T}(T_A)$, or in the form $\text{Ext}_A^1(B T_A, M_A)$ with $M_A \in \mathcal{F}(T_A)$. The indecomposable modules of the form $\text{Hom}_A(B T_A, M_A)$ all have projective dimension 0 or 1, those of the form $\text{Ext}_A^1(B T_A, M_A)$ have projective dimension 1 or 2. In particular, the global dimension of a tilted algebra always is ≤ 2 .


As an example, consider the path algebra A of the quiver $0 \rightarrow 0 \rightarrow 0 \leftarrow 0 \leftarrow 0$ of type E_6 . The Auslander-Reiten quiver of A has the following form, the indecomposables being denoted by their dimension type:



we have marked one tilting module $T_A = \bigoplus_{i=1}^6 T_i$. Let us indicate the subcategories $\mathcal{F}(T_A)$ by , and $\mathcal{T}(T_A)$ by :



Now $B = \text{End}(T_A)$ is the algebra given by the quiver $\begin{matrix} 0 & \rightarrow & 0 & \rightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & 0 & \rightarrow & 0 \end{matrix}$ with all commutativity relations, its Auslander-Reiten quiver is as follows (here X is marked by dotted , and Y by hatched):



There is a rather useful criterion in order to decide that an algebra B is a tilted algebra. Namely, assume that the Auslander-Reiten quiver $\Gamma(B)$ of B has a component C which contains no oriented cycles, such that any module in C is of the form $\tau^{-n}P$ for some $n \in \mathbb{N}$ and some projective module P (then C may be called a "preprojective" component; in particular, if B is connected and of finite representation type, then $\Gamma(B)$ itself is a preprojective component if and only if $\Gamma(B)$ has no oriented cycles.) Now, if C contains an indecomposable faithful module N_B , then B is a tilted algebra: there exists a hereditary algebra A , a tilting module T_A , and an indecomposable injective module I_A such that $B = \text{End}(T_A)$, and $N_B = \text{Hom}_A({}_B T_A, I_A)$. In fact, usually there are many different choices possible for A, T_A and I_A , as we will see below.

A very special kind of a tilting module is the slice module of a finite complete slice in a component C of $\Gamma(B)$ which contains all indecomposable projective modules. Recall that a set U of modules in C is said to form a complete slice provided the following three conditions are satisfied

- (i) For any X in C , there exists a unique module of the form $\tau^z X$ with $z \in \mathbb{Z}$ which belongs to U .
- (ii) If $X_0 \rightarrow X_1 \rightarrow \dots \rightarrow X_r$ is a chain of non-zero maps and indecomposable modules, and both X_0, X_r belong to U , then all X_i belong to U .
- (iii) There is no oriented cycle $U_0 \rightarrow U_1 \rightarrow \dots \rightarrow U_r \rightarrow U_0$ of irreducible maps with all U_i in U .

Note that the endomorphism ring of the slice module of a finite complete slice in a component which contains all indecomposable projective modules always is hereditary. In case we start with a complete slice in the preprojective component of a hereditary algebra A , say with slice module U_A , and $B = \text{End}(U_A)$, then we will say that B is obtained from A by an admissible change of orientation. Such a change of orientation can be established also by a sequence of the following operation: Let $P(1), \dots, P(n)$ be the indecomposable projective A -modules, and assume $P(1)$ is simple. Let $U_A = \tau^{-1}P(1) \oplus \bigoplus_{i=2}^n P(i)$. Then U_A is the slice module of a complete preprojective slice, and the quiver of $\text{End}(U_A)$ is obtained from that of A by reversing all arrows involving the point $P(1)$ (Recall that the quiver of a hereditary algebra A is obtained from the full subquiver of $\Gamma(A)$ consisting of the indecomposable projective modules by reversing all arrows). The corresponding functor $\text{Hom}_A(U_A, -) : M_A \longrightarrow M_B$ usually is called a (Bernstein-Gelfand-Ponomarev) reflection functor (see [1]). Also note that in case the quiver of the hereditary algebra A is a tree Δ then any change of orientation of Δ can be realized by a sequence of reflection functors, thus is admissible.

Lemma: Let A be hereditary, T_A a tilting module, and $B = \text{End}(T_A)$. Let I_A be indecomposable injective, and assume $N_B = \text{Hom}_A(B, T_A, I_A)$ is a faithful B -module. Let A' be obtained from A by an admissible change of orientation. Then there exists a tilting module $T_{A'}$, and an indecomposable injective A' -module $I_{A'}$, such that $B = \text{End}(T_{A'})$ and $N_B = \text{Hom}_{A'}(B, T_{A'}, I_{A'})$.

Proof: Since N_B is faithful, we have $\text{Hom}_B(Q_B, N_B) \neq 0$, $\text{Hom}_B(N_B, J_B) \neq 0$ for any indecomposable projective module Q_B and any indecomposable injective module J_B . Consider the complete slice U consisting of all indecomposable modules X_B which have a path of irreducible maps from X_B to N_B such that any such path is sectional. It has been shown in [5] 7.2 that the corresponding slice module U_B is a tilting module, and $A_O = \text{End}(U_B)$ is hereditary with a unique simple injective module I_O such that $N_B = \text{Hom}_{A_O}(B, U_B, I_O)$. Starting from U , we construct inductively various slices, always replacing an indecomposable module X_B which is a source in one slice (and $X_B \neq N_B$), by $\tau^{-1}X_B$. Note that always $\tau^{-1}X_B \neq 0$, due to the fact that N_B is faithful. The corresponding slice modules have as endomorphism rings just hereditary algebras obtained from A_O (and therefore also from A) by all possible admissible changes of orientations.

We note the following consequence: In case the underlying graph of the quiver of A is a tree, then we can construct all possible tilted algebras $\text{End}(T_A)$ which are of finite representation type and have an indecomposable faithful representation, by considering instead of A any fixed algebra obtained from A by a change of orientation.

Finally, we remark that in case B is of finite representation type and $\Gamma(B)$ has no oriented cycles, then an indecomposable module N_B is faithful if and only if it satisfies the following rather weak condition: $\text{Hom}_B(P_B, N_B) = 0$ for a projective module P_B implies $P_B = 0$.

2. Tilted algebras of finite representation type

The following proposition shows that for a fixed hereditary algebra A , there are only finitely many multiplicity-free tilting modules T_A with $\text{End}(T_A)$ of finite representation type. (Recall that a module M_A is called multiplicity-free, if for a decomposition $M_A = \bigoplus M_i$ with M_i indecomposable, these modules M_i are pairwise non-isomorphic, and this happens if and only if $\text{End}(M_A)$ is basic). Thus we can derive from A only a finite number of Morita equivalence classes of algebras of finite representation type as endomorphism rings of tilting modules. In fact, the proof of the proposition provides an inductive procedure in order to obtain all these algebras.

Proposition (2.1) Let A be a hereditary algebra. There are only finitely many multiplicity-free tilting modules T_A with $\text{End}(T_A)$ of finite representation type, and any such tilting module has both indecomposable preprojective and indecomposable preinjective direct summands.

The proof will use the following lemma which is of use also in other situations:

Lemma (2.2) Let A be a connected hereditary algebra of infinite representation type, and T_A a tilting module. Then: T_A has no non-zero preprojective direct summand if and only if all preprojective modules belong to $F(T_A)$. Also, T_A has no non-zero preinjective direct summand if and only if all preinjective modules belong to $\mathcal{T}(T_A)$.

Proof of the lemma. A module M_A belongs to $F(T_A)$ iff $\text{Hom}_A(T_A, M_A) = 0$. Now for M_A preprojective, and $\text{Hom}_A(X_A, M_A) \neq 0$, it follows that X_A has a non-zero preprojective direct summand. Thus, if T_A has no non-zero preprojective direct summand, then all preprojective modules belong to $F(T_A)$. Conversely, if $T(i)$ is an indecomposable preprojective direct summand of T_A , then clearly $\text{Hom}(T_A, T(i)_A) \neq 0$, thus $T(i)$ is not in $F(T_A)$. Similarly, a module M_A belongs to $\mathcal{T}(T_A)$, if and only if $0 = \text{Ext}^1(T_A, M_A) = D \text{Hom}_A(M_A, \tau T_A)$. If M_A is preinjective, and $\text{Hom}(M_A, Y_A) \neq 0$, then Y_A has a non-zero preinjective direct summand. Thus, if T_A has no non-zero preinjective direct summand, the same is true for τT_A , and therefore all preinjective modules are in $\mathcal{T}(T_A)$. Conversely, assume $T(j)$ is an indecomposable preinjective direct summand of T . Since A is connected and of infinite representation type, $T(j)$ is not projective, thus $\tau T(j)$ is non-zero. It is a preinjective module with $\text{Hom}(\tau T(j)_A, \tau T_A) \neq 0$, thus does not belong to $\mathcal{T}(T_A)$.

Proof of the proposition: The second assertion is an immediate consequence of the previous lemma. Let us now prove the first assertion by induction on the number n of simple A -modules. For $n = 1$, there is just one tilting module. Now let A be an algebra with $n > 1$ simple modules, let $P(1), \dots, P(n)$ be the

indecomposable projective A -module.

First, let us construct all tilting modules with a non-zero projective direct summand. For any non-empty subset $I \subseteq \{1, \dots, n\}$, let $P(I) = \bigoplus_{i \in I} P(i)$. Note that there exists an idempotent $e(I)$ of A such that $P(I) \approx Ae(I)$. Let $A(I') = A/\langle e \rangle$, with $\langle e \rangle$ the twosided ideal generated by e . Consider $M_{A(I')}$ as the full subcategory of M_A of all modules X_A satisfying $\text{Hom}_A(P(I), X_A) = 0$. Now $A(I')$ is hereditary, and has a smaller number of simple modules, thus, by induction, there are only finitely many multiplicity-free tilting modules $Y_{A(I')}$ with $\text{End}(Y_{A(I')})$ of finite representation type. Take any such tilting module Y , considered as A -module, which does not contain an indecomposable injective A -module as direct summand, and form $P(I) \oplus \tau_A^{-1}Y_A$. Then clearly this is a tilting module, since it is the direct sum of n pairwise non-isomorphic direct summands and satisfies the Ext-condition. Conversely, let T_A be a multiplicity-free tilting module with $B = \text{End}(T_A)$ of finite representation type, containing at least one indecomposable projective direct summand. A maximal projective direct summand of T_A is of the form $P(I)$ for some non-empty subset $I \subseteq \{1, \dots, n\}$. Let $T_A = P(I)_A \oplus X_A$. Now X has no non-zero projective direct summands, thus $X \approx \tau_A^{-1}\tau_A X$, and $\text{Hom}(P(I), \tau X) = D \text{Ext}(X, P(I)) = 0$, thus $\tau_A X$ is, in fact, an $A(I')$ -module, and, of course, has no indecomposable injective A -module as direct summand. Also, $\text{End}(\tau_A X) \approx \text{End}(X)$ is of the form eBe for some idempotent e , thus also $\text{End}(\tau_A X)$ is of finite representation type.

Let T_A be a multiplicity-free tilting module with $\text{End}(T_A)$ of finite representation type. Then we know that T_A contains both non-zero preprojective and non-zero preinjective direct summands, thus only finitely many of the modules $\tau_A^z T_A$, $z \in \mathbb{Z}$ have n indecomposable direct summands. Since any such class $\{\tau_A^z T_A\}_z$ contains a tilting module with a non-zero projective direct summand, we conclude that altogether there are only finitely many multiplicity-free tilting modules T_A with $\text{End}(T_A)$ of finite representation type.

3. Tilting modules T_A for A tame

Lemma (3.1). Let A be a hereditary tame algebra. Then there are only finitely many basic algebras of the form $B = \text{End}(T_A)$, with T_A a tilting module. Also, if T_A is a tilting module, then T_A has an indecomposable summand which is preprojective or preinjective.

Proof. Note that we may assume A to be connected. Let us first show the second assertion. A tilting module cannot be regular, since the dimension vectors of regular modules lie in a proper subspace of $G_0(A) \otimes Q$, namely in the hyperplane of vectors of zero defect.

However, the dimension vectors of the indecomposable direct summands of a tilting module generate $G_0(A) \otimes Q$. This shows that a tilting module has an indecomposable summand which is preprojective or preinjective.

Assume now that T_A is a multiplicity free tilting module and contains a non-zero preprojective direct summand. Then there is $n \in \mathbb{N}$ such that $\tau^n T$ has a non-zero projective direct summand, but $\tau^i T$ has no non-zero projective direct summand, for all $i < n$. Then $\tau^n T$ also is a tilting module and $\text{End}(T_A) \approx \text{End}(\tau^n T)$. Thus we can assume that $T_A = eA \oplus T'$ for some non-zero idempotent e and a module T' without non-zero projective direct summands. Then $\text{Hom}_A(eA, \tau^{-1} T') = 0$, thus $\tau^{-1} T'$ is an $A/\langle e \rangle$ -module, and in fact an $A/\langle e \rangle$ -tilting module. Since there are only finitely many $A/\langle e \rangle$ -tilting modules, there are only finitely many multiplicity-free tilting modules T_A with a non-zero projective direct summand.

Similarly, there are only finitely many multiplicity-free tilting modules T_A with a non-zero injective direct summand, and this gives all basic algebras which occur as endomorphism rings of tilting modules with a non-zero preinjective direct summand.

If A is a hereditary algebra, we denote its quadratic form on $G_0(A)$ by q .

Proposition (3.2). Let A be a connected tame hereditary algebra. Let T_A be a tilting module and $B = \text{End}(T_A)$. Then the following properties are equivalent:

- (i) $\mathcal{T}(T_A)$ is infinite.
- (ii) $\mathcal{T}(T_A)$ contains infinitely many indecomposable preprojective modules, and the modules $\text{Hom}_A({}_B T_A, X_A)$ with X_A indecomposable preprojective and in $\mathcal{T}(T_A)$ form a component of the Auslander-Reiten quiver of B .
- (iii) There exists a simple homogeneous module which belongs to $\mathcal{T}(T_A)$.
- (iv) All homogeneous modules belong to $\mathcal{T}(T_A)$.
- (v) If R is indecomposable and regular, and $q(\underline{\dim} R) = 0$, then there exists i with $\tau^i R$ in $\mathcal{T}(T_A)$.
- (vi) All preinjective modules belong to $\mathcal{T}(T_A)$.
- (vii) T_A has no non-zero preinjective direct summand.

Proof. The implications (ii) \Rightarrow (i) and (iv) \Rightarrow (iii) are trivial, the equivalence of (vi) and (vii) has been shown in lemma (2.2). Also, the implication (v) \Rightarrow (iv) is obvious, since any homogeneous module H satisfies $q(\underline{\dim} H) = 0$ and $\tau H \approx H$.

(i) \Rightarrow (vii): Assume T_A has an indecomposable preinjective direct summand, say $\tau^m I$ for some $m \in \mathbb{N}$ and some indecomposable injective module I . Let e be an idempotent of A such that $eA/\text{rad}(eA) \approx \text{soc}(I)$. If Y_A is generated by T_A , then $\text{Ext}_A^1(T, Y) = 0$, thus

$$\begin{aligned} 0 &= \text{Ext}_A^1(\tau^m I, Y) \approx D \text{Hom}_A(Y, \tau^{m+1} I) \\ &\approx D \text{Hom}_A(\tau^{-m-1} Y, I) \\ &\approx \text{Hom}_A(eA, \tau^{-m-1} Y), \end{aligned}$$

and therefore either Y is one of the finitely many indecomposable modules with $\tau^{-m-1} Y = 0$, or else $\tau^{-m-1} Y$ is indecomposable and one of the finitely many indecomposable $A/\langle e \rangle$ -modules, and $Y = \tau^{m+1} \tau^{-m-1} Y$. Altogether we see that there are only finitely many possibilities for Y .

(iii) \Rightarrow (vii): Again, assume T_A has an indecomposable preinjective direct summand $\tau^m I$. If S_A is simple homogeneous, and in $\mathcal{T}(T_A)$, then

$$0 = \text{Ext}_A^1(\tau^m I, S) \approx D \text{Hom}(S, \tau^{m+1} I) \approx D \text{Hom}(S, I),$$

since $\tau S \approx S$. However this contradicts the fact that $\underline{\dim} S$ has no zero component.

(vii) \Rightarrow (v): Decompose $T_A = T' \oplus T''$, with T' preprojective and T'' regular. Let R be indecomposable regular, with $q(\underline{\dim} R) = 0$. Let S be simple regular, with an epimorphism $R \twoheadrightarrow S$, and let t be the τ -period of S . Not all $\tau^i S$, $1 \leq i \leq t$, can be regular composition factors of T'' , since $\text{Ext}^1(T'', T'') = 0$. Thus, there is some $\tau^j S$ which is not a regular composition factor of T'' . The top regular composition factor of $\tau^j R$ is $\tau^j S$, thus any non-trivial homomorphism $\tau^j R \rightarrow T''$ would imply that T'' has $\tau^j S$ as regular composition factor. Therefore

$$\begin{aligned} 0 &= \text{Hom}(\tau^j R, T'') \approx \text{Hom}(\tau^{j+1} R, \tau T'') \\ &\approx D \text{Ext}^1(T'', \tau^{j+1} R). \end{aligned}$$

Of course, we also have $\text{Ext}^1(T', \tau^{j+1} R) = 0$, since T' is preprojective and $\tau^{j+1} R$ is regular. Thus $\text{Ext}^1(T, \tau^{j+1} R) = 0$, and therefore $\tau^{j+1} R \in \mathcal{T}(T_A)$.

(iv) \Rightarrow (ii): Let S be a simple homogeneous module. Now, no indecomposable regular module with regular composition factors S can be in $\text{add } T_A$, since $\text{Ext}^1(S, S) \neq 0$. Note that $\text{Hom}(X, S) \neq 0$ for a module X , implies that X has an indecomposable direct summand which is either preprojective or regular with regular composition factors S . Since S is generated by T_A , we conclude that T_A has an indecomposable preprojective direct summand; in particular, there exists an indecomposable preprojective module X in $\mathcal{T}(T_A)$. Now, given X indecomposable preprojective and in $\mathcal{T}(T_A)$, we want to construct an indecomposable preprojective module X'

in $\mathcal{T}(T_A)$ with a non-zero and non-invertible map $X \rightarrow X'$. Note that $\text{Ext}^1(S, X) = D \text{Hom}(X, \tau S) \neq 0$, since $S \approx \tau S$ satisfies $\text{Hom}(P, S) \neq 0$ for all indecomposable projective modules P , and $X \approx \tau^{-n}P$ for some $n \in \mathbb{N}$, and some P . Let

$$(*) \quad 0 \longrightarrow X \xrightarrow{m} E \xrightarrow{P} S \longrightarrow 0$$

be a non-split exact sequence. No direct summand of E can be isomorphic to S , since otherwise the sequence would split. Thus E has to be preprojective. Also, with X and S also $E \in \mathcal{T}(T_A)$. Thus, for X' we may take any indecomposable direct summand of E .

Next, assume X is indecomposable preprojective and in $\mathcal{T}(T_A)$, Y an indecomposable module in $\mathcal{T}(T_A)$, and there is a given a map $f: X \rightarrow Y$ with $\text{Hom}_A({}_B T_A, f)$ irreducible in M_B . We claim that Y also has to be preprojective. First, assume Y is regular. Choose a simple homogeneous module S which is not a regular composition factor of Y , and form a non-split exact sequence $(*)$. The exact sequence induced by f must split, since $\text{Ext}^1(S, Y) = 0$, thus f can be factored as $f = f'm$ through E and neither m is split mono, nor f' is split epi. Since $E \in \mathcal{T}(T_A)$, we obtain a factorization of $\text{Hom}_A({}_B T_A, f)$ with the same properties, thus $\text{Hom}_A({}_B T_A, f)$ cannot be irreducible. Similarly, if Y is preinjective, then any simple homogeneous module S satisfies $\text{Ext}^1(S, Y) = 0$, thus again $\text{Hom}_A({}_B T_A, f)$ cannot be irreducible. It follows now that the set \mathcal{C} of B -modules of the form $\text{Hom}_A({}_B T_A, X_A)$ with X_A indecomposable projective and in $\mathcal{T}(T_A)$ is a complete component of the Auslander-Reiten quiver of B . Namely, assume there is an irreducible map $N_B \rightarrow N'_B$ with N_B in \mathcal{C} , and N'_B indecomposable. Now, either $N'_B = \text{Hom}_A({}_B T_A, Y_A)$ or $N'_B = \text{Ext}_A^1({}_B T_A, Y_A)$ for some indecomposable module Y_A . In the second case, the irreducible map $N_B \rightarrow N'_B$ is involved in a connecting sequence and $N_B \approx \text{Hom}_A({}_B T_A, I_A)$ for some indecomposable injective module, contrary to the assumption $N_B \in \mathcal{C}$. Thus $N'_B = \text{Hom}_A({}_B T_A, Y_A)$, and then, by the previous considerations, Y_A is preprojective.

Proposition (3.2)*. Let A be a connected tame hereditary algebra, and T_A a tilting module with $B = \text{End}(T_A)$. Then the following properties are equivalent:

- (i) $F(T_A)$ is infinite.
- (ii) $F(T_A)$ contains infinitely many indecomposable preinjective modules, and the modules $\text{Ext}_A^1({}_B T_A, X_A)$ with X_A indecomposable preinjective and in $F(T_A)$ form a component of the Auslander-Reiten quiver of B .
- (iii) There exists a simple homogeneous module which belongs to $F(T_A)$.
- (iv) All homogeneous modules belong to $F(T_A)$.
- (v) If R is indecomposable and regular, and $q(\underline{\dim} R) = 0$, then there exists i with $\tau^i R$ in $F(T_A)$.
- (vi) All preprojective modules belong to $F(T_A)$.

(vii) T_A has no non-zero preinjective direct summand.

Proof. The proof is similar to the previous one, again the implications (ii) \Rightarrow (i), (iv) \Rightarrow (iii) and (v) \Rightarrow (iv) are obvious, and the equivalence of (vi) and (v) follows from (2.2).

(i) \Rightarrow (vii), and (iii) \Rightarrow (vii): Let P be indecomposable preprojective, $n \in \mathbb{N}$, and assume $\tau^{-n}P \in \text{add } T_A$. If Y_A is indecomposable and in $F(T_A)$, then $0 = \text{Hom}_A(\tau^{-n}P, Y) \approx \text{Hom}_A(P, \tau^n Y)$, thus there are only a finite number of possibilities for Y . If S_A is simple homogeneous, then $0 \neq \text{Hom}(P, S) \approx \text{Hom}(\tau^{-n}P, S)$, thus $S \notin F(T_A)$.

(vii) \Rightarrow (v): Decompose $T_A = T'' \oplus T'''$ with T'' regular and T''' preinjective. Let R be indecomposable regular, with $q(\underline{\dim} R) = 0$, let S be a simple regular submodule of R , say with τ -period t . Since $\text{Ext}^1(T'', T''') \neq 0$, there exists some i , such that $\tau^i S$ is not a regular composition factor of T'' . Then any homomorphism $\phi: T'' \rightarrow \tau^i R$ is zero, since otherwise the image would contain $\tau^j S$, and $\tau^j S$ would appear as regular composition factor of T'' . Thus $\text{Hom}(T'', \tau^i R) = 0$, and of course, also $\text{Hom}(T''', \tau^i R) = 0$, thus $\tau^i R \in F(T_A)$.

(iv) \Rightarrow (ii): Let S be simple homogeneous. Since $S \in F(T_A)$, we know that S is cogenerated by τT_A , and this immediately implies that τT_A has an indecomposable preinjective direct summand, which clearly has to belong to $F(T_A)$. Now, given any indecomposable preinjective module $X \in F(T_A)$. Then $\text{Ext}^1(X, S) \neq 0$, since S is homogeneous, and X preinjective, thus consider a non-split extension.

$$(*) \quad 0 \rightarrow S \xrightarrow{m} E \xrightarrow{p} X \rightarrow 0.$$

Clearly E has to be preinjective and to belong to $F(T_A)$. In this way, we construct inductively infinitely many indecomposable preinjective modules in $F(T_A)$.

Let X be indecomposable preinjective and in $F(T_A)$, and Y is indecomposable in $F(T_A)$, and assume then is given a map $f: Y \rightarrow X$ such that $\text{Ext}_A^1({}_B T_A, f)$ is irreducible in M_B . Then Y has to be preinjective. Namely, assume Y is regular or preprojective, let S be simple homogeneous which, in case Y is regular, is not a regular composition factor of Y . Then $\text{Ext}^1(Y, S) = 0$, thus the map f can be lifted to E : there exists $f': Y \rightarrow E$ with $f = pf'$. This is a non-trivial factorization of f , and therefore we obtain also a non-trivial factorization of $\text{Ext}_A^1({}_B T_A, f)$, contrary to the assumption that $\text{Ext}_A^1({}_B T_A, f)$ is irreducible. This shows that the set of modules $\text{Ext}_A^1({}_B T_A, X_A)$ with X_A indecomposable preinjective and in $F(T_A)$ is closed under irreducible maps, taking into account that the only indecomposable B -modules which are of the form $\text{Ext}_A^1({}_B T_A, X_A)$ with X_A indecomposable and also are codomains of irreducible maps with domain $\text{Hom}_A({}_B T_A, Y_A)$ are the modules $\text{Ext}_A^1({}_B T_A, P_A)$ with P indecomposable projective, and then $X_A \approx P_A$, thus X_A is not preinjective.

(3.3) We now can distinguish the following possibilities for a tilting module T_A . Always, A denotes a connected tame hereditary algebra, and $B = \text{End}(T_A)$.


- (I) T_A is preprojective or T_A is preinjective.
- (II) $T_A = T' \oplus T''$ with $T' \neq 0$ preprojective and $T'' \neq 0$ regular.
- (II*) $T_A = T'' \oplus T'''$ with $T'' \neq 0$ regular and $T''' \neq 0$ preinjective.
- (III) $T_A = T' \oplus T'' \oplus T'''$ with $T' \neq 0$ preprojective, T'' regular, and $T''' \neq 0$ preinjective.


It is clear that these cases are mutually disjoint. Also, they exhaust all possibilities, since according to 3.1, a tilting module cannot be regular. Let us consider the various cases in more detail.

Case (I). The endomorphism algebras of preprojective modules can also be obtained as endomorphism algebras of preinjective modules, and vice versa. Namely, for any m , the full subcategory of all preprojective modules X satisfying $\tau^m X = 0$ is equivalent to the full subcategory of all preinjective modules Y satisfying $\tau^{-m} Y = 0$ (this follows from the explicit description of these categories given in [2]). Thus, being interested in tilted algebras, we may restrict, in case (I), to the endomorphism algebras of preprojective tilting modules.

Let T_A be a preprojective tilting module. If an indecomposable A -module X is not in $\mathcal{T}(T_A)$, then it is a predecessor of one of the indecomposable direct summands of τT_A . [Namely, X has a non-zero factor module in $F(T_A)$, thus $\text{Hom}_A(X, \tau T_A) \neq 0$.] In particular, the set Z of indecomposable A -modules not in $\mathcal{T}(T_A)$ is finite and contains only preprojective modules. Of course, the indecomposable A -modules in $F(T_A)$ belong to Z .


The Auslander-Reiten quiver of B is obtained from that of A by deleting the finitely many modules from Z , and using the relative Auslander-Reiten sequences from $\mathcal{T}(T_A)$ and joining to the component of preinjective A -modules the indecomposable modules from $F(T_A)$, starting with the connecting sequences and continuing with the relative Auslander-Reiten sequences inside $F(T_A)$.





It is clear that the endomorphism algebras $B = \text{End}(T_A)$ of preprojective tilting modules are "concealed hereditary algebras" in the sense of [6], the equivalent cofinite full subcategories of M_A and M_B being given by $\mathcal{T}(T_A)$ and its image under $\text{Hom}_A({}_B T_A, -)$.

Case (II). This is situation 3.2, thus $\mathcal{T}(T_A)$ contains infinitely many indecomposable preprojective modules, and their images under $F = \text{Hom}_A({}_B T_A, -)$ form a complete connected component of $\Gamma(B)$. Also $\mathcal{T}(T_A)$ contains all preinjective modules, and as usual, their images under F form only part of a component C of $\Gamma(B)$, we have to add the indecomposable modules of the form $\text{Ext}_A^1({}_B T_A, P)$, with P projective, and close under relative Auslander-Reiten sequences from $F(T_A)$. According to 3.2*, the category $F(T_A)$ is finite, thus C will consist precisely of the images of the indecomposable preinjective modules under F and the images of the indecomposable modules in $F(T_A)$ under $\text{Ext}_A^1({}_B T_A, -)$. The remaining components of $\Gamma(B)$ correspond bijectively to the regular components of $\Gamma(A)$. In fact, the homogeneous components of $\Gamma(A)$ are preserved under F , since all homogeneous modules belong to $\mathcal{T}(T_A)$. The remaining regular components of $\Gamma(A)$ may contain also indecomposable modules not in $\mathcal{T}(T_A)$, these have to be deleted - however, according to 3.2(v), $\mathcal{T}(T_A)$ contains quite a few modules from any regular component (of course, this also follows from the fact that all components of $\Gamma(B)$ have to be infinite).



Clearly, the endomorphism algebras $B = \text{End}(T_A)$, with T_A of type (II), are "domestic regular enlargements" (in the sense of [6]) of a tame concealed hereditary algebra C . Here, C is given by the algebra $C = \text{End}(T'_A)$.

Case (II*): This case is dual to the previous case (II), we obtain as endomorphism algebra a "domestic regular coenlargement" of a tame concealed hereditary algebra.

Case (III): According to 3.2 and 3.2*, both $\mathcal{T}(T_A)$ and $\mathcal{F}(T_A)$ are finite categories, thus $B = \text{End}(T_A)$ is of finite representation type.


4. Examples

(4.1) A of type A_n


First, let us consider only tilting modules T_A such that the endomorphism ring $\text{End}(T_A)$ has an indecomposable faithful module.

Lemma. Let A be of type A_n , and T_A a tilting module, and assume that $B = \text{End}(T_A)$ has an indecomposable faithful module. Then T_A is a slice module (thus, B is hereditary of type A_n , again).

Proof. We may consider the Auslander-Reiten quiver $\Gamma(A)$ of A as a subset of $\mathbb{Z}A_n$, the elements of $\mathbb{Z}A_n$ being indexed as follows:



We will use, as in [3], rectangles of the form $q_{(z,r)} = \{(z', r') \mid z+r-n \leq z' \leq z, z+1 \leq z'+r' \leq z+r\}$ for some fixed $(z, r) \in \mathbb{Z}A$, and will call $q_{(z,r)}$ the rectangle ending at (z, r) , or starting at $(z+r-n, n-r+1)$, and also write $(z+r-n, n-r+1)^q$ instead of $q_{(z,r)}$.



The importance of these rectangles lies in the following facts: let X, Y be indecomposable representations of A . Then

(1) $\text{Hom}(X, Y) \neq 0$ iff $\underline{\dim} X \in q_{\underline{\dim} Y}$, and, of course, this is equivalent to $\underline{\dim} Y \in \underline{\dim} X^q$.

(2) $\text{Ext}^1(X, Y) \neq 0$ iff $\underline{\dim} Y \in q_{\underline{\dim} \tau X}$, iff $\underline{\dim} X \in \underline{\dim} \tau^{-1} Y^q$.


Now assume T_A is a tilting module such that $\text{End}(T_A)$ has an indecomposable faithful module. Applying, if necessary, some τ^* , we may assume that T_A has a non-zero projective direct summand, and this implies that any indecomposable faithful B -module, with $B = \text{End}(T_A)$, is of the form $\text{Hom}_A(T_A, X_A)$ for some indecomposable module X_A . Note that the fact that $\text{Hom}_A(T_A, X_A)$ is faithful means that

$\text{Hom}_A(T(i)_A, X_A) \neq 0$ for any indecomposable direct summand $T(i)$ of T_A . Let $x = \underline{\dim} X$, $t(i) = \underline{\dim} T(i)$, thus $t(i) \in q_x$ for all x . In case there exists a sequence of non-zero maps

$$T(i) = T(i_1) \rightarrow T(i_2) \rightarrow \dots \rightarrow T(i_s) = T(j),$$

then clearly $t(j)$ belongs to the set $\{(z', r') \mid z \leq z', z+r \leq z'+r'\}$ where $t(i) = (z, r)$, thus $t(j) \in q_x \cap t(i)^q$.

Now, let $T(i_1), \dots, T(i_u)$ be the sources of $\text{add } T_A$. Then, by the previous considerations, all $t(j)$ belong to $q_x \cap \bigcup_{\alpha=1}^u t(i_\alpha)^q$, and since $\text{Ext}^1(T(j), T(i_\alpha)) = 0$ for all α , we also have $t(j) \notin \bigcup_{\alpha=1}^u \tau^{-1} t(i_\alpha)^q$.



We see that from any τ -orbit of $\mathbb{Z}A_n$, at most one element can be of the form $t(j)$, thus, since there are precisely n different $t(j)$, it follows that $\{T(j) \mid j\}$ is a complete slice.


Let us classify now all tilting modules T_A , for $A = A(n)$ the path algebra of the quiver $1 \xrightarrow{2} 2 \xrightarrow{3} 3 \dots \xrightarrow{n-1} n-1 \xrightarrow{n} n$ (thus A is the algebra of all upper triangular $n \times n$ -matrices, for some n).

Lemma. Let $d = (d_1, \dots, d_n)$ be an integral vector with $d_i > 0$ for all i . Then there exists a tilting module of dimension type d if and only if for any pair $i < k$ with $d_i = d_k$, there exists j with $i < j < k$ and $d_i > d_j$.

Proof. Suppose d does not satisfy this condition. Then it is easy to construct a non-trivial semi-invariant on the affine space of all A -modules of dimension type d , thus there cannot be a tilting module of dimension type d , see [4].

Conversely, assume the condition is satisfied, and let $d_j = \min_{1 \leq i \leq n} d_i$. Let $T(1)$ be the unique indecomposable representation of dimension type $\ell = (1, \dots, 1)$, note that $T(1)$ is both projective and injective, and let $d' = d - d_j \cdot \ell$. The support of d' has precisely $n-1$ elements, and the restriction of d' to its support satisfies the corresponding condition, thus, by induction, there is a module T'_A with $\underline{\dim} T'_A = d'$ and $\text{Ext}^1_A(T'_A, T'_A) = 0$. Also, by induction, the number of isomorphism classes of indecomposable direct summands of T' is precisely $n-1$. It follows that $T_A = (d_j \oplus T(1)) \oplus T'_A$ is a tilting module.


The proof above gives a complete description of all tilting modules. Of course, we may restrict to multiplicity-free ones: Recall that we may associate to any indecomposable module X_A an interval $[\alpha_X, \beta_X]$, namely the support of $\underline{\dim} X$ in $\{1, 2, \dots, n\}$. Let $\alpha_i = \alpha_{T(i)}$, $\beta_i = \beta_{T(i)}$. The interval $[\alpha_i, \beta_i]$ are obtained inductively by deleting in any given interval one point, and considering the remaining points: they form either one or two new intervals: one interval, in case the deleted point was an end point, and two interval otherwise. There is an epimorphism $T(i) \rightarrow T(j)$ if and only if $\alpha_i = \alpha_j$ and $\beta_i \geq \beta_j$, and there is a monomorphism $T(i) \rightarrow T(j)$ if and only if $\alpha_i \geq \alpha_j$ and $\beta_i = \beta_j$. Altogether, we see that the endomorphism ring $B = \text{End}(T_A)$ of T_A is a full connected subquiver of the following "genealogical" tree




with all possible relations $\varphi\psi = 0$. Conversely, given the path algebra of a full connected subquiver of the genealogical tree containing n points, with all relations $\varphi\psi = 0$, there is a unique multiplicity-free tilting module with this algebra as endomorphism algebra.

(4.2) A of type E_6 .

The following list contains all tilted algebras $B = \text{End}(T_A)$, where A is hereditary of type E_6 , and T_A a tilting module, such that there exists an indecomposable faithful B -module. We have written down the quiver with relation of B . As relations, one always has all possible commutativity relations, and some additional zero-relations indicated by a dotted line joining the starting point and the end point of the relation. Non-oriented arrows can be oriented arbitrarily.





(4.3) A of type \tilde{A}_n .

Lemma. Let A be of type \tilde{A}_n , and T_A a preprojective tilting module. Then T_A is a slice module.

Proof. Without loss of generality, we may assume that all simple projective A -modules are direct summands of T_A , applying, if necessary, a suitable number of reflection functors. Let a_1, \dots, a_r be the sinks, and b_1, \dots, b_r the sources of the quiver of A . Since $P(a_j)$ is a direct summand of T_A , for $1 \leq j \leq r$, we have

$$0 = \text{Ext}^1(T(i), P(a_j)) = D \text{Hom}(P(a_j), \tau T(i))$$


for all direct summands $T(i)$ of T_A . The defect $\partial(\tau T(i))$ of $\tau T(i)$ can be calculated as follows:







$$\partial(\tau T(i)) = \sum_{j=1}^r \dim \text{Hom}(P(b_j), \tau T(i)) - \sum_{j=1}^r \dim \text{Hom}(P(a_j), \tau T(i)) \geq 0,$$

thus $\tau T(i)$ can only be preprojective in case $\tau T(i) = 0$. As a consequence, all $T(i)$ are projective, and therefore T_A is the slice module of a complete slice.

(4.4) A of type \tilde{E}_6 .

The following list contains all tilted algebras $B = \text{End}(T_A)$, where A is of type \tilde{E}_6 and T_A is a preprojective tilted algebra (again, given by the quiver of B , with all commutativity relations and additional zero relations).





R e f e r e n c e s .

- [1] Dlab, V., Ringel, C.M.: Indecomposable representations of graphs and algebras. Memoirs Amer. Math. Soc. 173 (1976).
- [2] Dlab, V., Ringel, C.M.: The preprojective algebra of a modulated graph. Proceedings Ottawa Conference 1979. To appear (Springer Lecture Notes).
- [3] Gabriel, P., Riedtmann, Chr.: Group representations without groups. Comment. Math. Helv. 54 (1979), 240-287.
- [4] Happel, D.: Relative invariants and subgeneric orbits of quivers of finite and tame type.I. To appear.
- [5] Happel, D., Ringel, C.M.: Tilted algebras. To appear.
- [6] Ringel, C.M. Tame algebras. Proceedings Ottawa Conference 1979. To appear (Springer Lecture Notes).

D. Happel
C.M. Ringel
Fakultät für Mathematik
Universität Bielefeld
Universitätsstr. 1
D - 4800 Bielefeld 1
West-Germany.