

1 **The structure of the Middle Stone Age of eastern Africa**

2 Blinkhorn, J.^{1,2} & Grove, M.³

3 1 – Department of Geography, Royal Holloway, University of London, Egham, Surrey, UK

4 2 – Department of Archaeology, Max Planck Institute for the Science of Human History, Jena,
5 Germany

6 3 – Department of Archaeology, Classics and Egyptology, University of Liverpool, 12-14
7 Abercromby Square, Liverpool, UK

8 **Abstract:**

9 The Middle Stone Age (MSA) of eastern Africa has a long history of research and is
10 accompanied by a rich fossil record, which, combined with its geographic location, have led it
11 to play an important role in investigating the origins and expansions of *Homo sapiens*. Recent
12 evidence has suggested an earlier appearance of our species, indicating a more mosaic origin
13 of modern humans, highlighting the importance of regional and inter-regional patterning and
14 bringing into question the role that eastern Africa has played. Previous evaluations of the
15 eastern African MSA have identified substantial variability, only a small proportion of which is
16 explained by chronology and geography. Here, we examine the structure of behavioural,
17 temporal, geographic and environmental variability within and between sites across eastern
18 Africa using a quantitative approach. The application of hierarchical clustering identifies
19 enduring patterns of tool use and site location through the MSA as well as phases of significant
20 behavioural diversification and colonisation of new landscapes, particularly notable during
21 Marine Isotope Stage 5. As the quantity and detail of technological studies from individual

22 sites in eastern Africa gathers pace, the structure of the MSA record highlighted here offers a
23 roadmap for comparative studies.

24 **Key Words:** Middle Stone Age; eastern Africa; behavioural variability;

25 **Corresponding Author:** J. Blinkhorn (james.blinkhorn@rhul.ac.uk)

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41 **1. Introduction**

42 Currently, our understanding of the geography of modern human origins is in a state of flux.
43 Until recently, the earliest fossil evidence for *Homo sapiens* was found in eastern Africa dating
44 to ~195 thousand years ago (ka) associated with Middle Stone Age (MSA) technologies
45 (McDougall et al., 2005). Renewed dating of fossil specimens from Jebel Irhoud, North Africa,
46 now present significantly older evidence for the earliest *Homo sapiens* ca 300ka, broadly
47 contemporaneous with the earliest evidence for MSA technologies across Africa (Hublin et al.,
48 2017; Potts et al. 2018). This is supported by genetic studies from southern Africa which
49 indicate the differentiation of modern human populations within the region at a similar time
50 frame (Schlebusch et al., 2017). As a result, eastern Africa no longer presents a discrete source
51 region for the origins of *Homo sapiens*. Nevertheless, due to its pivotal geographic location,
52 eastern Africa remains a potential source region for modern human dispersals out of Africa,
53 offering access to two key routes of expansion into Eurasia via the Bab al Mandeb strait to
54 Arabia or the Nile Valley to the Levant (Groucutt et al., 2015; Lamb et al., 2018). Biological
55 evidence (fossils; genetic studies) increasingly supports a pattern of geographically structured
56 populations amongst early *Homo sapiens* in Africa (Scerri et al. 2018). As a result, eastern Africa
57 may have played a central role mediating interaction between populations split between
58 northern and southern Africa. However, the ability to resolve the nature and configuration of
59 such population structures within Africa is restricted by the sparse fossil record, poor
60 preservation of ancient DNA in the region and limited ability to extrapolate from contemporary
61 populations. Examining the structure of behavioural records offers a complementary approach
62 to understanding the nature of past population structures within Africa (Scerri et al., 2014).
63 Here, we illuminate the structure and variability of MSA stone tool assemblages across eastern
64 Africa using a rigorous quantitative approach, combining data from a newly collated,

65 comprehensive database of stone tool typology and chronology with geographic and
66 environmental datasets.

67 The MSA of eastern Africa, broadly spanning 30-300ka, has a substantial research history. Clark
68 (1988) reviews the earlier history of research and offers an overview of MSA occupations of
69 the region. Critically, he notes that although certain aspects of technology are commonplace,
70 such as Levallois technology or retouched points, they vary across eastern Africa and no
71 feature can be considered ubiquitous. Clark provided an innovative combination of
72 descriptions of sites and stone tool assemblages from across the region with geographic,
73 ecological and environmental maps to illustrate how behavioural diversity was grounded
74 within a diverse physical environment. This perspective on regional variability remains as
75 relevant as ever and has proved robust to the increasing chronological resolution that has
76 since developed.

77 Two more recent syntheses of the eastern African MSA record have played important roles in
78 establishing the nature of behavioural variability in the region. Twenty years after Clark's
79 review, Basell (2008) presented a synthesis of chronometrically dated MSA assemblages
80 accompanied by a qualitative overview of assemblage composition. This overview clearly
81 illustrated the diversity of stone tool use, highlighting considerable overlap between
82 assemblages, and again stressing the absence of any single *fossil directeur* of the eastern
83 African MSA. Extending Clark's focus on the interactions between ecology and behaviour,
84 Basell (2008) highlights the placement of MSA sites within ecotonal settings, permitting access
85 to wooded ecologies, in contrast to the previously assumed central importance of savannahs.
86 Furthermore, the roles of volcanism and tectonics are also recognised alongside patterns of
87 climate change as affecting the habitability of the region and permitting the identification of

88 potential regional refugia. In conclusion, Basell (2008) hypothesised that following contraction
89 of MSA occupations during the high aridity of Marine Isotope Stage (MIS) 6, regional
90 expansion and movement into new environments in MIS 5 corresponded with increased
91 mobility and changes in stone tool use, promoted by climatic, volcanic and tectonic push and
92 pull factors.

93 Tryon and Faith (2013) augment the description of patterns of stone tool technology with the
94 introduction of a quantitative appraisal of the presence/absence of a range of artefact types.
95 Again, descriptions of stone tool technologies broadly echo earlier suggestions for
96 considerable diversity across eastern Africa within different artefact types and the absence of
97 a single unifying type. Notably, these authors suggest that the lack of regionally distinct and
98 derived typological traits is likely to hamper efforts to identify human expansions from the
99 region. Tryon and Faith (2013) evaluated the presence and absence of a range of stone tool
100 and other artefact types using correspondence analysis from dated assemblages,
101 differentiating early (MIS 6 and earlier) from later (MIS 5 and later) assemblages. Alongside
102 considerable overlap between early MSA and some later MSA assemblages, they identify a
103 subset of later MSA assemblages that appear distinct, associated with the presence of blades
104 and backed pieces, as well as beads, grindstones, ochre and anvils. These latter categories
105 appear critical in resolving between earlier and later assemblages (Tryon and Faith 2013: Figure
106 4). In addition, Tryon and Faith (2013) demonstrated a weak but significant negative
107 relationship between geographic distance and assemblage similarity, suggesting that
108 geography does have some effect on the observed patterning. Whether this is due to
109 geographic distance per se, or to habitat differences within the region, remains an open
110 question.

111 These three reviews highlight a number of common themes in their appraisal of eastern African
112 MSA sites, such as the lack of clear intra-regional structure in behaviour, the importance of
113 ecotonal site locations, and the absence of regionally specific stone tool use. Typology remains
114 a key means to evaluate variability across the breadth of the MSA record, although it is not
115 entirely unproblematic. Not only have a wide range of terms been employed to describe stone
116 tools in the MSA of eastern Africa over its extensive research history, but it is broadly
117 acknowledged that significant technological diversity can exist within stone tool types.
118 Elsewhere in Africa, where a comprehensive, technological study of stone tools across a wide
119 area have been conducted within a single analytical system, it has been possible to resolve
120 distinctive, regionalised patterns of behavioural variability within stone tool types (Scerri et al.,
121 2014). To date, no such analysis has been undertaken in eastern Africa.

122 The use of a broad typological approach may somewhat limit the detail of insight into the
123 precise nature of inter-assemblage relationships, but in advance of a fine-grained systematic
124 appraisal of technological approaches, it provides the ability to objectively compare large
125 numbers of sites and to elucidate generic patterns. Any quantitative archaeological analysis
126 faces a trade-off between resolution at the assemblage scale and the number of assemblages
127 that are available for inclusion. The presence / absence approach developed by Tryon and
128 Faith (2013), and extended considerably below, sacrifices fine-scale resolution in favour of
129 analysing the largest possible number of assemblages. This approach is particularly apposite
130 for the eastern African MSA, as it has been established by numerous authors that there exist
131 no 'typical' assemblages that fully characterise this region and period (such as the Aterian of
132 North Africa or Howieson's Poort of South Africa [e.g. Clark 1988; Tryon & Faith 2013]). The
133 goal of this paper is to build upon these previous syntheses of behavioural variability in the
134 MSA of eastern Africa by extending the application of quantitative approaches. In particular,

135 we aim to illuminate the structure of eastern African MSA behaviour, in terms of the typological
136 composition of stone tool assemblages, the variability of site locations with regards to their
137 geographic and environmental features, and how these change through time.

138 **2. Datasets**

139 A broad synthesis of published literature reporting MSA sites was undertaken to compose the
140 dataset for the proposed analyses. Where possible, this involved consulting primary reports
141 on stone tool assemblages, although in rare instances this was not possible. In order to
142 produce as large a database as possible, typological data were also synthesised from
143 secondary sources, principally Basell (2008) and Tryon and Faith (2013), and details of site
144 locations were collected. Chronological data for the assemblages was also collated, but the
145 presence of secure dating was not a prerogative for inclusion within the dataset.

146 Typological terminology used to report MSA assemblages from eastern Africa has varied
147 considerably over the region's extensive research history. In part, this may have stemmed from
148 theoretical differences underlying the methods and goals of stone tool analysis: whether types
149 represent finished tools for either cultural or functional purposes, or whether they occur as
150 points within a reduction continuum. Other factors include the introduction of formal
151 definitions of key technological systems, such as Levallois methods (Boeda, 1994), post-dating
152 the excavation and reporting of key sites. Finally, there is considerable variation in the level of
153 detail available on MSA assemblages, ranging from very basic typologies simply indicating
154 proportions of cores, flakes, tools and debris, to detailed technological descriptions resulting
155 from *chaîne opératoire* studies (e.g. Douze, 2012).

156 The goal here is not to present a new composite typology for studying eastern African MSA
157 assemblages, but to homogenize methodologically diverse reports of stone tool assemblages

158 into a single framework for analysis; this is essential for a thorough examination of the
159 structure of behavioural diversity in the region. Rather than using the typology as an
160 immutable representation of past behaviour, we use it to structure our analysis, and note that
161 tensions within and between typological categorisations may offer profitable lines of future
162 enquiry. Typologies reported frequently conflate reduction methods (e.g. blade production),
163 artefact form (e.g. denticulate), and artefact function (e.g. chopper). This mixture is retained
164 for this analysis for consistency with previous approaches and to enable evaluation of the
165 breadth of MSA behaviour. An important caveat is that the identification and grouping of
166 artefact types is contingent upon the level of resolution afforded by primary reports of stone
167 tool assemblages. It is also important to acknowledge the potential role of raw material
168 variability and flaking mechanics in structuring both past behaviour and the typologies
169 employed by archaeologists studying the MSA of eastern Africa. Again, the ability to evaluate
170 the impact of raw material variability upon stone tool use is constrained by the nature of
171 primary reporting of assemblages and is beyond the scope of the present analysis.

172 An extensive review of the eastern African literature indicates that over 1000 discrete stone
173 tool types have been reported; inevitably, some categories overlap, and many are closely
174 comparable with one another. Preliminary categorisation aimed to standardise these types to
175 the most common terms with minimal data loss, ensuring the use of terms employed by more
176 than 2 separate authors and across more than 2 sites. In order to preserve information this
177 required the splitting of some terms (e.g. "flakes and blades") into more than one category
178 (e.g. "flakes", "blades"). Categories that were nearly ubiquitous (e.g. flakes; core; tool) were
179 then removed as their widespread occurrence offers limited means to resolve patterns of
180 behaviour within the analytical framework adopted. Similarly, some stone tool types (e.g. core

181 management flakes) offer no useable information to resolve between alternate reduction
182 technologies or uses and have been excluded from the analysis.

183 Secondary categorisation aimed to further standardise the use of terminology while limiting
184 the levels of subsets within groups. For example, amongst heavy tools, artefacts may have
185 been reported by a single type (e.g. pick) or include further technological data (e.g. bifacial
186 pick). Similarly, many cores are simply described as Levallois cores, whereas others are
187 described as bidirectional recurrent Levallois cores. In both cases the latter type is a subset of
188 the former and has been subsumed into the more extensive category.

189 In rare instances, single artefacts could not be grouped meaningfully with other types or were
190 best grouped with types that had already been removed (e.g. a single instance of a "basally
191 modified retouched tool") and were also excluded from the data set. Finally, cores, flakes and
192 retouched pieces from particular technological types were combined, as the presence of one
193 (e.g. a retouched Levallois point) suggests the presence of the other two (Levallois point core
194 and Levallois points), and thus serves as an index of the wider technology. A total of twenty-
195 six types were identified for use in the analysis and are described below. These broadly reflect
196 but expand upon other recent, predominantly qualitative, syntheses (Basell 2008, Tryon & Faith
197 2013). Following the categorisation of typological terminology described above, only
198 assemblages that preserved at least two different types were preserved in the data set for the
199 analyses presented below (see Table SI.1 for a full list of sites and references). A total of 125
200 assemblages from 57 sites were identified.

201 **2.1 Reduction technologies**

202 Three types used in the analysis conflate combinations of core and flake types from the same
203 technological systems. *Bipolar technologies* involve striking a core while placed on an anvil.

204 *Core on Flake* technologies exploit flakes as masses of stone for further debitage production,
205 including Kombewa cores and flakes, where flaking is orientated along the original axis of
206 percussion, leading to secondary flaking removing the original bulb of percussion. *Point*
207 *technology* involves the production of triangular flakes, including pseudo-Levallois points. A
208 further four types combine cores, flakes and retouched types from particular technological
209 systems. *Blade technologies* focus upon the production of elongate flakes, typically at least
210 twice as long as they are wide. Following Boeda (1994), Levallois technologies involve
211 hierarchical shaping of core volumes and convexities to predetermine flake shapes. Here, we
212 differentiate *Levallois Flake*, *Blade* and *Point* production. Although not all primary sources
213 clearly differentiate Levallois and non-Levallois blade and point technologies, it is unclear from
214 the literature whether this results from the analytical terms used or represents an actual
215 absence of such artefact types. As all four types do occur in some assemblages, we do not
216 conflate them here, recognising they may illuminate patterns of differentiation of reduction
217 technologies that could be further bolstered by reappraisal of the original assemblages.

218 **2.2 Core Technologies**

219 In addition to those specified above, a number of discrete core reduction methods are
220 included for analysis. Cores that exhibit a distinct platform, but lack other formal preparation
221 are classified as *Platform Cores*, including Single, Multiple and Bidirectional forms. *Discoidal*
222 *Cores* are centripetally flaked from a platform onto a peaked surface and appear as either
223 unifacial or bifacial forms. Here, *Radial Cores* are used to group cores reported either as such
224 or as prepared cores, as both indicate the use of prepared platforms to exploit centripetally
225 flaked, relatively flat core surfaces. In some instances, this may overlap with modern definitions

226 of Levallois flake cores, especially from older reports, although it is worth noting they are still
227 reported as discrete types (e.g. Tryon et al., 2015).

228 **2.3 Retouched Tools**

229 Fourteen different forms of retouched pieces are included for analysis. Some methods of
230 retouching are distinct and form types used in the analysis, such as *Burins*. The use of backing,
231 or abrupt retouch, in the production of diverse *Microliths*, is a further example of a distinct
232 method of retouch, and here different forms of microlithic tools (e.g. crescents, trapezoids) are
233 not differentiated from one another. Bifacially retouched tools are frequently reported in
234 eastern African MSA, and here *RT Bifacial* is used where no further detail regarding tool form
235 is used (e.g. bifacial scraper or point). A number of retouched types are recognised based on
236 tool form. *Retouched (RT) Points* (including unifacial and bifacial forms) have played a key role
237 in the research history of the eastern African MSA, and are widespread, although not
238 ubiquitous. *RT Knives* are often of similar sizes and shapes to retouched points in plan but are
239 only retouched on one edge that is opposite a distinctly thicker edge. *Borers* incorporates all
240 tool forms with a distinct drill-like bit, such as awls. *Scaled Pieces* are typically reported as
241 retouched tools, and although the pattern of crushed retouch they preserve may reflect a
242 distinctive use, it may reflect their production through bipolar reduction. The remaining four
243 retouched types are some of the most common, being *Scrapers*, *Denticulates*, *Notches* and
244 *Notched Tools* (i.e. notched scrapers or notched denticulates). While extensive typologies are
245 reported, especially for scrapers, differences between these tool forms may reflect cultural
246 preferences for particular shapes, their use for alternate tasks, or the accumulation of
247 increasingly invasive retouching through an artefacts life history.

248 **2.4 Heavy Tools**

249 Five types comprise heavy tools represented at MSA eastern Africa sites; these are *Biface*,
250 *Chopper*, *Handaxe*, *Pick* and *Large Cutting Tools (LCT)*. These types have typically been
251 differentiated by their form, with the latter combining tools reported as *LCT* with informal
252 heavy tools as well as tool types which were reported at very low incidence, including core
253 axes and cleavers. In a similar manner to retouched tools, these types may also reflect
254 reduction continua. *Choppers* and *LCTs* may have undergone more limited flaking than *Bifaces*.
255 *Handaxes* typically refer to tear-drop shaped bifaces, whereas *Picks* may reflect one of these
256 forms that has undergone considerably more extensive reduction or use.

257 **2.5 Geography and Environments**

258 Site locations were either collected as co-ordinates from the literature or, where necessary,
259 georeferenced from maps (Table SI.1). Raw data used in the analyses includes SRTM DEM
260 (Jarvis, A., H.I. Reuter, A. Nelson, 2008) data for altitude, and two bioclimatic variables (mean
261 annual temperature, mean annual precipitation) for modern conditions (1970-2000; Hijmans
262 et al., 2005), as well as modelled data for these variables for the Last Glacial Maximum (LGM)
263 21 ka (Braconnot et al., 2007) and the Last Interglacial (LIG) in MIS 5e (Otto-Bliesner et al.,
264 2006). The physiographic landscape of eastern Africa is impacted by ongoing tectonic activity
265 (see Chorowicz, 2005), that complicates the use of the modern landscape to directly
266 characterise those of the past. However, the modern landscape remains the most suitable
267 analogue to past geographic settings, with differences from past conditions partially mitigated
268 by sampling across a 50km radius (see below). Equally, modelled past environmental
269 conditions for the LGM and LIG are used to provide possible extremes of variability observed
270 within a glacial-interglacial cycle and the impact this could have upon human populations,
271 rather than used directly to represent specific conditions during the LIG and LGM.

272 **3. Methods**

273 Dissimilarity matrices and hierarchical clustering are employed to identify patterns of
274 association between both assemblages and typological variables, as well as for spatial datasets,
275 including geographic and environmental data. Dissimilarity (or distance) matrices are
276 produced from pairwise comparisons of cases using an appropriate metric resulting in a
277 measure of dissimilarity. Hierarchical clustering is a form of cluster analysis that iteratively
278 merges cases into a group or splits a group into cases using a measure of closeness, with the
279 results commonly presented in the form of a dendrogram. These are common forms of
280 multivariate analysis, that are regularly described in both introductory and more advanced
281 textbooks (e.g. Krzanowski, 2000; Manly & Alberto 2016). Two alternate metrics are used in
282 calculating the dissimilarity matrices used and are described below. The complete linkage
283 method was employed for hierarchical clustering in all instances, as it is the only method
284 suitable for presence/absence data.

285 **3.1 Behaviour**

286 Jaccard's coefficient is used to calculate dissimilarity between cases of assemblages and lithic
287 types, as it is optimised for analysing presence/absence data. This coefficient treats mutual
288 presence of a particular type in two assemblages as evidence of similarity but gives no weight
289 to mutual absence. The focus on occurrence rather than absence data is particularly suitable
290 given the imperfect nature of archaeological sampling.

291 Two alternate approaches to hierarchical clustering were employed. Divisive clustering was
292 undertaken with the assumption that all cases form part of a cohesive group, i.e. that the MSA
293 reflects a shared behavioural background across the dataset. Each divisive step of the
294 clustering algorithm maximises dissimilarity between cases until each assemblage is separated

295 into an individual cluster. Divisive clustering was undertaken using the DIANA tool in the
296 cluster package of R. In contrast, agglomerative clustering begins with the assumption that all
297 cases are distinct, i.e. that there is no common MSA behavioural background across the
298 dataset. Each agglomerative step minimizes dissimilarity between cases until a single group
299 (i.e. the MSA) is formed. Agglomerative clustering was undertaken using the *hclust* tool in the
300 stats package of R. These alternate approaches to clustering offer complementary information
301 as to how behavioural diversity in the eastern African MSA is structured through time, space
302 and with respect to environmental conditions.

303 **3.2 Geography and Environments**

304 Examining the diversity of site locations focuses upon a number of geographic and
305 environmental parameters, identified above. In addition to altitude, a further geographic
306 dataset representing the energetic consumption for a 50kg human walking within the
307 landscape was created. The SRTM DEM was used to generate a slope raster in ArcGIS 10.3,
308 which was then translated into a raster representing energy consumption in joules for crossing
309 1m at the different slopes encountered, producing an isotropic cost surface based upon results
310 of energy consumption presented by Minnetti and colleagues (2015).

311 In order to understand the landscape in which sites are situated, rather than the individual
312 sites alone, 50km radius buffers around the site were used to sample geographic and
313 environmental datasets, informed by home range sizes from hunter-gather populations
314 (Binford, 2001) and patterns of raw material use (Blegen, 2017; Faith et al., 2015). Individual
315 raster datasets were created for each site buffer for the raster datasets and histogram data
316 were collated using the Zonal Histogram tools in ArcGIS 10.3. Geographic, modern and MIS
317 5e environment data sets resulted in ~9000 cells of data for each site, while LGM data sets

318 resulted in ~3000 cells of data for each site. The exported data were transformed from
319 histogram data to probability distributions. Using the HistDAWass package (Irpino, 2017) in R,
320 dissimilarity matrices were calculated using the L2 Wasserstein distance, which enables
321 characterisation of the scale, skewness and kurtosis of histogram-based data. Hierarchical
322 clustering of the dissimilarity matrices was then employed to examine grouping of sites
323 according to geographic (SRTM DEM and energy consumption cost surface) and three
324 environmental (modern, LGM, LIG) datasets using complete-linkage clustering algorithms.

325 **3.3 Chronology**

326 Chronological data is employed as an additional means to describe patterns of variability
327 amongst behavioural, geographic and environmental datasets, focusing on change through
328 time. Given the wide variety in reporting of ages for MSA assemblages in eastern Africa, and
329 the variability of uncertainty associated with different methods, we confine discussion of
330 chronology in the main text to Marine Isotope Stages. Assemblages are assigned to the MIS
331 with the greatest overlap with reported age constraints from either directly dated assemblages,
332 or those bracketed by overlying and underlying units. Where only minimum or maximum age
333 brackets occur, they are assigned to the stage in which the date occurs. In some instances,
334 undated assemblages have been ascribed to a particular MIS by previous studies based upon
335 site geomorphology (Basell 2008; Tryon & Faith 2013), and these are included here. Marine
336 Isotope Stages are used in the text as a shorthand to describe changing patterns of stone tool
337 typology through time, as well as an index of climatic conditions, rather than as a definitive
338 assessment of eastern African MSA chronology.

339 **4. Results**

340 Hierarchical clustering allows for the clustering of both stone tool types (based on how
341 frequently they are co-present in different assemblages) and assemblages (based on how
342 many tool types they share). Figure 2 shows a binary heatmap of presence and absence data
343 for the 26 stone tool types in each of the 125 MSA assemblages for which data were collated.
344 The results of the analyses described above are presented in three sections. Firstly, the results
345 of clustering of the stone tool types are reported, illuminating which constellations of artefacts
346 are commonly found in association with one another. Secondly, the results of clustering the
347 assemblages are presented, illustrating patterns of similarity and difference in the
348 combinations of artefact types present between assemblages. Thirdly, the results of clustering
349 of geographic and environmental datasets will be presented to examine diversity in the
350 landscapes occupied by eastern African MSA hominins, and correlations with behavioural
351 clusters explored.

352 **4.1 Stone Tool Types**

353 Agglomerative clustering identifies three discrete basal clusters of stone tool types (AT1-3),
354 whereas divisive clustering identifies four discrete basal clusters of stone tool types (DT1-
355 4)(Figure 3). The largest clusters produced by either method (AT1 and DT1) share twelve of
356 fifteen artefact types in common, structured into two (divisive) and three (agglomerative) sub-
357 clusters. Both methods identify *Levallois Flake Technology*, *Blade Technology*, *Platform Cores*,
358 *Discooidal Cores*, *Scrapers* and *RT Points* in one of these sub-clusters, and *Levallois Blade* and
359 *Levallois Point Technology*, *Point Technology*, *Denticulates*, *Cores on Flakes*, and *Choppers* in a
360 second sub-cluster. The third sub-cluster of AT1 is comprised of *Burins*, *Notched Tools* and *RT*
361 *Knife*, which form the discrete basal cluster DT2 using the divisive approach. *Borers*, *Notches*
362 and *LCT's* augment the first subcluster of DT1, whereas using agglomerative methods they

363 form part of AT2. Both agglomerative and divisive methods identify the association of *Bipolar*
364 *Technology, Microliths, Radial Cores, RT Bifacial* and *Scaled Pieces* in discrete basal clusters AT2
365 and DT3. *Bifaces, Handaxes* and *Picks* are identified as discrete basal clusters by both
366 agglomerative (AT3) and divisive (DT4) clustering methods.

367 **4.2 Assemblages**

368 Agglomerative clustering identifies seven discrete basal clusters of stone tool assemblages
369 (A1-A7) while divisive clustering identifies eight separate groups (D1-D8)(Figure 4). No direct
370 overlaps occur in the composition of the clusters identified by alternate methods, although
371 the three members of A2 (Abdur_N_C_S; Garba3_S1; Karungu_GS) are augmented by a fourth
372 site (Omo_KHS_gully) to form D8. Beyond this, numerous pairwise combinations of
373 assemblages are identified by both methods leading to repeated partial overlaps in
374 assemblage clusters identified by alternate methods. At some sites with multiple assemblages,
375 including Koobi Fora, Naisiusiu, Nasera, Olorgesailie, Porc Epic and Prospect Farm, all
376 assemblages form part of the same clusters identified using both clustering methods, although
377 elsewhere, such as at Mumba, Mochena Borago, Lukenya Hill, Koné, alternate assemblages
378 contribute to different clusters. The typological composition of each assemblage cluster is
379 presented in Figure 5 and the most commonplace traits are described in Tables SI.2 and SI.3.

380 *4.2.1 Distribution of assemblage clusters*

381 The largest agglomerative and divisive clusters (A1 and D1) appear at sites that are widely
382 distributed across eastern Africa, and a similar lack of spatial structuring is also apparent
383 amongst the majority of the smaller clusters. Amongst agglomerative clusters, A3 is particularly
384 notable for appearing in a concentration of sites in the Turkana Basin, supplemented by two
385 sites from the northern rift valley and one from the southern rift. Amongst divisive clusters, D5

386 assemblages are only found in the northern rift. Although assemblages from cluster D7 do
387 appear in both the Turkana Basin and northern rift, they are much more numerous in the
388 southern rift.

389 *4.2.2 Chronology*

390 The earliest MSA assemblages date from MIS 9, appearing in two different clusters using both
391 agglomerative (A7 and A1) and divisive (D6 and D7) methods. Amongst agglomerative
392 clusters, A7 is notable for spanning MIS 9-2, with particular concentration of sites apparent
393 from late MIS 5 to early MIS 3. The greatest diversity of agglomerative clusters appears in MIS
394 5, which is the only period when all seven are present, with five agglomerative clusters found
395 in both MIS 7 and MIS 3. Of the seven agglomerative clusters, five appear in both the Middle
396 and Late Pleistocene, with A2 only present during MIS 5, and A6 only apparent in the Late
397 Pleistocene. A1 is notable as it includes the largest number of assemblages during the humid
398 phases of MIS 7, 5 and 3.

399 No single divisive cluster is found in all Marine Isotope Stages, but both D6 and D7 first appear
400 in MIS 9 and are represented in all subsequent stages, apart from MIS 6. While present in MIS
401 8, D1 comprises the largest number of assemblages in MIS 7 and in each stage of the Late
402 Pleistocene. Amongst the smaller clusters, two (D4 and D8) are found in the Middle Pleistocene
403 and MIS 5, but not are not apparent in the latter stages of the Late Pleistocene, whereas two
404 clusters (D2 and D3) first appear in MIS 5 and are only found in the Late Pleistocene.

405 **5. Sites**

406 5.1 Geography

407 Six geographic clusters (G1-G6) are broadly delineated into higher (G1-3) and lower (G4-6)
 408 altitude groups (Figures 7 [left] and SI.2a; Table 1). Abdur (G6) is a unique environment, with
 409 particularly high cost of movement relating to the high slope of the region that spans the
 410 coastal plain to high-altitude hilly terrain. This makes it distinct from other sites located on or
 411 near coastal plains (G5), or those within the low altitude and broadly flat Turkana basin (G4).
 412 The three high altitude clusters exhibit distinct altitude ranges, the lowest of the three, G2,
 413 typically exhibiting a lower cost of movement than either G1 or G3.

Cluster	Description
G1	High Altitude (~1320-1720m)
G2	Middle Altitude (~1075-1320m)
G3	Highest Altitude (~1890-2340m)
G4	Low Altitude (~450-720m)
G5	Coastal (~150-200m)
G6	Near Coastal (~790m)

414 **Table 1:** Key characteristics of geographic clusters.

415 The three largest agglomerative assemblage clusters (A1, A4, A7) appear in the four major
 416 geographic clusters (G1-4) (Figure SI.2). While A7 appear evenly split amongst the middle and
 417 high altitudes of G1-3 and A4 predominately occurs in the high-altitude settings of G1, A1 is
 418 broadly split between the higher altitudes of G1 and G3 and the low altitudes at G4. The
 419 broadest range of agglomerative assemblage clusters is associated with the highest altitude
 420 contexts of G3, with six of seven clusters present. All but one assemblage from coastal contexts
 421 are associated with the smaller agglomerative assemblage clusters, namely A6, A5 and A2.
 422 Seven difference divisive assemblage clusters are found associated with the middle and low
 423 altitude settings of G2 and G4, whereas six divisive assemblage clusters are found associated

424 with the high-altitude settings of both G1 and G3. D7 and D1 appear in the most diverse range
425 of geographic contexts, occurring in all four main geographic clusters, comprising the majority
426 of assemblages associated with the middling altitudes of G2, as well as in one of the coastal
427 groups (G5). D1 is notable for having the highest number of assemblages associated with both
428 the high-altitude contexts of G1 and low altitude contexts of G4.

429 MSA sites in eastern Africa are most consistently found in higher altitude settings of G1,
430 spanning MIS 9-2, and G3, spanning MIS 8-3. While the majority of occupations during MIS 9-
431 8 relate to these upland contexts, the majority of assemblages in MIS 7 occur in the lowland
432 contexts of G4. During MIS 5, occupation appears to be evenly split amongst the five largest
433 clusters and marks the first evidence for occupation of both the coastlines (G5 and G6) as well
434 as low latitude contexts (G2). Occupation in MIS 4 is concentrated in the highest altitudes of
435 G3, whereas in MIS 3 this focus shifts to slightly lower contexts associated with G1.

436 **5.2 Modern Environments**

437 Five clusters are identified based on characteristics of mean annual precipitation and
438 temperature, split between more seasonally variably hot and arid environments (E1 and E2)
439 and less seasonal and more humid environments (E3-5) (Figure 7 [right]). E2 presents one
440 extreme of environments in eastern Africa, comprising arid environments with less than
441 400mm annual rainfall, and mean annual temperatures ranging between 20-30°C. E1
442 comprises sites in semi-arid to sub-humid settings (400-800mm annual rainfall), that are
443 slightly cooler than E2, but with mean annual temperatures mostly above 18°C. Amongst the
444 remaining three clusters, E5 is distinct, presenting the other extreme of eastern African
445 environments with very high levels of humidity of around 1500mm annual precipitation. The

446 remaining two clusters exhibit similar temperature characteristics of 15-25°C, but are split
 447 between semi-humid (800-1000mm) and humid (1000-1200mm) precipitation regimes.

Cluster	Description
E1	450-750mm; > 18°C
E2	<400mm; 20-30°C
E3	950-1185mm; 15-25°C
E4	800-940mm 15-25°C
E5	1500mm;

448 **Table 2:** Mean characteristics of environmental clusters.

449 The distribution of these groups is shown in Figure SI.3. Although many of the clusters appear
 450 widely distributed, elements of spatial structure are evident in the modern environmental
 451 characteristics of eastern African MSA sites. The most northerly sites in the region all fall within
 452 the arid group of clusters E1 and E2, and stand in stark contrast to the more humid habitats of
 453 the central Ethiopian rift, where sites form part of E3 and E4. A similar shift is observed with
 454 sites clustering around Lake Turkana form the major component of the most arid cluster, E2.
 455 A more mixed pattern can be observed in the Kenyan rift, with a greater frequency of sites
 456 from all three humid clusters, as well as closer overlap with semi-arid cluster E1.

457 Environmental clusters E1, E2 and E3 all include members from six of seven of the
 458 agglomerative assemblages. Three agglomerative assemblage clusters appear across the full
 459 suite of modern environmental clusters, with A1 assemblages appearing evenly split across E1-
 460 4, A7 assemblages appearing sparsely in E2 compared to other environments, while A5
 461 assemblages are predominately found in E2.

462 At least one assemblage from each divisive assemblage cluster occurs in E2, with 7 divisive
 463 clusters present in E1, 6 clusters present in E3, with E4 split between five divisive assemblage

464 clusters. Both D6 and D7 appear in all five environmental clusters, with D6 concentrated in the
 465 semi-arid habitats of E1, while D7 is concentrated in more humid environments of E3 and E4.
 466 D1 and D2 assemblages both appear fairly evenly split across the four major environmental
 467 clusters (E1-4).

468 The earliest MSA assemblages in eastern Africa are predominately found in either semi-arid
 469 (E1) or semi-humid environments (E4) during MIS 9 and 8. Indeed, the semi-humid habitats of
 470 E4 have been occupied in each MIS from MIS 9-2, including the only examples of occupation
 471 dating from MIS 6 and 4. Occupation of either wetter (E3) or drier (E2) habitats appears
 472 clustered in more humid stages of MIS 7, 5 and 3. While evidence for occupying the more
 473 humid settings of E3 during the Middle Pleistocene is restricted to a single site, the majority
 474 of MIS 7 occupations are associated with the arid settings of E2. In the Late Pleistocene,
 475 occupation appears fairly evenly distributed between E1, 3 and 4, with fewer sites appearing
 476 in arid E2.

477 **5.4 Past Environments**

478 *5.4.1 Arid conditions*

Cluster	Details
LGM1 – semi humid	12-20°C; 675-880mm
LGM2 – semi arid	13-22°C; 400-650mm
LGM3 - arid	17-26°C; 90-310mm
LGM4 - humid	11-19°C; 1000-1130mm

LGM5 – v humid	15-22°C; 1330-1530mm
----------------	----------------------

479 **Table 3:** Mean characteristics of LGM environment clusters.

480 Using environmental conditions during the LGM as a proxy for drier and cooler phases in
481 eastern Africa, five major clusters are identified, split in to two groups, with the first group of
482 arid to semi-humid settings (LGM1-3) typically more seasonal than the second group of humid
483 habitats (LGM4-5) (Figure 11 [left]). Amongst the first group, cluster 3 is distinct, and is
484 composed of sites with extremely arid and hot environments. Clusters LGM1 and LGM2 share
485 similar ranges of temperature and are split between semi-arid and semi-humid environments.
486 Within the second group, sites in LGM5 exhibit greater humidity and warmer temperatures
487 than LGM4. Although the relationship between more arid and more humid clusters has
488 changed in comparison to modern conditions, limited change is observed within the
489 distribution of the clusters themselves, suggesting a region-wide decrease in humidity rather
490 than reorganisation of environmental conditions associated with more arid phases (Figure
491 SI.4). For instance, little difference is seen in the distribution of the most arid sites, while the
492 Kenyan rift remains a more environmentally diverse region, with sites from different LGM
493 clusters appearing in closer proximity than seen in the Ethiopian rift.

494 Under peak glacial conditions, assemblages from all agglomerative assemblage clusters
495 appear in highly arid and semi-arid settings (LGM2 and LGM3), with A5 assemblages
496 predominantly associated with highly arid LGM3 environments. Semi-humid habitats (E1) are
497 dominated by the presence of the three largest agglomerative assemblage clusters, A1, A4
498 and A7. Both A1 and A5 are notable for appearing in all five LGM clusters.

499 Turning to divisive assemblage clusters, at least one instance of each cluster is found in the
500 arid contexts of LGM3, with seven of eight clusters present in semi-arid LGM2 contexts. The

501 largest assemblage cluster (D1) is predominately found in the semi-arid and semi-humid
502 clusters E1 and E2, a pattern shared with D2. Both D6 and D7 are found in a broad range of
503 environmental settings, spanning LGM1-4, with the former more numerous in LGM2, and the
504 latter occurring more frequently in LGM1, as well as three of five assemblages in LGM5
505 habitats.

506 The modelled environmental parameters for the LGM offer a bracket for past climatic
507 variability during arid phases, but may also offer more suitable analogues for other glacial
508 stages than either present conditions or modelled LIG environments. The earliest MSA sites
509 are found associated with the semi-arid habitats of LGM2, which show some occupation in all
510 MIS except for MIS 4. It is noteworthy that the most arid landscapes (LGM3) are only occupied
511 during interglacial phases (MIS 7,5 and 3). While limited evidence is available for occupation
512 of semi-humid environments (LGM1) in the Middle Pleistocene, this cluster is particularly
513 populous in the Late Pleistocene, including all assemblages dating from MIS 4.

514 *5.4.2 Humid Conditions*

Cluster	Details
LIG1	14-25°C; 370-740mm
LIG2	22-28°C; 145-275mm
LIG3	14-25°C; 765-1005mm
LIG4	14-21°C; 1060-1350mm

515 **Table 4:** Mean characteristics of LIG environment clusters.

516 Using environmental conditions during the last interglacial (MIS 5e ~125ka) as a proxy for hot
517 and humid phases in eastern Africa, four major clusters (LIG1-4) can be observed, principally
518 split on differences of humidity into two pairwise groups (Figure 11 [right]). The first pairwise
519 group comprises LIG1, with a broad range of humidity from semi-arid to sub humid and
520 temperatures ranging between 15-25°C, and LIG2, which are hot (all but one >25°C) and arid.
521 The second pairwise group exhibits a broadly similar range of mean annual temperatures, that
522 also overlaps the range observed for LIG1, but exhibit greater humidity, with LIG4 showing
523 greater humidity than LIG3.

524 Some changes in the distribution of clusters can be observed with respect to modern
525 conditions (Figure SI.5). The most arid cluster (LIG2) is now predominately clustered within the
526 Turkana basin, whereas semi-arid to sub-humid LIG1 is widely dispersed in the Kenyan Rift as
527 well as appearing in the Horn region. Sites appearing in different clusters are again found in
528 closer juxtaposition within the Kenyan Rift, whereas greater homogeneity is observed in the
529 Ethiopian Rift.

530 Assemblages from all agglomerative assemblage clusters appear in the humid LIG3 cluster,
531 with all but one assemblage cluster appearing in the most humid cluster (LIG4). The presence
532 of assemblages in the most arid environments (LIG2) is typically sparse and concentrated in
533 A3. The majority of assemblages in A4, A5 and A6 clusters occur in the semi-arid to sub-humid
534 environments of LIG1, with assemblages from the largest clusters (A1, A7) evenly spread across
535 LIG1, LIG3 and LIG4.

536 Amongst divisive assemblage clusters, seven out of eight clusters are represented in the semi-
537 arid to sub-humid environments of LIG1 and the humid settings of LIG3, with six divisive
538 assemblage clusters apparent in other environments (LIG2, LIG4). Within the largest

539 assemblage cluster (D1), assemblages are relatively evenly spread between alternate
540 environmental clusters, though least populous in LIG2. Clusters D4, D6 and D7 are found in all
541 four LIG environmental clusters, with D6 occurring in greater numbers in LIG1, while the D7
542 appears more concentrated in LIG3.

543 The modelled environmental conditions of the Last Interglacial (MIS 5e) offer an alternate
544 bracket for environmental conditions in eastern Africa to the LGM presented above, relating
545 to humid phases, which may serve as suitable analogues for MIS 9, 7, 5, and 3. There is
546 longstanding evidence for MSA occupation of the semi-arid to sub-humid environments of
547 LIG1, spanning MIS 9 to MIS 2, which includes all known sites in MIS 9 as well as the largest
548 proportion of sites from MIS 3. Occupations of the humid environments of LIG3 also span the
549 Middle and Late Pleistocene, but are heavily concentrated in MIS 5, representing the majority
550 of sites occupied during this stage, with sparse occurrences stretching between MIS 8 and 3.
551 Concentrated occupation of the most arid environments of LIG2 occurs in MIS 7, with two sites
552 occupied in MIS 5 and a single site in MIS 8 in these environments. MSA occupations are not
553 seen in the most humid environments (LIG4) until the Late Pleistocene, appearing in MIS 5 and
554 representing all but one assemblages in MIS 4.

555 **6. Discussion**

556 Through the use of hierarchical clustering, we have set out a detailed, quantitative appraisal of
557 behavioural diversity in the MSA of eastern Africa, both with regards to the constellations of
558 stone tool types used, and the landscape contexts in which they are found. Here, the focus has
559 been to use clusters of stone tools, rather than individual artefact types, to describe patterns
560 of behavioural variability. The use of both agglomerative and divisive approaches to
561 hierarchical clustering of behavioural datasets offer two alternate perspectives on the structure

562 of stone tool use of the MSA in eastern Africa. Divisive clustering starts from an assumption of
563 homogeneity, and subsequently identifies the divisions within the dataset, with the top-level
564 clusters used here marking the minimum number of divisions within the data. As a result,
565 divisive clustering offers an approach to understand the structure of shared technological
566 repertoires across the MSA of eastern Africa. Agglomerative clustering starts from an
567 assumption of heterogeneity, and subsequently identifies bottom-up pairings within the
568 dataset, until the top-level clusters used here are formed. The differential expression of shared
569 technological repertoires in response to distinct geographic and environmental factors may
570 therefore be best approached using agglomerative clustering.

571 **6.1 Stone Tool Types**

572 Both methods of clustering identify two common combinations of MSA behaviour in the
573 largest stone tool type clusters (AT1; DT1), the first (a) including *RT Points, Scrapers, Blade*
574 *technology, Levallois Flake Technology, Platform cores and Discoidal cores*, and the second (b)
575 comprising *Levallois Blade and Levallois Point technologies, Core on Flake technologies, Point*
576 *Technologies, Denticulates and Choppers*. Under divisive clustering, these groups form a single,
577 large cluster which offers a concise identification of the most common, co-occurring features
578 of MSA assemblages. When the frequency of their occurrence is taken into account (see
579 heatmap), these results potentially indicate that (a) comprises the most widespread
580 manifestation of MSA technology, whereas (b) presents the most common means to augment
581 or diversify this technology.

582 Under both clustering methods, *Bipolar Tech and Microliths* form pairwise clusters, as do *Radial*
583 *Cores and RT Bifacials*, and these four types group with *Scaled Pieces* to form a common cluster.
584 While the common co-occurrence of *Bipolar Technology* and *Microliths* is widely noted, their

585 association with alternate flaking and retouching methods that is identified here has not been
586 similarly stressed in previous analyses.

587 Both approaches to clustering indicate that well defined heavy tool types - *Picks, Bifaces* and
588 *Handaxes* - form a distinct group. This common grouping may be explained either as reflecting
589 positions on a shared reduction continuum, or as relating to specific functional demands for
590 diverse, specialised heavy tools. In contrast, the poorly defined heavy tool types - *LCT* and
591 *Choppers* - appear embedded within more diverse stone tool clusters. The requirement for
592 heavy tools within MSA assemblages appears widespread, but the need for more formalised
593 tools may relate to distinct behaviours in response to specific functional demands. At an
594 alternate end to the spectrum of stone tool types, smaller retouched tools types are distributed
595 across these clusters rather than grouping together. If these are interpreted as positions along
596 a reduction continuum, then it appears retouched tools with differing use life histories are
597 associated with differing constellations of other lithic technologies, which could be a result of
598 practical factors, such as distance from raw material resources, or functional or stylistic
599 constraints.

600 The most noticeable difference between clustering approaches relates to the associations of
601 *Burins, Notched Tools* and *RT Knives* as a discrete subgroup on one hand, and *Borers, Notches*
602 and *LCT's* as a distinct subgroup on the other. *Burins, Notched Tools* and *RT Knives* form a
603 discrete basal cluster under a divisive approach, suggesting they represent distinctive
604 strategies amongst MSA technologies, whereas their association with AT1 under an
605 agglomerative approach indicates that they are frequently deployed alongside the most
606 commonplace MSA stone tool types. Likewise, the inclusion of *Borers, Notches* and *LCT's* in
607 DT1 suggests they are a consistent feature of the most common and widespread manifestation

608 of MSA stone tool technologies, but their separation from AT1 may indicate that in practice
609 they are less commonplace or occur in distinct behavioural contexts.

610 **6.2 Assemblages**

611 Seven major clusters were identified by agglomerative clustering. This bottom-up approach to
612 identifying common patterns of behaviour identifies one cluster (A1) which comprises ~37%
613 of all assemblages, suggesting widespread, common practices in using different stone tool
614 technologies. Although no single tool type is shared amongst all assemblages in the cluster,
615 *Levallois Flake Technology*, *Scrapers* and *RT Points* are all abundant. Similarly, the second
616 largest cluster (A7) has no single ubiquitous stone tool types amongst all members, but
617 *Levallois Flake* and *Levallois Blade Technologies* occur in high frequencies. Amongst the smaller
618 clusters, *Handaxes* (A2), *Scrapers* (A3), *Platform Cores* (A4), *RT Points* (A5) and *Microliths* (A6)
619 are found amongst all members of their respective clusters. Notably, even the smallest
620 agglomerative clusters are comprised of tool types spread across the agglomerative tool type
621 clusters, rather than being restricted to a single group.

622 Under divisive clustering the smallest number of clusters that MSA assemblages can be divided
623 into is eight. Amongst these, four large clusters comprise the majority of assemblages (D1,
624 D2, D6 and D7), and offer a characterisation of the most common modalities of MSA
625 behaviour, whereas the four smaller clusters (n<10 assemblages) highlight more uncommon
626 combinations of stone tool technologies. No single tool type appears in all assemblages in the
627 largest cluster (D1), although *Scrapers*, *Levallois Flake Technologies* and *RT Points* appear in
628 over 90% of these assemblages. Members of D2 and D6 all include scrapers, with distinct
629 patterns of both other stone tool types present and their frequency. *Levallois Flake Technology*
630 is ubiquitous in D7, but other tool types are present in low frequencies. Amongst the smaller

631 clusters, D4 is notable for having three tools types, *Levallois Flake Technology*, *Blade*
632 *Technology* and *Scrapers* in all assemblages, whereas other clusters are unified by the presence
633 of *Microliths* (D3), *Bifaces* (D5) and *Handaxes* (D8). In a similar manner to the agglomerative
634 clusters, divisive assemblage clusters comprise tool types that are spread across the divisive
635 tool clusters.

636 Subtle variation in the constellation of artefact occurrence that define the different assemblage
637 clusters and their members occurs between the two methods that precludes easy, direct
638 comparisons. As above, the clusters identified through divisive methods perhaps best
639 characterise the broad groups of technological practice, whereas agglomerative methods
640 identify groups with distinct expressions of varied technological practices. A number of
641 assemblages consistently cluster together under both methods, with 28 assemblages shared
642 by the largest agglomerative (A1) and divisive (D1) assemblage clusters. In a similar manner, 8
643 of 10 assemblages in A3 are found together in D2, while 12 of 21 assemblages in A4 occur
644 together in D1, and three quarters of assemblages in D4 occur together in A7. While multiple
645 assemblages from single sites often cluster together under one method, a more limited
646 number of sites have multiple assemblages that cluster together consistently under both
647 methods, namely Koimilot (Kaphurin Formation), Koobi Fora, Naisiusiu, Nasera, Ologesailie,
648 Porc Epic and Prospect Farm.

649 The methods used here do not neatly identify individual stone tool types, or even
650 combinations of tool types, that are ubiquitous across all assemblages, supporting previous
651 assessments that the MSA is a polythetic group, lacking hard and fast rules to ascribe group
652 membership. Instead, the results demonstrate that stone tool assemblages typically contain
653 artefacts spread across clusters of commonly co-occurring stone tool types, but rarely express

654 all the same elements. Similarly, the use of clustering has enabled quantitative assessment,
655 from a wider range of sites and more numerous stone tool variables, of qualitative appraisals
656 of patterns of spatial variability, suggesting little intra-regional variability.

657 **6.3 Sites**

658 Clustering using geographic as well as modern and modelled past climate data sets readily
659 identify structure both within these data, and with respect to their distribution. Amongst the
660 variables analysed, gross differences in altitude and precipitation present the main differences
661 between clusters formed, with energy expenditure and annual temperature overlapping
662 considerably between groups and presenting more subtle modulation to cluster formation.
663 Notably, both precipitation and altitude have tangible impacts upon the make-up, distribution
664 and seasonality of both faunal and floral populations (Siepielski et al 2017; Rosenzweig 1995).

665 Across datasets, the Kenyan rift presents the most heterogeneous habitats for MSA sites. Sites
666 within the region fall in all three of the higher altitude clusters identified by analysing
667 geographic variables, yet for environmental datasets, sites in the Kenyan rift occur across major
668 divisions of clusters, such that even with the expansion of arid conditions, modelled for the
669 LGM, a balance of arid to humid habitats are found. In contrast, broad latitudinal banding of
670 habitats can be noted in site contexts from the Turkana basin and through the Ethiopian rift.
671 The Turkana basin, presenting low-altitude contexts which consistently present the hottest,
672 most arid site environments, presents a clear break in the landscape structure of eastern Africa,
673 separating the mountainous and humid Kenyan and Ethiopian Rift. To the north, high altitude
674 site contexts stretch through the Ethiopian Rift and the Horn, separating the northernmost
675 lower altitude sites from the Turkana basin. Similarly, the sites in the middle Ethiopian Rift
676 experience higher levels of humidity either than the Turkana basin sites to the south, or the

677 Horn and lowland sites to the north. As well as their distinct topographic setting on the Kenyan
678 coast, Panga ya Saidi and Mtongwe repeatedly fall within more humid environmental clusters,
679 which under some conditions diverge from sites in the Kenyan Rift at a comparable latitude.
680 Similarly, the environmental characteristics of Nyara River, the southernmost site under
681 consideration here, frequently stand in contrast to the rest of eastern Africa. These three sites
682 all fall within more heavily forested regions today, and persistent humidity under past
683 conditions may have perpetuated this distinction from the majority of eastern African MSA site
684 habitats.

685 Only limited evidence for distinct spatial structure amongst stone tool assemblages in MSA
686 eastern Africa is apparent. This may well reflect the influence of the mosaic geographic and
687 environmental make-up of the region, with both considerable distances occurring between
688 areas presenting similar habitats, such as in higher altitude areas of the Kenyan and Ethiopian
689 Rift, as well as distinctly different landscapes occurring in relative proximity in the former.
690 Similarly, no clear-cut patterns of certain assemblage clusters associating solely with single
691 geographic or environmental site clusters were identified. Rather, patterns of emphasis,
692 instead of exclusivity, are observed in associations between the two. Although certain
693 packages of stone tools may have been most frequently used in some environments, the
694 results here support scenarios in which they were not bound to those environments.

695 **6.4 Chronology**

696 Larger numbers of assemblages appear in interglacial stages (MIS 9,7,5 and 3), with warmer
697 and more humid conditions potentially favouring both population growth and geographical
698 expansion across the eastern African landscape. Whilst there may be a taphonomic component
699 to this pattern, it is not simply a chronological preservation bias: for example, fewer sites are

700 documented for the later MIS 4 than for the earlier MIS 5. Furthermore, the increase in numbers
701 of sites in humid periods is accompanied by correlated increases in both the variety of tool
702 forms present and the range of environments inhabited. Figure 9 demonstrates this
703 geographical expansion, whilst Figures 10, 12, and 13 show that a wider range of environments
704 is inhabited during MIS 5 and MIS 3 in particular, regardless of which environmental dataset
705 is employed for reconstruction. MIS 5 is shown to be the most varied stage in terms of
706 technology, geography, and environments inhabited; indeed, this is the only stage in which all
707 of the 26 tool types are present. Sites during MIS 5 also demonstrate an increased habitation
708 of coastal areas and a spread into mid-altitude areas. Collectively, these results support earlier
709 suggestions that during humid stages hominin populations expanded into a broader variety
710 of geographical regions, encountered a broader array of environments, and produced lithic
711 assemblages of more diverse composition (e.g. Basell 2008).

712 Focusing on more archaeologically abundant phases, MIS 7, 5, and 3, there are a limited
713 number of important chronological patterns in the tool forms present. Of the 16 types that are
714 recorded in *all* of these stages, some steadily increase through time (i.e. the proportion of
715 assemblages in which they are present follows a pattern showing MIS7<MIS5<MIS3). These
716 types include *Blade Technology*, *Radial Cores*, *Notched Tools*, and *Bipolar Technology*. Types
717 that decline include an associated group of *Levallois Flake* and *Levallois Blade Technology*, and
718 *Discoidal Cores*, as well as *LCTs*, *Point Technology*, *Denticulates*, and *RT Knives*. Some of these
719 types are closely associated in the dendrograms of Figure 3, whilst others are more distantly
720 related. It remains a likely, therefore, that the gradual replacement of older tool forms by newer
721 technologies is due to a complex interaction of reduction method and function, as well as the
722 need to deal with changes in the resources encountered in an increasing variety of habitats.

723 Investigating the appearance of the largest assemblage clusters with respect to geographic
724 and modern environmental clusters through time highlights a number of familiar patterns. In
725 particular these are (1) continuity of using particular clusters of behaviour through time within
726 the same geographic and environmental contexts, and (2) expanding the use of existing
727 behavioural clusters in new contexts under favourable climates. Rarely do assemblage groups
728 appear in very different settings (e.g. both extreme arid and humid) outside of interglacial
729 phases, but overall these are patterns of emphasis rather than exclusivity. Refining our
730 knowledge of spatial, environmental, and behavioural distinctions through time will remain
731 vitally important to establishing how strong these patterns are, and where informative
732 exceptions to them occur.

733 **7. Conclusions**

734 The Middle Stone Age of eastern Africa exhibits a diverse range of behaviour, in terms of the
735 stone tools used by past populations and the geographic and environmental contexts which
736 they inhabited. Through the application of a quantitative approach, we have been able to
737 explore behavioural variability in greater detail than ever before and set out how this diversity
738 is structured in time, space and across environments. Hierarchical clustering is an ideal method
739 to employ to examine complex patterns of presence and absence across multiple, though at
740 times sparse, typological variables. Importantly, adopting this approach has enabled us to
741 effectively integrate data from sites that have been overlooked from previous, qualitative
742 assessments, and especially those that lack clear chronometric control. The synthesis we
743 present is also unique in the means of integrating data regarding the geographic and
744 environmental contexts of the sites, and particularly for evaluating the variability of the wider

745 landscapes surrounding sites. Combining these complementary approaches has enabled us to
746 identify new patterns in the structure of behavioural variability of the eastern African MSA.

747 The typology employed here is not presented as an authoritative description of MSA stone
748 tool technology but is used as a means to evaluate patterns of variability. It is noteworthy that
749 both top-down (divisive) and bottom-up (agglomerative) methods of identifying clusters of
750 co-occurring stone tool types identify one major cluster with two common, distinct
751 components, from a range of common pair-wise associations. This offers some insight into
752 what may count as a 'classic' MSA repertoire, with the introduction of additional elements
753 resulting from different interactions with the environment as well as patterns of cultural
754 transmission through time and across space. These findings may be consolidated upon further
755 by refining the typology employed; such research will necessitate renewed analyses of existing
756 assemblages where possible, to differentiate, for example, preferential from recurrent Levallois
757 technologies.

758 Engaging with the breadth of MSA data that is available exhibits how varied constellations of
759 stone tools were used to occupy a variety of geographic and environmental contexts. Typically,
760 we identified patterns of emphasis, rather than exclusivity, both for the configuration of stone
761 tools found together and where they have been used, offering support to qualitative
762 approaches that have struggled to identify clear patterns. Nevertheless, this quantitative
763 approach has been able to clarify a number of trends, particularly with respect to changes
764 through time. The use of some constellations of stone tools and occupation of some landscape
765 contexts appear to span most of the timeframe of the MSA, indicative of enduring behavioural
766 adaptations, and potentially highlighting refugia for periods of enhanced climatic stress.
767 Previous studies have split the MSA into early (MIS6 and older) and late (MIS5 and later)

768 components (e.g. Tryon & Faith 2013), emphasising the appearance of new elements of
769 behaviour in MIS 5. Rather than seeing a shift from early to late MSA behaviour, we have
770 illuminated that MIS5 is a phase in which Middle Pleistocene MSA behaviours continue to
771 occur but are significantly augmented with new combinations of stone tools appearing
772 alongside the colonisation of significantly different landscapes that are characteristic of the
773 Late Pleistocene. Further, targeted study is required to identify whether any particular stone
774 tool technologies offered different functionality that enabled such diversification, and the
775 results presented here can be significantly refined in the future by the inclusion of additional
776 technological data.

777 This study illustrates the importance of combining archaeological assemblage data with
778 information on the geographic, topographic, and environmental differences between the
779 locations from which those assemblages are recovered. Only with more comprehensive data
780 of this kind can we begin to link archaeological patterns with their potential abiotic causes. It
781 is clear from the analyses above that there do exist patterns of differences between sites, and
782 that there are partial explanations of such differences to be found in more detailed
783 considerations of chronology, geography, and various elements of the local environmental
784 setting. Environments inevitably change both across space and through time, and it is
785 therefore essential to consider all these components simultaneously if we are to arrive at a
786 comprehensive understanding of the archaeological record. It is also clear from the analyses
787 reported here that there is much unexplained variation within the MSA of eastern Africa. Much
788 of this no doubt arises from the stochastic nature of human behaviour and may relate both to
789 individual differences and to the patterns of contact and conflict between subpopulations that
790 were almost certainly isolated, both genetically and culturally, for long periods during the later
791 Middle and Late Pleistocene.

792 It is now apparent that considerable chronological overlap occurs between eastern Africa's
793 MSA industries with both the preceding Late Acheulean (e.g. Mieso dating to ~212ka [de la
794 Torre et al. 2014]) and the succeeding Later Stone Age (e.g. Panga ya Saidi dating from 67ka
795 [Shipton et al. 2018]). Beyond changes in emphasis in typological inventories, two key, inter-
796 related elements associated with both major transitions in eastern African prehistory are raw
797 material choice and artefact size. Due to the availability of comparable data, it has not been
798 possible to examine these features as potential contributing factors to changing constellations
799 of stone tool types. Similarly, to engage with the breadth of MSA assemblages in eastern Africa,
800 it has not been feasible to examine varying constellations of artefact typology relating to the
801 nature of different site types to explore, for instance, differences between logistical foraging
802 compared to residential sites. Both offer potential avenues to extend the application of the
803 quantitative analyses that are presented here.

804 Research into both the genetic and cultural foundations of *Homo sapiens* populations will
805 continue to add to the debate concerning the patterns of material culture observed here, and
806 in archaeological contexts from elsewhere in Africa and beyond. Analyses of the kind reported
807 above, however, enable us to formulate questions regarding contact and isolation between
808 groups that may be particularly amenable to future analyses of genetic and cultural
809 transmission. For example, these analyses demonstrate that many tool forms – borers, LCTs,
810 Levallois blade and point technologies, notched tools, and retouched knives – are *always*
811 present to some degree in the humid Marine Isotope Stages in eastern Africa, and yet they are
812 *never* present in the drier glacial stages. With various taphonomic caveats accepted, we can
813 ask whether such technologies are reinvented during the geographic and demographic
814 expansions associated with each humid stage, or whether they remain as unexpressed
815 elements of the repertoire throughout dry stages when, for reasons as yet unclear, they are

816 never physically realised. Similarly, Figures 8 and 9 establish geographic and environmental
817 clusters that are inhabited throughout the period studied here, and such refugial areas should
818 contain both the minimal distillation of MSA technology required for survival through arid
819 periods and the core populations from which subsequent expansions result. There is therefore
820 likely to be a close relationship between the smaller populations that survive less hospitable
821 stages and the contraction of technological diversity found here among assemblages from
822 MIS 8, 6, and 4. This relationship merits further study by archaeologists, but also by geneticists
823 and researchers working on the dynamics of cultural evolution in fluctuating environments.

824 **Acknowledgements:**

825 This research has been supported by the Natural Environment Research Council as part of
826 Grant: NE/K014560/1, "A 500,000-year environmental record from Chew Bahir, south Ethiopia:
827 testing hypotheses of climate-driven human evolution, innovation, and dispersal", which forms
828 part of the Hominin Sites and Paleolake Drilling Project. We thank two anonymous reviewers
829 and the editor for their thoughtful comments that have helped to improve the presentation of
830 this research.

831 The authors declare no competing interests.

832 **References Cited:**

833 Basell, L.S., 2008. Middle Stone Age (MSA) site distributions in eastern Africa and their
834 relationship to Quaternary environmental change , refugia and the evolution of Homo
835 sapiens. *Quaternary Science Reviews* 27, 2484–2498. doi:10.1016/j.quascirev.2008.09.010

836 Binford, L., 2001. *Constructing frames of reference: an analytical method for archaeological*
837 *theory building using ethnographic and environmental data sets.* University of California
838 Press.

839 Blegen, N., 2017. The earliest long-distance obsidian transport: Evidence from the ~ 200 ka
840 Middle Stone Age Sibilo School Road Site , Baringo , Kenya. *Journal of Human Evolution*
841 103, 1–19. doi:10.1016/j.jhevol.2016.11.002

842 Boeda, E., 1994. Le concept Levallois, Variabilité des méthodes. CNRS.

843 Braconnot, P., Otto-Bliesner, B., Harrison, S., Jousaume, S., Peterchmitt, J.-Y., Abe-Ouchi, a.,
844 Crucifix, M., Driesschaert, E., Fichet, T., Hewitt, C.D., Kageyama, M., Kitoh, a., Lâiné, a.,
845 Loutre, M.-F., Marti, O., Merkel, U., Ramstein, G., Valdes, P., Weber, S.L., Yu, Y., Zhao, Y.,
846 2007. Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial
847 Maximum – Part 1: experiments and large-scale features. *Climate of the Past* 3, 261–277.
848 doi:10.5194/cp-3-261-2007

849 Chorowicz, J., 2005. The east African rift system. *Journal of African Earth Sciences* 43(1-3),
850 379-410.

851 Clark, J.D., 1988. The Middle Stone Age of East Africa and the beginnings of regional identity.
852 *Journal of World Prehistory* 2, 235–305. doi:10.1007/BF00975618

853 Douze, 2012. Le Early Middle Stone Age d'Éthiopie et les changements techno-économiques
854 à la période de l'émergence des premiers Homo sapiens. L'Université Boredaux 1.

855 Faith, J.T., Tryon, C.A., Peppe, D.J., Beverly, E.J., Blegen, N., Blumenthal, S., Chritz, K.L., Driese,
856 S.G., Patterson, D., 2015. Paleoenvironmental context of the Middle Stone Age record
857 from Karungu , Lake Victoria Basin , Kenya , and its implications for human and faunal
858 dispersals in East Africa. *Journal of Human Evolution* 83, 28–45.
859 doi:10.1016/j.jhevol.2015.03.004

860 Groucutt, H.S., Petraglia, M.D., Bailey, G., Scerri, E.M.L., Parton, A., Clark-Balzan, L., Jennings,

861 R.P., Lewis, L., Blinkhorn, J., Drake, N.A., Breeze, P.S., Inglis, R.H., Devès, M.H., Meredith-
862 Williams, M., Boivin, N., Thomas, M.G., Scally, A., 2015. Rethinking the dispersal of *Homo*
863 *sapiens* out of Africa. *Evolutionary Anthropology: Issues, News, and Reviews* 24, 149–
864 164. doi:10.1002/evan.21455

865 Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G., Jarvis, A., 2005. Very high resolution
866 interpolated climate surfaces for global land areas. *International Journal of Climatology*
867 25, 1965–1978. doi:10.1002/joc.1276

868 Hublin, J., Ben-ncer, A., Bailey, S.E., Freidline, S.E., Neubauer, S., Skinner, M.M., Bergmann, I.,
869 Cabec, A. Le, Benazzi, S., Harvati, K., Gunz, P., 2017. New fossils from Jebel Irhoud,
870 Morocco and the pan-African origin of *Homo sapiens*. *Nature Publishing Group* 546,
871 289–292. doi:10.1038/nature22336

872 Irpino, A., 2017. HistDAWass: Histogram-Valued Data Analysis. doi:doi:10.1007/s11634-014-
873 0176-4

874 Jarvis, A., H.I. Reuter, A. Nelson, E.G., 2008. Hole-filled SRTM for the globe Version 4.
875 Available from the CGIAR-CSI SRTM 90m Database (<http://srtm.csi.cgiar.org>) 15.

876 Krzanowski, W., 2000. *Principles of multivariate analysis* (Vol. 23). Oxford: Oxford University
877 Press.

878 Lamb, H.F., Bates, C.R., Bryant, C.L., Davies, S.J., Huws, D.G., Marshall, M.H., Roberts, H.M.,
879 2018. 150,00-year palaeoclimate record from northern Ethiopia supports early , multiple
880 dispersals of modern humans from Africa. *Scientific Reports* 1–7. doi:10.1038/s41598-
881 018-19601-w

882 McDougall, I., Brown, F.H., Fleagle, J.G., 2005. Stratigraphic placement and age of modern

883 humans from Kibish, Ethiopia. *Nature* 433, 733–736. doi:10.1038/nature03258

884 Manly, B.F. and Alberto, J.A.N., 2016. *Multivariate statistical methods: a primer*. CRC Press.

885 Minetti, A.E., Gaudino, P., Seminati, E., Cazzola, D., Minetti, A.E., Moia, C., Roi, G.S., Susta, D.,
886 Ferretti, G., Holt, N.C., Askew, G.N., Minetti, A.E., Moia, C., Roi, G.S., Susta, D., Ferretti, G.,
887 Alberto, E., Moia, C., Roi, G.S., Susta, D., Ferretti, G., 2015. Energy cost of walking and
888 running at extreme uphill and downhill slopes *Energy cost of walking and running at*
889 *extreme uphill and downhill slopes* 1039–1046. doi:10.1152/jappphysiol.01177.2001

890 Otto-Bliesner, B.L., Marshall, S.J., Overpeck, J.T., Miller, G.H., Hu, A., 2006. Simulating Arctic
891 climate warmth and icefield retreat in the last interglaciation. *Science (New York, N.Y.)*
892 311, 1751–3. doi:10.1126/science.1120808

893 Rosenzweig, M. L., 1995. *Species diversity in space and time*. Cambridge: Cambridge
894 University Press.

895 Scerri, E.M.L., Drake, N.A., Jennings, R., Groucutt, H.S., 2014. Earliest evidence for the structure
896 of *Homo sapiens* populations in Africa. *Quaternary Science Reviews* 101, e179–e179.
897 doi:10.1016/j.quascirev.2014.07.019

898 Scerri, E.M.L., Thomas, M.G., Manica, A., Gunz, P., Stock, J., Stringer, C.B., Grove, M., Groucutt,
899 H.S., Timmermann A., Rightmire, G.P., d’Errico, F., Tryon, C., Drake, N.A., Brooks, A.,
900 Dennell, R., Durbin, R., Henn, B., Lee-Thorpe, J., deMenocal, P., Petraglia, M.D.,
901 Thompson, J., A., Scally, A., Chikhi, L., 2018. Did our species evolve in subdivided
902 populations across Africa, and why does it matter? *Trends in Ecology and Evolution (in*
903 *press)*.

904 Schlebusch, C.M., Malmström, H., Günther, T., Sjödin, P., Coutinho, A., Edlund, H., Munters,

905 A.R., Vicente, M., Steyn, M., Soodyall, H., Lombard, M., Jakobsson, M., 2017. Southern
906 African ancient genomes estimate modern human divergence to 350,000 to 260,000
907 years ago. *Science*. doi:10.1126/science.aao6266

908 Siepielski, A.M., Morrissey, M.B., Buoro, M., Carlson, S.M., Caruso, C.M., Clegg, S.M., Coulson,
909 T., DiBattista, J., Gotanda, K.M., Francis, C.D. and Hereford, J., 2017. Precipitation drives
910 global variation in natural selection. *Science* 355(6328), 959-962.

911 Shipton, C., Roberts, P., Archer, W., Armitage, S., Bitu, C., Blinkhorn, J., Courtney-Mustaphi, C.,
912 Crowther, A., Curtis, R., d'Errico, F., Douka, K., Faulkner, P., Groucutt, H., Helm, R., Herries,
913 A., Jembe, S., Kourampas, N., Lee-Thorp, J., Marchant, R., Mercader, J., Marti, A.,
914 Prendergast, M., Rowson, B., Tengeza, A., Tibesasa, R., White, T., Petraglia, M., Boivin, N.,
915 2018. 78,000-year-old record of Middle and Later Stone Age innovation in an East
916 African tropical forest. *Nature Communications* 9, 1832.

917 de la Torre, I., Mora, R., Arroyo, A. and Benito-Calvo, A., 2014. Acheulean technological
918 behaviour in the middle Pleistocene landscape of Mieso (East-Central Ethiopia). *Journal*
919 *of Human Evolution* 76, 1-25.

920 Tryon, C.A., Crevecoeur, I., Faith, J.T., Ekshtain, R., Nivens, J., Patterson, D., Mbua, E., Spoor, F.,
921 2015. Late Pleistocene age and archaeological context for the hominin calvaria from
922 GvJm-22. *Proceedings of the National Academy of Sciences* 112, 2682–2687.
923 doi:10.1073/pnas.1417909112

924 Tryon, C.A., Faith, J.T., 2013. Variability in the Middle Stone Age of Eastern Africa. *Current*
925 *Anthropology* 54 (S8), S234–S254. doi:10.1086/673752

926

927

928 **Figure Captions:**

929 **Figure 1:** Map illustrating the distribution of key sites and locations mentioned in the text on
930 an SRTM digital elevation model (Jarvis et al. 2008).

931 **Figure 2:** Binary heatmap of presence (black) and absence (grey) of twenty-six stone tool
932 types recorded in 125 eastern African MSA assemblages.

933 **Figure 3:** Dendrograms illustrating hierarchical clustering of stone tool type variables
934 amongst eastern African MSA sites using (left) agglomerative and (right) divisive
935 approaches.

936 **Figure 4:** Dendrograms illustrating hierarchical clustering of stone tool assemblages
937 amongst eastern African MSA sites using (left) agglomerative and (right) divisive
938 approaches.

939 **Figure 5:** Heatmaps illustrating percentage presence of artefact types within groups
940 identified by agglomerative (top) and divisive (bottom) clustering ranging from 100%
941 (red) to 0% (white).

942 **Figure 6:** Jitter plot illustrating the number of assemblages in each agglomerative cluster
943 found within each Marine Isotope Stage.

944 **Figure 7:** Jitter plot illustrating the number of assemblages in each divisive assemblage
945 cluster found within each Marine Isotope Stage.

946 **Figure 8:** Dendrograms illustrating hierarchical clustering of eastern African MSA sites based
947 upon geographic characteristics (altitude and energy)(left), and modern environments

948 (mean annual temperature and precipitation)(right) within a 50km radius of site
949 locations.


950 **Figure 9:** Jitter Plot illustrating the occupation of alternate geographic clusters (G1-6) split
951 between Marine Isotope Stages, with long term continuity of occupation evident in G1
952 and G3, and significant expansion observable in MIS 5.


953 **Figure 10:** Jitter plot illustrating occupation of alternate modern environmental clusters (E1-
954 5) split between Marine Isotope Stages, with long-term continuity of occupation
955 evident in E4, pulsed occupations of E2 in odd-numbered stages (7,5,3), and
956 expansions into new environments in MIS 5 and 3.

957 **Figure 11:** Dendrograms illustrating hierarchical clustering of eastern African MSA sites
958 based upon modelled environmental parameters (mean annual temperature and
959 precipitation) for the Last Glacial Maximum (21ka) as a proxy for arid conditions (left)
960 and for the last interglacial (MIS5e) as a proxy for humid conditions (right).

961 **Figure 12:** Jitter plot illustrating occupation of alternate arid environmental clusters (LGM1-5)
962 split between Marine Isotope Stages. Although repeated occupations of LGM2 and
963 LGM3 are observed throughout the MSA of eastern Africa, only LGM1 preserves
964 evidence for occupation during MIS4.

965 **Figure 13:** Jitter plot illustrating occupation of alternate humid environment clusters (LIG1-4)
966 split between Marine Isotope Stages. Both LIG1 and LIG3 indicate occupation within
967 both the Middle and Late Pleistocene. Occupation of LIG2 and LIG4 overlaps during
968 MIS 5 but are otherwise split between the Middle and Late Pleistocene respectively.







Biface
 Bipolar
 BladeTech
 Borer
 Burin
 Chopper
 CoreonFlake
 Denticulate
 DiscoidalCore
 Handaxe
 LCT
 LevalloisBladeTech
 LevalloisFlakeTech
 LevalloisPointTech
 Microlith
 Notch
 NotchedTool
 Pick
 PlatformCore
 PointTech
 RadialCore
 RTBifacial
 RTKnife
 RTPoint
 ScaledPieces
 Scraper

Abdur_N_C_S
 AbuHugar
 AdumaA1
 AdumaA4C
 AdumaA5Ex
 AdumaA5ExSurf
 AdumaA8
 AdumaA8AC
 AdumaA8AG
 AdumaA8ASurf
 AdumaA8B
 AdumaVP1/1
 AdumaVP1/3
 AlajSpringsMSA
 AlaKanasa
 Asfet
 CartwrightsSite
 EyasiShore_77_81
 EyasiShore_N_surface
 EyasiShore_W_insitu
 EyasiShore_W_surf
 EYM_Endingi
 Gademotta_ETH72_1
 Gademotta_ETH72_6
 Gademotta_ETH72_8B
 Gademotta_GDM7
 Garba3_S1
 Garba3_S2
 GodaButchia0_70
 Gotera
 GudGud
 Guomde
 KapedoTuffs
 KapForm_KoimilotGnJh74_1
 KapForm_KoimilotGnJh74_2
 KapForm_SSRS
 Karungu_A3Ex
 Karungu_GS
 Karungu_Kisaaka_Main
 Karungu_Kisaaka_ZIG
 Karungu_Obware
 Kone_Loc1AEx
 Kone_Loc4AEx
 Kone_Loc5_Extension
 Kone_Loc5Excav
 Kone_Loc6
 Kone_LocCEx
 Koobi_FwJi1
 Koobi_FwJi3
 Koobi_FxJi61Ex
 Koobi_FxJi61Surf
 Koobi_FxJi66
 Koobi_GaJi17
 LaasGeel_SU_711
 Loichoro_1
 LukenyaHill_GvJm16_A
 LukenyaHill_GvJm46
 LukenyaHill_GvJm22_F170_205
 LukenyaHill_GvJm22_G210_225
 Magubike_MSA
 MalewaGorge
 Midishi2_LSU1II
 Midishi2_LSUIV
 Midishi2_LSU
 Midishi2_LSU
 Midishi2_LSU
 MochenaBorago_LowerT
 MochenaBorago_RGroup
 MochenaBorago_SGroup
 MochenaBorago_UpperT
 Mtongwe_ELG
 Mtongwe_EMG
 Mtongwe_EUG
 Muguruk_AggA
 Muguruk_AggB
 Muguruk_Member4
 Muguruk_Member6
 Mumba_L_VI_38
 Mumba_L_VI_81
 Mumba_L_VI_38
 Mumba_L_VI_A
 Mumba_MU_V_81
 Mumba_U_V_38
 Mumba_U_VI_A
 Mumba_VI_B
 Naisiusiu_1931
 Naisiusiu_1969insitu
 Naisiusiu_1972
 Nakechikchok
 Nasera_12_17
 Nasera_18_25
 Nasera_6_7
 Nasera_8/9_11
 Ndutu_14
 Ndutu_72
 Nyara_River
 Ologesallie_BOK1E
 Ologesallie_BOK2
 Ologesallie_BOK3
 Ologesallie_BOK4
 Ologesallie_GOK1
 Omo_AHS1-5
 Omo_AHS6_8
 Omo_AHSsurface
 Omo_BNS_L3
 Omo_BNS<50m
 Omo_KHS2/3
 Omo_KHSNully
 Omo_KHSNMKenya
 Omo_KHSSgully
 PangeyaSajdi_19_18
 PorcEpic_III
 PorcEpic_IV
 ProlongedDrift
 ProspectFarm_I
 ProspectFarm_II
 ProspectFarm_III
 ProspectFarm_IV
 Rusinga_Kakrigu
 Rusinga_Nyamita
 Rusinga_Nyamsingula
 Rusinga_Wakondo
 Shurmai_MSA
 Simbi
 Singa
 SonjoBuri

0.0 0.2 0.4 0.6 0.8 1.0







0.0 0.2 0.4 0.6 0.8 1.0




0.0 0.2 0.4 0.6 0.8 1.0


0.0 0.2 0.4 0.6 0.8 1.0














Agglomerative Cluster













Arid (LGM) Cluster





1 **The structure of the Middle Stone Age of eastern Africa: Supplementary Information**

2 **Table SI.1:** Assemblages used in analysis, identifying site/area name, assemblage name,
 3 latitude, longitude Marine Isotope Stage and references.

Site	Assemblage	N	E	MIS	Ref
Abdur	AN, AC and AS	15.13	39.68	5	Bruggemann 2004; Walter et al. 2000
Abu Hugar	Abu Hugar	12.86	33.99	NA	Clark 1988
Aduma	A1	10.393	40.537	5	Yellen 2005
Aduma	A4 Contact	10.393	40.537	5	Yellen 2005
Aduma	A5 Excavated	10.393	40.537	5	Yellen 2005
Aduma	A5 (excavated surface)	10.393	40.537	5	Yellen 2005
Aduma	A8	10.393	40.537	5	Yellen 2005
Aduma	A8A Contact	10.393	40.537	5	Yellen 2005
Aduma	A8A Gravel	10.393	40.537	5	Yellen 2005
Aduma	A8A surface	10.393	40.537	5	Yellen 2005
Aduma	A8B	10.393	40.537	5	Yellen 2005
Aduma	VP 1/1	10.393	40.537	5	Yellen 2005
Aduma	VP 1/3	10.393	40.537	5	Yellen 2005
Aladi Springs	Aladi Springs	9.25	40.75	2	Gossa et al. 2012
Ala Kanasa	Ala Kanasa	10.52	40.53	NA	Clark 1988
Asfet	Asfet	15.29	39.91	5	Beyin 2013
Cartwrights Site	Cartwrights Site	-0.63	36.46	NA	Waweru 2007
Eyasi Shore	77_81	-3.54	35.28	5	Mehlman 1989
Eyasi Shore	Nordostbucht surface	-3.54	35.28	5	Mehlman 1989
Eyasi Shore	Westbucht in situ	-3.54	35.28	5	Mehlman 1989
Eyasi Shore	Westbucht surface	-3.54	35.28	5	Mehlman 1989
Enkapune ya Muto	RBL4	0.00	36.15	NA	Basell 2008
Gademotta	ETH_72_1	7.55	38.57	7	Douze 2012
Gademotta	ETH_72_6	7.55	38.57	6	Douze 2012
Gademotta	ETH_72_8B	7.55	38.57	8	Douze 2012
Gademotta	GDM7	7.55	38.57	NA	Sahle et al. 2013
Garba 3	Sample 1	8.41	38.34	5	Mussi 2013
Garba 3	Sample 2	8.41	38.34	5	Mussi 2013
Goda Butchia	0-70cm	9.541	41.62	3	Pleurdeau et al. 2014
Gotera	Gotera	4.70	37.40	NA	Basell 2008
GudGud	GudGud	10.87	48.26	3	Basell 2008
Guomde	Guomde	3.61	36.52	8	Basell 2008
KapedoTuffs	Combined	1.07	36.08	5	Tryon et al. 2008
Kapthurin Formation	Koimilot GnJh74 Locus 1	0.517	35.975	8	Tryon 2003

Kapthurin Formation	Koimilot GnJh74 Locus 2	0.517	35.975	7	Tryon 2003
Kapthurin Formation	SSRS	0.657	35.99	8	Blegen et al. 2017
Karungu	Aringo 3 Excavation	-0.83	34.18	5	Faith et al. 2015
Karungu	General Survey	-0.80	34.13	NA	Faith et al. 2015
Karungu	Kisaaka-Main	-0.80	34.13	3	Faith et al. 2015
Karungu	Kisaaka-ZTG	-0.80	34.13	5	Faith et al. 2015
Karungu	Obware	-0.87	34.22	NA	Faith et al. 2015
Kone	Loc1_A_TrialEx	8.43	39.66	4	Kurashina 1978
Kone	Loc4_A_TrialEx	8.43	39.66	4	Kurashina 1978
Kone	Loc5_Extension	8.43	39.66	4	Kurashina 1978
Kone	Loc5_1975ex	8.43	39.66	4	Kurashina 1978
Kone	Loc6	8.43	39.66	4	Kurashina 1978
Kone	LocC_testEx	8.43	39.66	4	Kurashina 1978
Koobi_Fora	FwJi 1	4.28	36.25	NA	Kelly 1996
Koobi_Fora	FwJi 3	4.30	36.22	NA	Kelly 1996
Koobi_Fora	FxJi 61 Excav	4.04	36.43	NA	Kelly 1996
Koobi_Fora	FxJi 61 Surf	4.04	36.43	NA	Kelly 1996
Koobi_Fora	FxJj 66	4.05	36.42	NA	Kelly 1996
Koobi_Fora	GaJj 17	3.97	36.29	NA	Kelly 1996
Laas Geel	SU_711	9.68	44.27	3	Gutherez et al. 2014
Lolchoro	1	-2.97	35.96	NA	Seitsonen 2005
LukenyaHill_GvJm16	A	-1.65	37.14	NA	Merrick 1975
LukenyaHill_GvJm46	LukenyaHill_GvJm46	-1.65	37.14	3	Kelly 1996; Basell 2008
Lukenya Hill GvJm22	F - 170 - 205	-1.65	37.14	NA	Tryon et al. 2015
Lukenya Hill GvJm22	G - 210-225	-1.65	37.14	NA	Tryon et al. 2015
Magubike	TP1/2/3	-7.76	35.47	3	Bushozi 2011
Malewa Gorge	Malewa Gorge	-0.77	36.39	7	Basell 2008
Midishi 2	LSU III	10.60	47.73	2	Tryon & Faith 2013
Midishi 2	LSU IV	10.60	47.73	3	Tryon & Faith 2013
Midishi 2	LSU V	10.60	47.73	3	Tryon & Faith 2013
Midishi 2	LSU VI	10.60	47.73	3	Tryon & Faith 2013
Mochena Borago	Lower T	6.90	37.74	3	Brandt et al. 2017
Mochena Borago	R Group	6.90	37.74	3	Brandt et al. 2017
Mochena Borago	S Group	6.90	37.74	3	Brandt et al. 2017
Mochena Borago	Upper T	6.90	37.74	3	Brandt et al. 2017
Mtongwe	East_Lower_Group	-4.08	39.65	5	Tryon & Faith 2013
Mtongwe	East_Middle_Group	-4.08	39.65	5	Tryon & Faith 2013
Mtongwe	East Upper Group	-4.08	39.65	5	Tryon & Faith 2013
Muguruk	Agg A	-0.09	34.63	NA	McBrearty 1988
Muguruk	Agg B	-0.09	34.63	NA	McBrearty 1988

Muguruk	Member 4	-0.09	34.63	NA	McBrearty 1986
Muguruk	Member 6	-0.09	34.63	NA	McBrearty 1986
Mumba	Lower III 1938 G_H	-3.54	35.29	3	Mehlman 1989
Mumba	Lower_V_77_81	-3.54	35.29	3	Mehlman 1989
Mumba	Lower V_1938_G	-3.54	35.29	3	Mehlman 1989
Mumba	Lower VI_A	-3.54	35.29	5	Mehlman 1989
Mumba	Middle Upper V 77_81	-3.54	35.29	3	Mehlman 1989
Mumba	Upper V 1938 B	-3.54	35.29	3	Mehlman 1989
Mumba	Upper VI_A	-3.54	35.29	4	Mehlman 1989
Mumba	VI_B	-3.54	35.29	5	Mehlman 1989
Naisiusiu	1931 material	-3.25	34.86	3	Leakey et al. 1972
Naisiusiu	1969 in situ material	-3.25	34.86	3	Leakey et al. 1972
Naisiusiu	1972_surface	-3.25	34.86	3	Leakey et al. 1972
Nakechikchok	Nakechikchok	3.15	35.8	NA	Shea et al. 2010
Nasera	12_17	-2.73	35.35	3	Mehlman1989
Nasera	18_25	-2.73	35.35	3	Mehlman1989
Nasera	6_7	-2.73	35.35	2	Mehlman1989
Nasera	8/9_11	-2.73	35.35	3	Mehlman1989
Ndutu	Ndutu	-3.25	34.86	9	Eren et al. 2014
Ndutu	Ndutu	-3.25	34.86	9	Leakey et al. 1972
Nyara_River	Nyara_River	-8.97	33.46	NA	Basell 2008
Olorgesailie	BOK1E	-1.56	36.43	9	Brooks et al. 2018
Olorgesailie	BOK2	-1.56	36.43	9	Brooks et al. 2018
Olorgesailie	BOK3	-1.56	36.43	9	Brooks et al. 2018
Olorgesailie	BOK4	-1.56	36.43	9	Brooks et al. 2018
Olorgesailie	GOK1	-1.56	36.43	8	Brooks et al. 2018
Omo	AHS 1-5	4.46	35.97	7	Shea et al. 2008
Omo	AHS 6_8	4.46	35.97	7	Shea et al. 2008
Omo	AHS surface	4.46	35.97	7	Shea et al. 2008
Omo	BNS_L3	4.46	35.97	5	Shea et al. 2008
Omo	BNS<50m	4.46	35.97	5	Shea et al. 2008
Omo	KHS 2/3	4.46	35.97	7	Shea et al. 2008
Omo	KHS N gully	4.46	35.97	7	Shea et al. 2008
Omo	KHS NM Kenya	4.46	35.97	7	Shea et al. 2008
Omo	KHS S gully	4.46	35.97	7	Shea et al. 2008
Pange ya Saidi	19_18	-3.72	39.71	5	Shipton et al 2018
Porc Epic	III	9.56	41.88	3	Pleurdeau 2003
Porc Epic	IV	9.56	41.88	3	Pleurdeau 2003
ProlongedDrift	ProlongedDrift	-0.76	35.94	3	Waweru 2007
ProspectFarm	I	-0.93	36.11	NA	Anthony 1978

ProspectFarm	II	-0.93	36.11	5	Anthony 1978
ProspectFarm	III	-0.93	36.11	5	Anthony 1978
ProspectFarm	IV	-0.93	36.11	3	Anthony 1978
Rusinga	Kakrigu	-0.45	34.05	NA	Tryon & Faith 2013
Rusinga	Nyamita	-0.42	34.15	5	Tryon & Faith 2013
Rusinga	Nyamsingula	-0.40	34.18	5	Tryon & Faith 2013
Rusinga	Wakondo	-0.41	34.17	5	Tryon & Faith 2013
Shurmai	MSA	0.50	37.21	3	Dickson & Gang
Simbi	Simbi	-0.54	35.04	5	Basell 2008
Singa	Singa	11.99	34.20	NA	Basell 2008
SonjoBuri	SonjoBuri	-2.34	36.04	NA	Seitsonen 2005

4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

24 **Table SI.2:** Description of assemblage composition of clusters identified using an
 25 agglomerative approach.

A1 (n=46)	This cluster is unique in including at least one occurrence of every stone tool type, but no single type is ubiquitous. The most numerous types are <i>Levallois Flake Technology</i> (93.5%), <i>Scrapers</i> (89.1%), <i>RT Points</i> (73.9%), with <i>Discooidal Cores</i> (65.2%), <i>Blade Technology</i> and <i>Borers</i> (both 58.7%) occurring in more than half of the assemblages.
A2 (n=3)	All three members of this cluster include <i>Handaxes</i> , with 2/3 including <i>Blade Technology</i> , and 1/3 including <i>Levallois Flake Technology</i> and <i>Scaled Pieces</i> , but no other tool types are present.
A3 (n=11)	All members of this cluster include scrapers, with <i>Point Technology</i> (81.8%), and <i>Notches</i> and <i>Radial Cores</i> (both 63.6%) appear in more than half of the assemblages, with a total of 15 artefact types represented.
A4 (n=21)	All members of this cluster include <i>Platform Cores</i> , with 10 other types appearing in more than half of the assemblages, including <i>Scrapers</i> , <i>Radial Cores</i> and <i>Levallois Flake Technology</i> (all 85.7%), and <i>Blade Technology</i> and <i>Microliths</i> (both 71.4%) appearing in high frequency. Only 6 types are entirely absent.
A5 (n=10)	<i>RT Points</i> are found in all members of this cluster, with <i>Levallois Flake Technology</i> and <i>LCT's</i> occurring in 6/10 sites, and <i>Platform Cores</i> , <i>Levallois Point Technology</i> and <i>Bifaces</i> appearing in 5/10 sites, and 8 other types appear at lower frequencies.
A6 (n=8)	All members of this cluster include <i>Microliths</i> , with <i>Blade Technology</i> present in 5/8 of the sites, and <i>Discooidal Cores</i> , <i>Scrapers</i> and <i>Scaled Pieces</i> appear in half the assemblages. 16 tool types are absent from this cluster.
A7 (n=26)	Two types, <i>Levallois Flake Technology</i> and <i>Blade Technology</i> occur in 96.2% of assemblages, though no types are entirely ubiquitous. In addition to this, only scrapers appear in more than half of the assemblages, though only two tool types are completely absent



26
 27
 28
 29
 30
 31
 32
 33
 34

35 **Table SI.2:** Description of assemblage composition of clusters identified using an divisive
 36 approach.


D1 (n=45)	No types appear in all members of this assemblage, although <i>Scrapers</i> (97.8%), <i>Levallois Flake Technology</i> and <i>RT Points</i> (both 91.1%) occur in high frequency, followed by <i>Borers</i> (82.2%), <i>Blade Technology</i> (75.6%), and <i>Notches</i> (68.9%), with at least one occurrence of all stone tool types present.
D2 (n=17)	<i>Scrapers</i> are ubiquitous in this cluster, although beyond this only <i>Microliths</i> and <i>Radial Cores</i> (both 58.8%) appear in more than half the assemblages, whereas 9 tool types are entirely absent.
D3 (n=3)	All members of this cluster include <i>Microliths</i> , with 2/3 including <i>Blade Technology</i> , <i>Platform Cores</i> , <i>Bipolar Tech</i> and <i>Discooidal Cores</i> , and a single <i>Pick</i> occurs.
D4 (n=9)	<i>Blade Technology</i> , <i>Scrapers</i> and <i>Levallois Flake Technology</i> are ubiquitous in this group, with high proportions of both <i>Levallois Blade</i> and <i>Levallois Point Technologies</i> (88.9%).
D5 (n=4)	All four members of this cluster include <i>Bifaces</i> , <i>Levallois Point Technology</i> and <i>RT Points</i> , with $\frac{3}{4}$ including <i>LCT's</i> and <i>Platform Cores</i> , out of a total of 10 artefacts present.
D6 (n=20)	<i>Scrapers</i> are ubiquitous in this cluster, with <i>Levallois Flake Technology</i> (90%), <i>Discooidal Cores</i> (80%) and <i>Platform Cores</i> (70%) present in more than half of the assemblages, with only 6 artefact types entirely absent.
D7 (n=23)	All members of this cluster contain <i>Levallois Flake Technology</i> but none of the remaining 16 artefact types present appear in more than half of the assemblages.
D8 (n=4)	All four members of this cluster includes handaxes, with 2/4 including <i>Blade Technology</i> and <i>Levallois Flake Technology</i> , with single occurrences of <i>Levallois Flake Technology</i> , <i>Borers</i> , <i>Denticulates</i> , <i>Levallois Blade Technology</i> , <i>Point Technology</i> , <i>RT Knife</i> , and <i>Scaled Pieces</i> .

37
 38
 39
 40
 41
 42
 43
 44
 45


46 **Figure SI.1:** Distribution maps of sites grouped by hierarchical clustering using either
47 agglomerative (left) or divisive (right) methods, suggesting most cluster types are widely
48 (although at times sparsely) distributed, with limited spatial structuring.





50 **Figure SI.2:** (left) Distribution of Geographic Clusters; (top right) jitter plot of assemblage
 51 clusters (agglomerative) by geographic clusters; (bottom right) jitter plot of assemblage
 52 clusters (divisive) by geographic clusters.





54 **Figure SI.3:** (left) Distribution of Modern Environment Clusters; (top right) jitter plot of
 55 assemblage clusters (agglomerative) by modern environment clusters; (bottom right) jitter
 56 plot of assemblage clusters (divisive) by modern environment clusters.



58 **Figure SI.4:** Distribution of Arid Environmental Clusters; (top right) jitter plot of assemblage
 59 clusters (agglomerative) by arid environmental clusters; (bottom right) jitter plot of
 60 assemblage clusters (divisive) by arid environmental clusters.



62 **Figure SI.5:** Distribution of Humid Environmental Clusters; (top right) jitter plot of
 63 assemblage clusters (agglomerative) by humid environmental clusters; (bottom right) jitter
 64 plot of assemblage clusters (divisive) by humid environmental clusters.



66 **References Cited:**

- 67 Anthony, B.W., 1978. *The Prospect Industry: A Definition*. Harvard University.
- 68 Basell, L.S., 2008. Middle Stone Age (MSA) site distributions in eastern Africa and their
69 relationship to Quaternary environmental change, refugia and the evolution of *Homo*
70 *sapiens*. *Quaternary Science Reviews* 27, 2484–2498. doi:10.1016/j.quascirev.2008.09.010
- 71 Beyin, A., 2013. A surface Middle Stone Age assemblage from the Red Sea coast of Eritrea:
72 Implications for Upper Pleistocene human dispersals out of Africa. *Quaternary*
73 *International* 300, 195–212. doi:http://dx.doi.org/10.1016/j.quaint.2013.02.015
- 74 Blegen, N., 2017. The earliest long-distance obsidian transport: Evidence from the ~ 200 ka
75 Middle Stone Age Sibilo School Road Site, Baringo, Kenya. *Journal of Human Evolution*
76 103, 1–19. doi:10.1016/j.jhevol.2016.11.002
- 77 Brandt, S., Hildebrand, E., Vogelsang, R., Wolfhagen, J., Wang, H., 2017. *Journal of*
78 *Archaeological Science: Reports* A new MIS 3 radiocarbon chronology for Mochena
79 Borago Rockshelter, SW Ethiopia: Implications for the interpretation of Late Pleistocene
80 chronostratigraphy and human behavior. *Journal of Archaeological Science: Reports* 11,
81 352–369. doi:10.1016/j.jasrep.2016.09.013
- 82 Brooks, A.S., Yellen, J.E., Potts, R., Behrensmeier, A.K., Deino, A.L., Leslie, D.E., Ambrose, S.H.,
83 Ferguson, J.R., d’Errico, F., Zipkin, A.M. and Whittaker, S., 2018. Long-distance stone
84 transport and pigment use in the earliest Middle Stone Age. *Science* 360(6384), 90-94.
- 85 Bruggemann, J.H., Bu, R.T., Guillaume, M.M.M., Walter, R.C., Cosel, R. Von, Ghebretensae, B.N.,
86 Berhe, S.M., 2004. Stratigraphy, palaeoenvironments and model for the deposition of
87 the Abdur Reef Limestone: context for an important archaeological site from the last
88 interglacial on the Red Sea coast of Eritrea. *Palaeogeography, Palaeoclimatology,*
89 *Palaeoecology* 203, 179–206. doi:10.1016/S0031-0182(03)00659-X
- 90 Bushozi, P., 2011. *Lithic technology and hunting behaviour during the Middle Stone Age in*
91 *Tanzania*. University of Alberta.
- 92 Clark, J.D., 1988. The Middle Stone Age of East Africa and the beginnings of regional identity.
93 *Journal of World Prehistory* 2, 235–305. doi:10.1007/BF00975618
- 94 Dickson, D.B., Gang, G.-Y., 2002. Evidence of the Emergence of “Modern” Behavior in the
95 Middle and Later Stone Age Lithic Assemblages at Shurmai Rockshelter (GnJm1) and
96 Kakwa Lelash Rockshelter (GnJm2) in the Mukogodo Hills of North-Central Kenya.
97 *African Archaeological Review* 19, 1–26. doi:10.1023/A:1014490818099
- 98 Douze, 2012. *Le Early Middle Stone Age d’Éthiopie et les changements techno-économiques*
99 *à la période de l’émergence des premiers Homo sapiens*. L’Université Boredeaux 1.
- 100 Eren, M.I., Durant, A.J., Prendergast, M., Mabulla, A.Z.P., 2014. Middle Stone Age archaeology

- 101 at Olduvai Gorge , Tanzania. *Quaternary International* 323, 292–313.
102 doi:10.1016/j.quaint.2013.12.042
- 103 Faith, J.T., Tryon, C.A., Peppe, D.J., Beverly, E.J., Blegen, N., Blumenthal, S., Chritz, K.L., Driese,
104 S.G., Patterson, D., 2015. Paleoenvironmental context of the Middle Stone Age record
105 from Karungu , Lake Victoria Basin , Kenya , and its implications for human and faunal
106 dispersals in East Africa. *Journal of Human Evolution* 83, 28–45.
107 doi:10.1016/j.jhevol.2015.03.004
- 108 Gossa, T., Sahle, Y., Negash, A., 2012. A reassessment of the Middle and Later Stone Age lithic
109 assemblages from Aladi Springs, Southern Afar Rift, Ethiopia. *Azania: Archaeological*
110 *Research in Africa* 47, 210–222. doi:10.1080/0067270X.2012.676314
- 111 Gutherz, X., Diaz, A., Ménard, C., Bon, F., Douze, K., Léa, V., Lesur, J., Sordoillet, D., 2014. The
112 Hargeisan revisited: Lithic industries from shelter 7 of Laas Geel, Somaliland and the
113 transition between the Middle and Late Stone Age in the Horn of Africa. *Quaternary*
114 *International* 343, 69–84. doi:10.1016/j.quaint.2014.04.038
- 115 Kelly, A.J., 1996. Intra-regional and inter-regional variability in the East Turkana (Kenya) and
116 Kenyan Middle Stone Age. Rutgers.
- 117 Kurashina, H., 1978. An examination of prehistoric lithic technology in East-Central Ethiopia.
118 University of California.
- 119 Leakey, M.D., Hay, R.L., Thurber, D.L., Protsch, R., Berger, R., 1972. Stratigraphy, Archaeology,
120 and Age of the Ndotu and Naisiusiu Beds, Olduvai Gorge, Tanzania. *World Archaeology*
121 3, 328–341.
- 122 McBrearty, S.A., 1986. The archaeology of the Muguruk site, western Kenya. University of
123 Illinois.
- 124 Mehlman, M.J., 1989. Late Quaternary archaeological sequences in northern Tanzania.
125 University of Illinois.
- 126 Merrick, H. V, 1975. Change in Late Pleistocene lithic industries in eastern Africa. University of
127 California.
- 128 Mussi, M., Altamura, F., Macchiarelli, R., Melis, R.T., Spinapolice, E.E., 2013. Garba III (Melka
129 Kunture , Ethiopia) : a MSA site with archaic Homo sapiens remains revisited. *Quaternary*
130 *International* 343, 28–39. doi:10.1016/j.quaint.2013.08.028
- 131 Pleurdeau, D., 2003. Le Middle Stone Age de la grotte du Porc-Épic (Dire Dawa , Éthiopie) :
132 gestion des matières premières et comportements techniques Porc-Epic cave (Dire
133 Dawa , Ethiopia) Middle Stone Age: raw materials economy and technical behaviour.
134 *L'anthropologie* 107, 15–48.
- 135 Pleurdeau, D., Hovers, E., Assefa, Z., Asrat, A., 2014. Cultural change or continuity in the late

136 MSA / Early LSA of southeastern Ethiopia: The site of Goda Buticha , Dire Dawa area.
137 Quaternary International 343, 117–135.

138 Potts, R., Behrensmeier, A.K., Faith, J.T., Tryon, C.A., Brooks, A.S., Yellen, J.E., Deino, A.L.,
139 Kinyanjui, R., Clark, J.B., Haradon, C.M. and Levin, N.E., 2018. Environmental dynamics
140 during the onset of the Middle Stone Age in eastern Africa. *Science* 360 (6384), 86–90.
141 doi: 10.1126/science.aao2200

142 Sahle, Y., Morgan, L.E., Braun, D.R., Atnafu, B., Hutchings, W.K., 2013. Chronological and
143 behavioral contexts of the earliest Middle Stone Age in the Gademotta Formation, Main
144 Ethiopian Rift. *Quaternary International*. doi:10.1016/j.quaint.2013.03.010

145 Seitsonen, O., 2005. Stone age sequence, lithic technology, and ancient lake level changes in
146 the North Tanzanian Rift valley area. University of Helsinki.

147 Shea, J.J., 2008. The Middle Stone Age archaeology of the Lower Omo Valley Kibish
148 Formation: Excavations, lithic assemblages, and inferred patterns of early *Homo sapiens*
149 behavior. *Journal of Human Evolution* 55, 448–485. doi:10.1016/j.jhevol.2008.05.014

150 Shea, J.J., Hildebrand, E.A., 2010. The Middle Stone Age of West Turkana , Kenya. *Journal of*
151 *Field Archaeology* 35, 355–364. doi:10.1179/009346910X12707321358630

152 Shipton, C., Roberts, P., Archer, W., Armitage, S., Bitu, C., Blinkhorn, J., Courtney-Mustaphi, C.,
153 Crowther, A., Curtis, R., d’Errico, F., Douka, K., Faulkner, P., Groucut, H., Helm, R., Herries,
154 A., Jembe, S., Kourampas, N., Lee-Thorp, J., Marchant, R., Mercader, J., Marti, A.,
155 Prendergast, M., Rowson, B., Tengeza, A., Tibesasa, R., White, T., Petraglia, M., Boivin, N.,
156 2018. 78,000-year-old record of Middle and Later Stone Age innovation in an East
157 African tropical forest. *Nature Communications* 9, 1832.

158 Tryon, C.A., 2003. The Acheulian to Middle Stone Age Transition: Tephrostratigraphic Context
159 for Archaeological Change in the Kapthurin Formation, Kenya. University of Connecticut.

160 Tryon, C.A., Faith, J.T., 2013. Variability in the Middle Stone Age of Eastern Africa. *Current*
161 *Anthropology* 54 (S8), S234–S254. doi:10.1086/673752

162 Tryon, C.A., Peppe, D.J., Faith, J.T., Van, A., Nightingale, S., Ogondo, J., Fox, D.L., 2012.
163 Archaeological Research in Africa Late Pleistocene artefacts and fauna from Rusinga and
164 Mfangano islands , Lake Victoria, Kenya. *Azania* 14–38.
165 doi:10.1080/0067270X.2011.647946

166 Tryon, C.A., Roach, N.T., Logan, M.A. V, 2008. The Middle Stone Age of the northern Kenyan
167 Rift: age and context of new archaeological sites from the Kapedo Tuffs. *Journal of*
168 *Human Evolution* 55, 652–664. doi:10.1016/j.jhevol.2008.03.008

169 Walter, R.C., Buffler, R.T., Bruggemann, J.H., Guillaume, M.M.M., Berhe, S.M., Negassi, B.,
170 Libsekal, Y., Cheng, H., Edwards, R.L., von Cosel, R., Néraudeau, D., Gagnon, M., 2000.
171 Early human occupation of the Red Sea coast of Eritrea during the last interglacial.

172 Nature 405, 65–69. doi:10.1038/35011048

173 Waweru, V.N., 2007. Middle Stone Age Technology at Cartwright’s Site, Kenya. University of
174 Conneticut.

175 Yellen, J., Brooks, A., Helgren, D., Tappen, M., Ambrose, S., Bonnefille, R., Feathers, J.,
176 Goodfriend, G., Ludwig, K., Renne, P., Stewart, K., 2005. The Archaeology of Aduma
177 Middle Stone Age Sites in the Awash Valley , Ethiopia. *PaleoAnthropology* 10, 25–100.

178