

University of Groningen

## Kinetics of TH2 biomarkers in sputum of asthmatics following inhaled allergen

Zuiker, Rob G. J. A.; Ruddy, Marcella K.; Morelli, Nicoletta; Mogg, Robin; Rivas, Veronica M.; van Dyck, Kristien; De Lepeleire, Inge; Tanen, Michael R. L.; Boot, J. Diderik; Kamerling, Ingrid M. C.

*Published in:*  
European clinical respiratory journal

*DOI:*  
[10.3402/ecrj.v2.28319](https://doi.org/10.3402/ecrj.v2.28319)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2015

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Zuiker, R. G. J. A., Ruddy, M. K., Morelli, N., Mogg, R., Rivas, V. M., van Dyck, K., ... Diamant, Z. (2015). Kinetics of TH2 biomarkers in sputum of asthmatics following inhaled allergen. *European clinical respiratory journal*, 2(1), [28319]. DOI: 10.3402/ecrj.v2.28319

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



## Kinetics of TH2 biomarkers in sputum of asthmatics following inhaled allergen

Rob G. J. A. Zuiker, Marcella K. Ruddy, Nicoletta Morelli, Robin Mogg, Veronica M. Rivas, Kristien van Dyck, Inge De Lepeleire, Michael R. L. Tanen, J. Diderik Boot, Ingrid M. C. Kamerling & Zuzana Diamant

To cite this article: Rob G. J. A. Zuiker, Marcella K. Ruddy, Nicoletta Morelli, Robin Mogg, Veronica M. Rivas, Kristien van Dyck, Inge De Lepeleire, Michael R. L. Tanen, J. Diderik Boot, Ingrid M. C. Kamerling & Zuzana Diamant (2015) Kinetics of TH2 biomarkers in sputum of asthmatics following inhaled allergen, *European Clinical Respiratory Journal*, 2:1, 28319, DOI: [10.3402/ecrj.v2.28319](https://doi.org/10.3402/ecrj.v2.28319)

To link to this article: <https://doi.org/10.3402/ecrj.v2.28319>



© 2015 Rob G. J. A. Zuiker et al.



Published online: 26 May 2015.




Submit your article to this journal [↗](#)



Article views: 89



View Crossmark data [↗](#)



ORIGINAL RESEARCH ARTICLE

## Kinetics of TH2 biomarkers in sputum of asthmatics following inhaled allergen

Rob G. J. A. Zuiker<sup>1†</sup>, Marcella K. Ruddy<sup>2†</sup>, Nicoletta Morelli<sup>3</sup>, Robin Mogg<sup>4</sup>, Veronica M. Rivas<sup>5</sup>, Kristien van Dyck<sup>6</sup>, Inge De Lepeleire<sup>6</sup>, Michael R. L. Tanen<sup>5</sup>, J. Diderik Boot<sup>7</sup>, Ingrid M. C. Kamerling<sup>1</sup> and Zuzana Diamant<sup>8,9,10\*</sup>

<sup>1</sup>Centre for Human Drug Research, Leiden, The Netherlands; <sup>2</sup>EMD Serono, Rockland, MA, USA; <sup>3</sup>VU Medical Center, Amsterdam, The Netherlands; <sup>4</sup>Janssen Pharmaceutical Companies of Johnson & Johnson, Spring House, PA, USA; <sup>5</sup>Merck Research Laboratories, Rahway, NJ, USA; <sup>6</sup>Merck Research Laboratories, Brussels, Belgium; <sup>7</sup>HAL Allergy B.V., Leiden, The Netherlands; <sup>8</sup>Skane University Hospital, Department of Respiratory Medicine and Allergology, Institute for Clinical Science, Skane University, Lund, Sweden; <sup>9</sup>Departments of Clinical Pharmacy & Pharmacology and General Practice, University Medical Center Groningen, Groningen, The Netherlands; <sup>10</sup>QPS Netherlands, Groningen, The Netherlands

**Background:** Allergen-induced late airway response offers important pharmacodynamic targets, including T helper 2 (TH2) biomarkers. However, detection of inflammatory markers has been limited in dithiothreitol-processed sputum.

**Objectives:** To test whether allergen-induced TH2 inflammatory markers can be reproducibly quantified by sensitive detection techniques in ultracentrifuged sputum and the effect of fluticasone (FP) on these endpoints.

**Methods:** Thirteen allergic asthmatics with dual allergen-induced airway responses, documented during a single-blind placebo run-in period, participated in a double-blind, two-period crossover study. Each period consisted of three consecutive days, separated by  $\geq 3$  weeks. Following randomization, subjects inhaled FP (500  $\mu\text{g}$  bid, five doses total) or placebo. On Day 2 in each study period, allergen challenge was performed and airway response measured by forced expiratory volume in 1 sec (FEV1) until 7 h post-challenge. Sputum was induced 24 h pre-allergen and 7 and 24 h post-allergen. Sputum samples were split into two portions: TH2 biomarkers were quantified by Meso Scale multiplex platform following ultracentrifugation, and cell differentials were counted on Giemsa–May–Grünwald-stained cytopins. Allergen-induced changes in inflammatory endpoints were compared between FP and placebo using a mixed model ANCOVA.

**Results:** Inhaled allergen induced dual airway responses in all subjects during both placebo periods with reproducible late asthmatic response (LAR) and increased sputum inflammatory biomarkers (IL-2, IL-4, IL-13, and eotaxin-1) and eosinophil counts. FP effectively blunted both the LAR and the inflammatory biomarkers.

**Conclusions:** Combining novel, sensitive quantification methods with ultracentrifugation allows reproducible quantification of sputum biomarkers following allergen challenge, reversed by FP. This approach allows non-invasive identification of pharmacodynamic targets for anti-asthma therapies.

Keywords: *allergen bronchial provocation test; asthma; sputum; TH2 inflammation; fluticasone*

\*Correspondence to: Zuzana Diamant, Skane University Hospital, Department of Respiratory Medicine & Allergology, Institute for Clinical Science, Skane University, Klinikgatan 18, SE-22181 Lund, Sweden, Email: z.diamant@gems.demon.nl

Received: 22 April 2015; Accepted in revised form: 22 April 2015; Published: 26 May 2015

Inhaled allergen challenge is a highly reproducible, integral disease model enabling the investigation of several features of asthma (1). Allergen challenge can

be applied to study the pathophysiology and, if complemented with (non-)invasive airway samplings, the immunobiology to allergic stimuli within the airways. In drug

<sup>†</sup>These authors have contributed equally to this work.

development, allergen challenge is an established tool predicting clinical efficacy of novel anti-allergic and anti-asthma treatments (2).

Non-invasive airway sampling by hypertonic saline-induced sputum (3) has been shown to yield reproducible increases in inflammatory cells and biomarkers following allergen-induced late asthmatic response (LAR) (4) with subsequent response to novel and existing anti-inflammatory therapies (2, 4–6). Although animal studies have provided evidence of TH2 cytokine response following allergen challenge, supported by some human studies applying bronchoscopy (7, 8), no consistent data exist on reproducible quantification of TH2 cytokines and chemokines in sputum. Accountable factors include degradation by standard sputum processing with dithiothreitol (DTT), which destroys the disulfide bounds of these inflammatory markers (9), overall low baseline concentrations, and relatively insensitive detection techniques. Some of these hurdles could be overcome by physical homogenization of sputum samples by ultracentrifugation, causing cellular disruption with subsequent release of intracellular products, in combination with sensitive detection techniques (10, 11).

Combining sputum ultracentrifugation with novel, sensitive quantification techniques using Meso Scale multi-array microplates (12) in the allergen challenge model, we aimed to study the following: 1) the feasibility of the quantification of TH2 cytokines and chemokines in sputum at 7 and 24 h post-challenge, 2) their reproducibility, and 3) their reversibility after a short course of inhaled fluticasone (FP). Furthermore, to allow comparison with other established markers of allergen-induced airway inflammation, we also measured the allergen-induced airway responses (i.e. the early asthmatic response [EAR] and LAR), exhaled nitric oxide (eNO), sputum cell differentials, and the provocative concentration of methacholine causing a fall in forced expiratory volume in 1 sec ( $FEV_1$ ) of 20% ( $PC_{20}FEV_1$  methacholine) at baseline and 24 h post-allergen, during all study periods.

## Methods

### Study population and design

Thirteen non-smoking subjects participated in a double-blind, two-way crossover study. Participants had clinically stable, mild to moderate allergic asthma (13), using *prn* short-acting  $\beta$ -2 agonists only and with dual airway responses to inhaled house dust mite (HDM), documented during the single-blind placebo run-in screening period. Each period consisted of three consecutive days, with a washout period  $\geq 3$  weeks between periods (Fig. 1). The screening, allowing to test the reproducibility of the variables, was identical to the subsequent treatment periods, during which subjects randomly received inhaled FP (metered dose inhaler [MDI], 500  $\mu$ g bid, a total of five doses) or matching placebo. On Day 1, baseline measurements including eNO, spirometry followed by methacholine challenge ( $PC_{20}FEV_1$  methacholine), and subsequent sputum induction ( $3 \times 5$  min NaCl 4.5%) were performed prior to study medication. On Day 2, 1 h post-study medication, subjects underwent a titrated allergen challenge (1). The subsequent airway response was repeatedly measured by  $FEV_1$  up to 7 h post-allergen. eNO was measured pre-allergen and at 3 h and 7 h post-allergen, the latter measurement followed by sputum induction. At 24 h post-allergen (Day 3), test procedures were repeated as on Day 1 (Fig. 2). All test procedures were conducted according to standardized, validated methods and at the same time of the day (within 2 h) (1, 14–16).

A dual airway response to inhaled HDM extract consisted of an EAR and a LAR, defined as a fall in  $FEV_1 \geq 15\%$  from baseline occurring between 0–3 h and 3–7 h post-allergen, respectively.

The study was approved by the Ethics Committee of Leiden University Medical Center, Leiden, The Netherlands, and all participants provided signed informed consent (EUDRACT number 2007-003671-40). All procedures were performed in accordance with the Helsinki Declaration of 1975, revised in 2008.





Fig. 1. Overview of the single-blind placebo run-in period and double-blind crossover study periods 1 and 2.



**Fig. 2.** Overview of study assessments. IS, induced sputum; eNO, exhaled nitric oxide. Time zero is time of first study medication dosing. The single-blind placebo run-in screening period and the two subsequent study periods were identical.

### Study medication and dosing rationale

FP 250  $\mu\text{g}/\text{puff}$  (Allen & Hanburys, Glaxo Wellcome Ltd, Middlesex, UK) and matching placebo (Armstrong Pharmaceuticals, Inc., Canton, MA, USA, packaged at Merck Frosst, Kirkland, Canada) were supplied in identical MDIs and inhaled per single puff through an AeroChamber (Volumatic; GlaxoSmithKline, Zeist, The Netherlands). The rationale for the dose regimen was based on a previous study showing substantial reductions in allergen-induced LAR, non-specific airway hyperresponsiveness (AHR), and sputum eosinophils already following one single dose of inhaled FP 250  $\mu\text{g}$  (6). Thus, to ensure optimal reversal of the allergen-induced inflammatory markers versus placebo, a total of five FP doses (500  $\mu\text{g}$  per dose) were administered throughout the active treatment period.

### Allergen challenge

The allergen challenge was performed using the 2 min tidal breathing method that has been previously validated (1). The run-in period served as a dose (range)-finding procedure, whereas during the first and second study periods each subject inhaled the same two or three cumulative doses of the allergen extract that had caused a fall in FEV<sub>1</sub> of at least 15% from baseline during the run-in period. Following diluent, incremental doubling concentrations (7.81–2,000 BU/mL) of HDM extract (*Dermatophagoides pteronyssinus*; SQ 503, ALK-BPT, ALK-Abelló, Almere, The Netherlands) in phosphate-buffered saline (PBS) (Invitrogen cat. no. 14040) were aerosolized by a calibrated jet-nebulizer (DeVilbiss 646, output 0.13 mL/min, Somerset, PA, USA) and inhaled at approximately 12 min intervals, until the EAR was reached (defined as a decrease in FEV<sub>1</sub> of  $\geq 15\%$  from post-diluent baseline within 1 h post-allergen). Airway response to inhaled allergen was measured by FEV<sub>1</sub> in duplicate on a calibrated spirometer (Vmax Spectra; Sensor Medics, Bilthoven, The Netherlands) according to standard procedures (17), at 10, 20, 30, 45, 60, 90, and 120 min and then hourly until

7 h after the last allergen inhalation. The highest technically valid measurement was expressed as percentage decrease from the post-diluent baseline FEV<sub>1</sub> and included in the analysis.

### Methacholine challenge

The methacholine challenge was performed using standard methodologies (15). Serial doubling concentrations of methacholine bromide (MBr; Janssen Pharmaceutical, Beerse, Belgium), diluted in normal saline (NaCl 0.9%) to serial doubling dilutions of 0.15–80  $\mu\text{mol}/\text{mL}$ , were aerosolized by a calibrated jet nebulizer (DeVilbiss 646) at 5-min intervals and inhaled by the subjects by tidal breathing for 2 min through the mouthpiece with the nose clipped. Airway response was measured by FEV<sub>1</sub> at 30 and 90 sec (and potentially at 180 sec as well) following each concentration, and the lowest technically satisfactory FEV<sub>1</sub> was incorporated into the analysis. Nebulization was continued until a  $\geq 20\%$  fall in FEV<sub>1</sub> from the post-diluent baseline.

After both bronchoprovocation tests, subjects received salbutamol through an AeroChamber, until the FEV<sub>1</sub> returned to within 10% of the baseline value.

### Exhaled nitric oxide

All eNO measurements were performed according to current guidelines (14) using a chemiluminescence analyzer (Ecomedics CLD88sp; Ecomedics, Duernten, Switzerland), which had to be replaced by a NIOX MINO<sup>®</sup> (Aerocrine AB, Solna, Sweden) during the study. The NIOX MINO was used for subjects 8, 9, 10, 11, 12, and 13 during both study periods. In a previous study at our institute, both analyzers yielded similar values (18).

### Sputum induction, processing, and analysis

Sputum induction was performed as previously described (16, 19) using a DeVilbiss Ultraneb 2000 ultrasonic nebulizer (Tefa Portanje, Woerden, The Netherlands) connected to a 100-cm long plastic tube, with an internal diameter of approximately 22 mm, connected to a two-way

valve (No. 2700; Hans-Rudolf, Kansas City, MO, USA) with a mouthpiece. Hypertonic saline (NaCl 4.5%) was nebulized and inhaled through the mouth, with the nose clipped, during three periods of 5 min. At approximately 7 min after each induction, spirometry was performed as a safety measure.

Collected sputum samples were divided into two portions of equal weight. The cell pellet of the first portion was processed as a full sample according to guidelines (16, 20), using 0.1% DTT (Sputolysin; Calbiochem, La Jolla, CA, USA). Cell viability and total cell count were assessed using trypan blue; sputum samples containing >80% squamous cells were excluded from analysis. Differential cell counts were performed by a qualified cytologist on May-Grünwald-Giemsa-stained, coded cytopins and expressed as a percentage of 500 nucleated, non-squamous cells.

The second sputum portion was used to quantify soluble inflammatory markers. At Merck Research Laboratories, defrosted samples were pretreated with a protease-inhibitor cocktail (50 µL per 200 mg sputum), prepared by dissolving one protease cocktail tablet (Complete Protease Inhibitor Cocktail tablets, Roche Applied Science no. 11697498001) into 50 mL of PBS. Prepared sputum samples were subsequently ultracentrifuged in an ultracentrifuge (Optima Max Ultracentrifuge, 130,000 rpm; Beckman Coulter, Inc., Fullerton, CA, USA) at 35,000 rpm (53,500 × *g*) for 90 min at 4°C. Subsequently, sputum supernatant was collected and analyzed.

### Cytokine and chemokine measurements

Quantification of soluble biomarkers in sputum samples was performed using an MSD (Meso Scale Discovery, Gaithersburg, MD, USA) singleplex kit (IL-13), an MSD duplex kit (eotaxin-3 and TARC), and two MSD multiplex assays (IL-1β, IL-2, IL-4, IL-5, IL-8, IL-10, IL-12p70, IFN-γ, TNF-α, eotaxin, IP-10, MCP-1, MCP-4, and MIP-1β). All concentrations were expressed as pg/mL.

### Statistical analysis

Data from all randomized subjects were included in the analysis.

The effect of FP versus placebo on the TH2 cytokines, chemokines, and other inflammatory markers at 7 and 24 h post-allergen was assessed using a mixed-effects analysis of variance (ANOVA) model. The model included fixed factors for sequence, treatment, and period and a random effect for subjects within sequence. Between-treatment differences were estimated by the difference in least-squares means from the model with 90% confidence interval (CI; one-sided alpha = 5%). Sputum cell differentials were analyzed using the actual change from baseline, whereas absolute cell counts were analyzed using the change from baseline for the square root transformed values. Geometric mean baseline sputum biomarker concentra-

tions were calculated; half of the lower limit of quantification was used in case of negative outcomes. Changes in sputum biomarker concentrations were analyzed after log transformation and expressed as fold change from baseline.

The airway response to inhaled allergen was expressed as percentage decrease in FEV<sub>1</sub> from the post-diluent baseline and plotted as time–response curves during all treatment periods. The difference in FEV<sub>1</sub> during both the EAR and the LAR was analyzed using the time-weighted average of percentage change and the maximum percentage change from baseline. Subject 1 had an initial FEV<sub>1</sub> decrease of slightly under 15% at 7 h post-allergen, but met the inclusion criterion at 8 h post-allergen and was included in the study. Therefore, for this subject FEV<sub>1</sub>, cytokines, chemokines, and eNO were consequently measured at 8 h during all periods. FEV<sub>1</sub> results at 8 h were not included in the analysis.

PC<sub>20</sub>FEV<sub>1</sub>methacholine was calculated by linear interpolation on a plot of log concentrations versus response using methacholine concentrations below and above a 20% fall in FEV<sub>1</sub>. The (allergen-induced and FP-reverted) changes in PC<sub>20</sub>FEV<sub>1</sub>methacholine were expressed in doubling doses. eNO was expressed as a fold change from baseline at 3, 7, and 24 h post-allergen.

Reproducibility of the allergen-induced airway responses and sputum inflammatory markers was assessed using data from the run-in and study placebo periods. The intraclass correlation coefficient (ICC) was calculated, and a two-sided paired *t*-test was performed.

### Sample size

In the absence of information about variability in TNF-α and IL-13 concentrations in sputum, eosinophil count was used as an approximate variable for sample size estimation (21). Power calculation showed that the study would have >90% power (α = 0.05, one-tailed) to detect a fivefold increase from baseline at 7 h post-allergen challenge with 12 completing subjects.

## Results

### Subjects

After completion of the run-in period, 15 subjects were considered eligible. Before randomization, two subjects were withdrawn: one had a positive cotinine test, whereas the other repeatedly presented with a clinically relevant bronchoconstriction (baseline FEV<sub>1</sub> < 70% predicted). Thus, 13 subjects were randomized and all completed the study (Table 1).

### Safety

No serious adverse events occurred. Headache and fatigue were the most frequently reported adverse events.



**Table 1.** Baseline characteristics of randomized subjects

Number of subjects	13
Age, years	25.9 (21–43)
Gender	4M/9F
BMI, kg/m <sup>2</sup>	24.4 (16.6–39.8)
FEV <sub>1</sub> , L	3.57 (2.92–4.50)
FEV <sub>1</sub> , % pred	94.0 (74.5–112.3)
PC <sub>20</sub> FEV <sub>1</sub> methacholine, µmol/mL	12.8 (0.8–81.5)
SPT HDM Wheal, mm	5.5 (2.5–10.5)
eNO, ppb	53.4 (11.2–160.8)

Numbers are expressed as mean (range). BMI, body mass index; FEV<sub>1</sub>, forced expiratory volume in 1 sec; SPT HDM, skin prick test for house dust mite; ppb, parts per billion.

All events were mild in intensity and classified as unrelated to the study medication or procedures.

### Allergen-induced airway responses

Inhaled HDM induced both an EAR and an LAR in all subjects during both placebo periods. Compared to placebo, FP significantly reduced the EAR and completely blunted the LAR (Fig. 3). The reproducibility of the allergen-induced LAR during both placebo periods was good, both in terms of the maximum percent fall in FEV<sub>1</sub> from baseline and as time-weighted average (3–7 h post-allergen), with an ICC of 79.7 and 69%, respectively (Table 2).

### Allergen-induced non-specific AHR

During both placebo periods, allergen challenge increased non-specific AHR, by decreasing PC<sub>20</sub>FEV<sub>1</sub>methacholine at 24 h post-allergen by, on average, 1.18 (90% CI: 1.73, 0.64) doubling doses. In contrast, FP increased 24 h

post-allergen PC<sub>20</sub>FEV<sub>1</sub>methacholine by on average 1.60 doubling doses (90% CI: 1.06, 2.15), resulting in a mean difference of 2.79 doubling doses (90% CI: 2.07, 3.51;  $p < 0.001$ ) between placebo and FP (Fig. 4).


### Sputum inflammatory cells

A sputum sample was obtained from all subjects on all occasions. The average squamous cell contamination was 36% (range 2–71%). Of the 117 samples, 16 were not analyzable. Inhaled allergen significantly increased sputum eosinophils both at 7 and 24 h post-challenge during both placebo periods. This effect was significantly reduced by FP (Table 3). The reproducibility for both sputum eosinophil count (ICC 76%) and percentage (ICC 88%) was high at 7 h post-allergen, but poor (ICC 0%) at 24 h.

### Sputum (TH2) cytokines and chemokines

During placebo treatment, inhaled allergen increased sputum inflammatory cytokines and chemokines both at 7 and 24 h post-allergen, yielding the most robust increase at 7 h (Table 4). FP significantly blunted the allergen-induced increases in sputum concentrations of IL-5, IL-13, TARC, eotaxin-3, MCP-1, eotaxin-1, and IL-4 at 7 h post-allergen challenge and of IL-5, IL-13, eotaxin-3, IL-12p70, and MCP-1 at 24 h post-allergen challenge. None of the other sputum soluble markers were significantly affected by FP compared to placebo treatment. At 24 h post-allergen, there were no differences in any sputum inflammatory markers, with the exception of TARC between both placebo treatments.

At 7 h post-allergen, many soluble markers were reproducible; in particular, IL-2, IL-4, IL-13, and eotaxin-1 showed ICC values greater than 50%, with more variation between subjects than within subjects. At 24 h,



**Fig. 3.** Time-response curves (mean  $\pm$  SEM) to inhaled allergen during run-in period, placebo treatment, and fluticasone treatment.

Table 2. Analysis of the airway response to inhaled allergen

FEV <sub>1</sub> endpoint	Treatment	% estimate <sup>a</sup> (90% CI)	Difference in % change between fluticasone and placebo (90% CI)	Reproducibility	
				One-sided p-value <sup>c</sup>	Change <sup>b</sup> (90% CI) Two-sided p-value <sup>c</sup> ICC (%)
EAR time-weighted mean	Placebo	-7.09 (-8.90, -5.28)	5.28 (3.28, 7.28)	<0.001	4.0 (0.8, 7.3) 0.020
	Fluticasone (500 mcg bid)	-1.81 (-3.62, 0.00)			
EAR maximum decrease	Placebo	-17.7 (-21.3, -14.1)	11.12 (8.26, 13.99)	<0.001	6.3 (1.0, 11.7) 0.025
	Fluticasone (500 mcg bid)	-6.56 (-10.1, -2.99)			
LAR time-weighted mean	Placebo	-13.8 (-17.0, -10.6)	15.08 (10.97, 19.18)	<0.001	2.6 (-0.8, 6.0) 0.121
	Fluticasone (500 mcg bid)	1.28 (-1.89, 4.44)			
LAR maximum decrease	Placebo	-25.9 (-31.1, -20.7)	24.00 (16.70, 31.31)	<0.001	0.0 (-4.7, 4.6) 0.982
	Fluticasone (500 mcg bid)	-1.89 (-7.06, 3.28)			
24 h	Placebo	-5.30 (-8.16, -2.44)	7.05 (4.45, 9.64)	<0.001	3.7 (0.4, 6.9) 0.030
	Fluticasone (500 mcg bid)	1.75 (-1.11, 4.61)			

<sup>a</sup>Percentage change from period baseline; <sup>b</sup>placebo period vs. (placebo) run-in period; <sup>c</sup>paired t-test; EAR, early asthmatic response; LAR, late asthmatic response; CI, confidence interval; ICC, intraclass correlation coefficient. <sup>\*</sup>p-value: fluticasone vs. placebo, one-sided alpha = 5%.




Fig. 4. Changes in airway hyperresponsiveness 24 h pre-allergen versus 24 h post-allergen during run-in period, placebo treatment, and fluticasone treatment.

none of the inflammatory markers had ICC values greater than 50%.

Cytokine baseline values on Day 1 for each treatment period are provided in Table 5.

### Change in eNO

Compared to baseline, eNO levels did not significantly increase at 3 and 7 h post-allergen and did not differ between placebo and FP. At 24 h post-allergen, however, a significant increase in eNO was measured (1.63-fold, 90% CI: 1.2, 2.3), which was blunted by FP (0.83-fold, 90% CI: 0.6, 1.2), resulting in a significant difference between placebo and FP of 49% ( $p=0.012$ , 90% CI: 19, 68).

### Discussion

In this study, we were able to reproducibly quantify several TH2 inflammatory cytokines and chemokines in sputum from allergic asthmatic subjects following inhaled allergen. The increase in these soluble sputum biomarkers was consistent with other established allergen-induced inflammatory responses and was most robust at 7 h post-allergen, coinciding with the maximal fall in FEV<sub>1</sub> during the LAR. FP significantly blocked both the allergen-induced airway response and the majority of the inflammatory markers in sputum. Although other researchers previously showed a similar inflammatory response in bronchoalveolar lavage (7) and in sputum (22–24), none of them has investigated such wide range of allergen-induced TH2-cytokines and chemokines or their reversibility with corticosteroid treatment.



*Table 3.* Analysis of the eosinophils and neutrophils to inhaled allergen

Sputum differential	Time (h)	Treatment	% estimate (90% CI) <sup>a</sup>	% change from baseline		Reproducibility		
				Fluticasone vs. placebo (90% CI)	<i>p</i> -value*	Change <sup>b</sup> (90% CI)	Two-sided <i>p</i> -value <sup>c</sup>	ICC (%)
Eosinophils (count)	7	Placebo	250.8 (165.5; 336.1)	-277 (-394; -160)	0.002	-37.4 (-168, 93.5)	0.528	88.1
		Fluticasone	-25.9 (-106; 54.5)					
	24	Placebo	217.9 (142.7; 293.0)	-271 (-377; -165)	0.001	-118 (-244, 8.17)	0.063	0.0
		Fluticasone	-52.9 (-128; 22.3)					
Eosinophils (percent)	7	Placebo	11.5 (5.7; 17.4)	-11.8 (-19.9; -3.8)	0.014			
		Fluticasone	-0.3 (-5.8; 5.2)					
	24	Placebo	6.8 (1.7; 11.8)	-7.8 (-12.3, -3.3)	0.007			
		Fluticasone	-1.1 (-6.1; 3.9)					
Neutrophils (count)	7	Placebo	173.6 (62.5, 284.6)	-101 (-253, 52.0)	0.124	58.8 (-206, 324)	0.623	15.1
		Fluticasone	73.0 (-31.7, 177.7)					
	24	Placebo	224.6 (33.1, 416.2)	-190 (-450, 70.6)	0.103	219 (-48.2, 487)	0.095	28.0
		Fluticasone	35.0 (-157, 226.5)					
Neutrophils (percent)	7	Placebo	-5.5 (-18.8, 7.9)	6.3 (-9.3, 21.8)	0.768			
		Fluticasone	0.8 (-11.9, 13.5)					
	24	Placebo	-5.3 (-20.9, 10.3)	-4.6 (-24.7, 15.5)	0.335			
		Fluticasone	-9.9 (-25.5, 5.6)					

<sup>a</sup>Sputum eosinophils and neutrophils expressed as % change from baseline at 7 and 24 h post-allergen challenge; <sup>b</sup>placebo period vs. run-in period; <sup>c</sup>paired *t*-test.

\**p*-value: fluticasone vs. placebo, one-sided alpha = 5%.

Table 4. Analysis of sputum cytokines and chemokines to inhaled allergen

C	Hours post -allergen challenge	Treatment	Estimate <sup>a</sup> (90% CI)	% change (fluticasone vs. placebo) (90% CI), <i>p</i> -value*	Reproducibility.		
					Change <sup>b</sup> (90% CI)	Two-sided <i>p</i> -value <sup>c</sup>	ICC (%)
IL-5	7	Placebo	5.67 (3.63; 8.86)	82.6 (71.4; 89.4)	1.27 (0.53; 3.06)	0.555	48.1
		Fluticasone	0.99 (0.63; 1.55)	<i>p</i> < 0.001			
	24	Placebo	2.57 (1.54; 4.30)	66.1 (29.9; 83.6)	0.69 (0.24, 2)	0.465	34.7
		Fluticasone	0.87 (0.52; 1.46)	<i>p</i> = 0.011			
IL-13	7	Placebo	9.76 (6.25; 15.25)	88.6 (81.2; 93.1)	1.13 (0.63; 2.02)	0.655	73.6
		Fluticasone	1.11 (0.71; 1.73)	<i>p</i> < 0.001			
	24	Placebo	2.11 (1.52; 2.94)	59.7 (37.0; 74.2)	0.66 (0.36, 1.19)	0.148	41.2
		Fluticasone	0.85 (0.61; 1.18)	<i>p</i> = 0.002			
TARC	7	Placebo	2.13 (1.68; 2.72)	41.9 (22.8; 56.3)	0.9 (0.49; 1.66)	0.709	43.3
		Fluticasone	1.24 (0.97; 1.58)	<i>p</i> = 0.003			
	24	Placebo	1.75 (1.32; 2.32)	21.4 (-9.4; 43.6)	0.49 (0.29, 0.82)	0.012	17.9
		Fluticasone	1.37 (1.04; 1.82)	<i>p</i> = 0.108			
Eotaxin-3	7	Placebo	2.24 (1.71; 2.94)	44.6 (18.4; 62.4)	1.25 (0.55; 2.81)	0.556	0.0
		Fluticasone	1.24 (0.94; 1.64)	<i>p</i> = 0.010			
	24	Placebo	2.56 (1.80; 3.63)	39.2 (12.0; 58.0)	0.73 (0.29, 1.81)	0.458	22.2
		Fluticasone	1.55 (1.09; 2.21)	<i>p</i> = 0.018			
MCP-1	7	Placebo	1.27 (1.03; 1.56)	28.2 (9.1; 43.3)	1.14 (0.91; 1.44)	0.230	32.2
		Fluticasone	0.91 (0.74; 1.12)	<i>p</i> = 0.015			
	24	Placebo	0.96 (0.81; 1.15)	22.8 (4.1; 37.8)	1.07 (0.82, 1.4)	0.571	8.0
		Fluticasone	0.74 (0.62; 0.89)	<i>p</i> = 0.028			
Eotaxin-1	7	Placebo	2.01 (1.43; 2.82)	47.5 (15.2; 67.6)	0.68 (0.36; 1.3)	0.219	52.0
		Fluticasone	1.05 (0.75; 1.48)	<i>p</i> = 0.018			
	24	Placebo	1.25 (0.89, 1.75)	-0.7 (-61.9, 37.3)	0.65 (0.26, 1.63)	0.329	0.0
		Fluticasone	1.26 (0.90, 1.76)	<i>p</i> = 0.511			
IL-4	7	Placebo	1.52 (1.21; 1.92)	34.0 (8.9; 52.2)	1.02 (0.74; 1.41)	0.899	72.3
		Fluticasone	1.01 (0.80; 1.27)	<i>p</i> = 0.021			
	24	Placebo	0.98 (0.92, 1.03)	-4.7 (-13.6, 3.5)	0.9 (0.72, 1.12)	0.312	0.0
		Fluticasone	1.02 (0.96, 1.08)	<i>p</i> = 0.836			
MCP-4	7	Placebo	1.34 (0.96, 1.87)	32.4 (-3.3, 55.7)	1.05 (0.6; 1.85)	0.839	0.0
		Fluticasone	0.91 (0.65, 1.27)	<i>p</i> = 0.063			
	24	Placebo	1.36 (0.98, 1.89)	14.6 (-36.2, 46.5)	0.87 (0.44, 1.7)	0.646	1.6
		Fluticasone	1.16 (0.83, 1.61)	<i>p</i> = 0.277			
IL-12	7	Placebo	1.47 (0.97, 2.24)	29.0 (-24.9, 59.6)	1.56 (0.91; 2.68)	0.095	38.5
		Fluticasone	1.05 (0.69, 1.59)	<i>p</i> = 0.149			
	24	Placebo	1.61 (1.07; 2.42)	50.6 (12.2; 72.2)	0.85 (0.46, 1.54)	0.555	26.5
		Fluticasone	0.79 (0.53; 1.19)	<i>p</i> = 0.025			
IP-10	7	Placebo	1.11 (0.85, 1.46)	19.2 (-16.6, 44.0)	1.12 (0.64; 1.95)	0.671	0.0
		Fluticasone	0.90 (0.69, 1.18)	<i>p</i> = 0.159			
	24	Placebo	1.04 (0.71, 1.53)	15.4 (-42.1, 49.7)	0.94 (0.52, 1.69)	0.809	0.0
		Fluticasone	0.88 (0.60, 1.30)	<i>p</i> = 0.286			
MIP-1β	7	Placebo	1.70 (1.16, 2.48)	17.8 (-40.5, 51.9)	1.52 (0.66; 3.52)	0.296	14.6
		Fluticasone	1.40 (0.96, 2.04)	<i>p</i> = 0.261			
	24	Placebo	1.70 (1.08, 2.67)	-11.9 (-97.1, 36.5)	0.8 (0.25, 2.55)	0.687	2.7
		Fluticasone	1.90 (1.21, 2.99)	<i>p</i> = 0.637			
IL-8	7	Placebo	1.01 (0.80, 1.28)	4.0 (-31.6, 30.0)	0.93 (0.64; 1.36)	0.697	39.3
		Fluticasone	0.97 (0.77, 1.23)	<i>p</i> = 0.409			
	24	Placebo	1.26 (0.95, 1.66)	8.4 (-35.4, 38.0)	0.88 (0.53, 1.46)	0.583	29.9
		Fluticasone	1.15 (0.87, 1.52)	<i>p</i> = 0.347			

Table 4 (Continued)

C	Hours post -allergen challenge	Treatment	Estimate <sup>a</sup> (90% CI)	% change (fluticasone vs. placebo) (90% CI), <i>p</i> -value*	Reproducibility.		
					Change <sup>b</sup> (90% CI)	Two-sided <i>p</i> -value <sup>c</sup>	ICC (%)
IL-10	7	Placebo	1.18 (0.94, 1.48)	-1.8 (-40.0, 26.0) <i>p</i> = 0.539	1.17 (0.85; 1.63)	0.300	30.5
		Fluticasone	1.20 (0.96, 1.50)				
	24	Placebo	1.24 (0.95, 1.61)	9.7 (-17.2, 30.3) <i>p</i> = 0.247	0.97 (0.61, 1.52)	0.867	11.0
		Fluticasone	1.12 (0.86, 1.45)				
IL-1 $\beta$	7	Placebo	1.02 (0.77, 1.34)	-10.6 (-60.0, 23.6) <i>p</i> = 0.684	1.02 (0.64; 1.62)	0.933	42.6
		Fluticasone	1.12 (0.85, 1.48)				
	24	Placebo	1.17 (0.80, 1.72)	3.2 (-45.1, 35.5) <i>p</i> = 0.443	1.16 (0.6, 2.28)	0.628	8.9
		Fluticasone	1.13 (0.77, 1.67)				
IL-2	7	Placebo	0.90 (0.70, 1.16)	-13.9 (-59.2, 18.5) <i>p</i> = 0.752	0.81 (0.64; 1.02)	0.069	72.4
		Fluticasone	1.03 (0.80, 1.32)				
	24	Placebo	1.01 (0.72, 1.42)	-19.6 (-79.6, 20.3) <i>p</i> = 0.778	0.82 (0.42, 1.6)	0.530	13.4
		Fluticasone	1.21 (0.86, 1.70)				
IFN- $\gamma$	7	Placebo	1.05 (0.78, 1.43)	-41.1 (-101, 1.0) <i>p</i> = 0.946	1.19 (0.64; 2.23)	0.552	0.0
		Fluticasone	1.49 (1.10, 2.02)				
	24	Placebo	1.03 (0.77, 1.38)	-10.8 (-66.7, 26.4) <i>p</i> = 0.670	0.84 (0.4, 1.75)	0.611	0.0
		Fluticasone	1.14 (0.86, 1.53)				
TNF- $\alpha$	7	Placebo	1.14 (0.88, 1.50)	-43.6 (-84.1, -12.0) <i>p</i> = 0.988	1.18 (0.73; 1.92)	0.460	0.0
		Fluticasone	1.64 (1.26, 2.15)				
	24	Placebo	1.21 (0.93, 1.56)	-4.3 (-43.0, 23.9) <i>p</i> = 0.593	0.86 (0.43, 1.71)	0.634	0.0
		Fluticasone	1.26 (0.97, 1.63)				

<sup>a</sup>Sputum cytokines and chemokines in fold change from baseline at 7 and 24 h post-allergen challenge after placebo and fluticasone treatment; <sup>b</sup>placebo period vs. run-in period; <sup>c</sup>paired *t*-test.

\**p*-value: fluticasone vs. placebo, one-sided alpha = 5%.

The use of sulphhydryl-reducing reagents, such as DTT, has complicated the detection of inflammatory cytokines and chemokines, and alternative processing techniques enabling the measurement of eotaxin, for example, have previously been published (9). In our study, sputum samples were ultracentrifuged (10), instead of being processed with DTT, to avoid potentially degrading effects on several TH2 cytokines and chemokines (9). Following this 'boosting' step, substantial allergen-induced increases in several cytokines and chemokines could be reproducibly quantified using sensitive detection techniques (Meso Scale multi-array microplates). However, reproducibility was lost for most soluble markers and sputum eosinophils at 24 h post-allergen.

In parallel with reproducible increases in the TH2-derived inflammatory markers, we were able to demonstrate reproducible changes in the established allergen-induced inflammatory outcome (4, 25, 26), including the late asthmatic airway response, non-specific AHR, and sputum eosinophils, underscoring the validity of our data. In agreement with previous evidence, we also found increased eNO levels at 24 h post-allergen (27), while no significant eNO increases could be observed at our cutoff

point during the LAR, specifically, at 7 h post-allergen. Although previous studies showed increased eNO levels at 9 and 10 h post-allergen ((27) and (28), respectively), the present findings can be explained by the use of two different measuring devices (for logistic reasons) and the time lag required for the synthesis of inducible NO synthase, responsible for the synthesis of NO (29).

Although no direct comparison was made in the present study with soluble markers from the DTT-processed sputum portion, the current approach yielded reproducible data. In addition, the observation that FP can reverse the allergen-induced increase in these inflammatory markers in parallel with its inhibitory effects on the other inflammatory events, including the airway responses and cellular markers, suggests that this approach is sensitive enough to offer evaluation of therapeutic interventions in asthmatic subjects.

In conclusion, combining novel, sensitive quantification methods with ultracentrifugation allows reproducible quantification of sputum biomarkers following an allergen-induced LAR, which can be reversed by FP. This approach allows non-invasive identification of pharmacodynamic targets for anti-asthma therapies.

Table 5. Mean\* baseline values of cytokines and chemokines during placebo and fluticasone treatment

Cytokine/chemokine	IL-5	IL-13	TARC	Eotaxin-3	MCP-1	Eotaxin-1	IL-4	MCP-4	IL-12	IP-10	MIP-1 $\beta$	IL-8	IL-10	IL-1 $\beta$	IL-2	IFN- $\gamma$	TNF- $\alpha$
Placebo Day 1 (-3 h) <sup>a</sup>	2.43	1.40	3.69	9.15	21.33	8.55	0.86	4.49	1.99	33.19	9.80	35.17	3.51	10.29	2.98	1.63	3.85
Fluticasone Day 1 (-3 h) <sup>b</sup>	2.41	1.41	3.61	8.78	22.85	9.07	0.85	5.50	2.49	39.63	8.91	32.45	3.17	8.73	2.69	1.63	3.48

<sup>a</sup>Geometric means (pg/mL); <sup>b</sup>measured -3 h before the start of treatment on study Day 1.

## Acknowledgments

The authors thank the CHDR nursing staff, Dr. Maud Veselic-Charvat (Department of Pathology, Leiden University Medical Centre, Leiden, The Netherlands), lung staff for their skillful execution of the lung function tests, and Prof. Dr. J. Burggraaf for his review of the manuscript.

## Conflict of interest and funding

All authors declare no conflict of interest. This work was supported by Merck Research Laboratories, New Jersey, USA.

## References

- Diamant Z, Gauvreau GM, Cockcroft DW, Boulet LP, Sterk PJ, de Jongh FH, et al. Inhaled allergen bronchoprovocation tests. *J Allergy Clin Immunol.* 2013; 132: 1045–55. doi: 10.1016/j.jaci.2013.08.023.
- Boulet LP, Gauvreau G, Boulay ME, O'Byrne P, Cockcroft DW. The allergen bronchoprovocation model: an important tool for the investigation of new asthma anti-inflammatory therapies. *Allergy.* 2007; 62: 1101–10. doi: 10.1111/j.1398-9995.2007.01499.x.
- Bakakos P, Schleich F, Alchanatis M, Louis R. Induced sputum in asthma: from bench to bedside. *Curr Med Chem.* 2011; 18: 1415–22.
- Gauvreau GM, Watson RM, O'Byrne PM. Kinetics of allergen-induced airway eosinophilic cytokine production and airway inflammation. *Am J Respir Crit Care Med.* 1999; 160: 640–7. doi: 10.1164/ajrcm.160.2.9809130.
- Diamant Z, Timmers MC, van der Veen H, Page CP, van der Meer FJ, Sterk PJ. Effect of inhaled heparin on allergen-induced early and late asthmatic responses in patients with atopic asthma. *Am J Respir Crit Care Med.* 1996; 153: 1790–5. doi: 10.1164/ajrcm.153.6.8665036.
- Parameswaran K, Inman MD, Watson RM, Morris MM, Efthimiadis A, Ventresca PG, et al. Protective effects of fluticasone on allergen-induced airway responses and sputum inflammatory markers. *Can Respir J.* 2000; 7: 313–9.
- Choy DF, Modrek B, Abbas AR, Kummerfeld S, Clark HF, Wu LC, et al. Gene expression patterns of Th2 inflammation and intercellular communication in asthmatic airways. *J Immunol.* 2011; 186: 1861–9. doi: 10.4049/jimmunol.1002568.
- Liu L, Jarjour NN, Busse WW, Kelly EA. Enhanced generation of helper T type 1 and 2 chemokines in allergen-induced asthma. *Am J Respir Crit Care Med.* 2004; 169: 1118–24. doi: 10.1164/rccm.200312-1659OC.
- Hadjicharalambous C, Dent G, May RD, Handy RL, Anderson IK, Davies DE, et al. Measurement of eotaxin (CCL11) in induced sputum supernatants: validation and detection in asthma. *J Allergy Clin Immunol.* 2004; 113: 657–62. doi: 10.1016/j.jaci.2004.01.757.
- Grebski E, Peterson C, Medici TC. Effect of physical and chemical methods of homogenization on inflammatory mediators in sputum of asthma patients. *Chest.* 2001; 119: 1521–5.
- Kelly MM, Keatings V, Leigh R, Peterson C, Shute J, Venge P, et al. Analysis of fluid-phase mediators. *Eur Respir J Suppl.* 2002; 37: 24–39s.
- Liu MY, Xydakis AM, Hoogveen RC, Jones PH, Smith EO, Nelson KW, et al. Multiplexed analysis of biomarkers related to obesity and the metabolic syndrome in human plasma, using

- the Luminex-100 system. *Clin Chem.* 2005; 51: 1102–9. doi: 10.1373/clinchem.2004.047084.
13. From the Global Strategy for Asthma Management and Prevention, Global Initiative for Asthma (GINA). 2015. Available from: <http://www.ginasthma.org/> [cited 22 April 2015].
  14. American Thoracic Society and European Respiratory Society. ATS/ERS recommendations for standardized procedures for the online and offline measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide. *Am J Respir Crit Care Med.* 2005; 171: 912–30. doi: 10.1164/rccm.200406-710ST.
  15. Crapo RO, Casaburi R, Coates AL, Enright PL, Hankinson JL, Irvin CG, et al. Guidelines for methacholine and exercise challenge testing – 1999. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med.* 2000; 161: 309–29. doi: 10.1164/ajrccm.161.1.ats11-99.
  16. Djukanovic R, Sterk PJ, Fahy JV, Hargreave FE. Standardised methodology of sputum induction and processing. *Eur Respir J Suppl.* 2002; 37: 1–2s.
  17. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J.* 2005; 26: 319–38. doi: 10.1183/09031936.05.00034805.
  18. Boot JD, de Ridder L, de Kam ML, Calderon C, Mascelli MA, Diamant Z. Comparison of exhaled nitric oxide measurements between NIOX MINO electrochemical and Ecomedics chemiluminescence analyzer. *Respir Med.* 2008; 102: 1667–71. doi: 10.1016/j.rmed.2008.06.021.
  19. Pizzichini E, Pizzichini MM, Leigh R, Djukanovic R, Sterk PJ. Safety of sputum induction. *Eur Respir J Suppl.* 2002; 37: 9–18s.
  20. Efthimiadis A, Spanevello A, Hamid Q, Kelly MM, Linden M, Louis R, et al. Methods of sputum processing for cell counts, immunocytochemistry and in situ hybridisation. *Eur Respir J Suppl.* 2002; 37: 19–23s.
  21. Gauvreau GM, Watson RM, Rerecich TJ, Baswick E, Inman MD, O’Byrne PM. Repeatability of allergen-induced airway inflammation. *J Allergy Clin Immunol.* 1999; 104: 66–71.
  22. Bettiol J, Sele J, Henket M, Louis E, Malaise M, Bartsch P, et al. Cytokine production from sputum cells after allergenic challenge in IgE-mediated asthma. *Allergy.* 2002; 57: 1145–50.
  23. Manise M, Holtappels G, Van Crombruggen K, Schleich F, Bachert C, Louis R. Sputum IgE and cytokines in asthma: relationship with sputum cellular profile. *PLoS One.* 2013; 8: e58388. doi: 10.1371/journal.pone.0058388.
  24. Peters MC, Mekonnen ZK, Yuan S, Bhakta NR, Woodruff PG, Fahy JV. Measures of gene expression in sputum cells can identify TH2-high and TH2-low subtypes of asthma. *J Allergy Clin Immunol.* 2014; 133: 388–94. doi: 10.1016/j.jaci.2013.07.036.
  25. Diamant Z, Van der Veen H, Kuijpers EA, Bakker PF, Sterk PJ. The effect of inhaled thiorphan on allergen-induced airway responses in asthmatic subjects. *Clin Exp Allergy.* 1996; 26: 525–32.
  26. Gauvreau GM, Lee JM, Watson RM, Irani AM, Schwartz LB, O’Byrne PM. Increased numbers of both airway basophils and mast cells in sputum after allergen inhalation challenge of atopic asthmatics. *Am J Respir Crit Care Med.* 2000; 161: 1473–8. doi: 10.1164/ajrccm.161.5.9908090.
  27. Kharitonov SA, O’Connor BJ, Evans DJ, Barnes PJ. Allergen-induced late asthmatic reactions are associated with elevation of exhaled nitric oxide. *Am J Respir Crit Care Med.* 1995; 151: 1894–9. doi: 10.1164/ajrccm.151.6.7767537.
  28. Boot JD, Tarasevych S, Sterk PJ, Schoemaker RC, Wang L, Amin D, et al. Reversal of the late asthmatic response increases exhaled nitric oxide. *Respir Med.* 2005; 99: 1591–4. doi: 10.1016/j.rmed.2005.03.016.
  29. Ricciardolo FL, Sterk PJ, Gaston B, Folkerts G. Nitric oxide in health and disease of the respiratory system. *Physiol Rev.* 2004; 84: 731–65. doi: 10.1152/physrev.00034.2003.