

University of Groningen

Risk-Assessment of Esophageal Surgery

Lammerts, Rosa; van Det, Marc J.; Geelkerken, Rob H.; Kouwenhoven, Ewout A.

Published in:
Thoracic and cardiovascular surgeon reports

DOI:
[10.1055/s-0038-1660833](https://doi.org/10.1055/s-0038-1660833)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lammerts, R. G. M., van Det, M. J., Geelkerken, R. H., & Kouwenhoven, E. A. (2018). Risk-Assessment of Esophageal Surgery: Diagnosis and Treatment of Celiac Trunk Stenosis. *Thoracic and cardiovascular surgeon reports*, 7(1), e21-e23. DOI: 10.1055/s-0038-1660833

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Risk-Assessment of Esophageal Surgery: Diagnosis and Treatment of Celiac Trunk Stenosis

Rosa G.M. Lammerts^{1,2} Marc J. van Det¹ Rob H. Geelkerken³ Ewout A. Kouwenhoven¹

¹Division of Upper GI Surgery, Department of Surgery, Ziekenhuisgroep Twente, Almelo, Overijssel, The Netherlands

²Department of Internal Medicine, University Medical Center Groningen, Groningen, The Netherlands

³Division of Vascular Surgery, Department of Surgery, Medisch Spectrum Twente, Enschede, Overijssel, The Netherlands

Address for correspondence Rosa G.M. Lammerts, MD, Division of Upper GI Surgery, Department of Surgery, Ziekenhuisgroep Twente, Zilvermeeuw 1, Almelo, Overijssel 7609 PP, The Netherlands (e-mail: r.g.m.lammerts@umcg.nl).

Thorac Cardiovasc Surg Rep 2018;7:e21–e23.

Abstract

Keywords

- ▶ atherosclerosis
- ▶ endovascular procedures/stents
- ▶ except PCI
- ▶ imaging (all modalities)
- ▶ perfusion
- ▶ stents
- ▶ esophagectomy
- ▶ complications

Anastomotic leakage of the gastric conduit following surgical treatment of esophageal cancer is a life-threatening complication. An important risk factor associated with anastomotic leakage is calcification of the supplying arteries of the gastric conduit. The patency of calcified splanchnic arteries cannot be assessed on routine computed tomography (CT) scans for esophageal cancer and, as such, in selected patients with known or assumed mesenteric artery disease, additional CT angiography of the abdominal arteries with 1 mm slices is strongly encouraged. If the mesenteric perfusion is compromised in patients with resectable esophageal cancer, angioplasty procedures with stenting of the mesenteric arteries could be performed to prevent possible ischemia of the gastric conduit.

Introduction

Patients undergoing esophagectomy are at risk of morbidity and even mortality due to perioperative complications. Leakage of the anastomosis between the proximal esophagus and the gastric conduit is a common complication. An important risk factor for anastomotic leakage (AL) is ischemia of the gastric conduit, and is entirely dependent on the patency of the gastroepiploic arcade for its perfusion. On routine computed tomography (CT) scans, calcification of the aorta and its visceral branches can be detected. However, for estimating the presence of significant stenosis, a CT angiography (CTA) of the abdominal aorta and visceral branches is required.¹ This report presents two patients evaluated for esophageal cancer, both with significant calcifications in the celiac trunk detected on routine CT scanning.

Case Reports

Case 1

A male patient, age 76, was diagnosed with cT2–3N0–1M0 distal esophageal cancer. Cardiovascular risk factors included diabetes mellitus and hypertension. The routine staging CT scan of the neck, chest, and abdomen demonstrated abundant calcifications both in the descending aorta, celiac artery (CA), and the superior mesenteric artery (SMA). Surprisingly, additional CTA did not reveal significant stenosis in the CA and SMA.

After neoadjuvant chemoradiotherapy, the patient underwent a minimally invasive Ivor Lewis esophagectomy with an uncomplicated postoperative course.

Case 2

A male patient, age 65, was diagnosed with cT3–N2–M0 distal esophageal cancer. Cardiovascular risk factors included

received
January 18, 2018
accepted
May 3, 2018

DOI <https://doi.org/10.1055/s-0038-1660833>.
ISSN 2194-7635.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms






Fig. 1 Transversal standard computed tomography (CT) of the celiac trunk.



Fig. 2 (A) Sagittal computed tomography (CT) angiography; subtotal occlusion of the celiac artery. (B) Transversal CT angiography; subtotal occlusion of the celiac artery.

hypertension, diabetes mellitus type II, a transient ischemic attack, coronary artery disease (percutaneous transluminal coronary angioplasty), and morbid obesity that was treated with a gastric bypass. The staging CT scan showed severe calcifications of the CA. Additional CTA demonstrated a subtotal occlusion of the CA without significant occlusion of the SMA (→Figs. 1 and 2A and B).

The patient was discussed in the multidisciplinary working group on mesenteric ischemia and angioplasty was advised. After neoadjuvant chemoradiotherapy and 6 weeks prior to the esophagectomy, percutaneous stent angioplasty (→Figs. 3A and B) of the CA was performed. Antithrombotic drugs (carbaspirin calcium 100 mg and clopidogrel 75 mg) were prescribed. The patient underwent a total minimally invasive Ivor Lewis esophagectomy with continuation of carbaspirin calcium. After a rapid recovery, the patient was




Fig. 3 (A) Digital subtraction angiography (DSA) before stenting. (B) DSA 1 day after percutaneous angioplasty with stenting.

discharged 6 days after operation. One year postoperatively the stent was still patent.

Discussion

AL after esophageal surgery is a life-threatening complication. This case report demonstrates that risk assessment and treatment of a potentially impaired perfusion of the gastric conduit is feasible. To our knowledge, this is the first report describing stent angioplasty prior to esophagectomy in a high-risk patient for AL due to significant calcifications.

Recently, atherosclerosis of the descending aorta and celiac trunk has been identified as a strong risk factor for AL.^{2,3} The relationship between calcifications and AL likely reflects a complex pathophysiological mechanism in generalized atherosclerosis.⁴ Notwithstanding, reducing this risk factor by assessing and quantifying the grade of the celiac trunk stenosis, and treating it when necessary with modern percutaneous endovascular techniques, appears to be possible within the waiting time between chemoradiotherapy and esophagectomy. Although larger patient data sets are needed to estimate the actual risk reduction of this strategy for AL, this report shows that it is highly important to identify patients at risk. We propose to classify patients with a stenosis in the CA > 70% detected by duplex or CT scan (arterial phase), as patients at risk, and advise to perform preventive percutaneous mesenteric artery stenting.^{1,5,6}

In conclusion, patients with severe calcifications of the mesenteric arteries on routine preoperative CT scan-

ning can benefit from further assessment and treatment to reduce the risk of AL. CTA can accurately estimate the existence and the grade of a stenosis, and endovascular treatment can be performed in the waiting time for esophagectomy.

References

- 1 Björck M, Koelemay M, Acosta S, et al; Esvs Guidelines Committee. Editor's choice - management of the diseases of mesenteric arteries and veins: clinical practice guidelines of the European Society of Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg* 2017; 53(04):460–510
- 2 van Rossum PSN, Haverkamp L, Verkooijen HM, van Leeuwen MS, van Hillegersberg R, Ruurda JP. Calcification of arteries supplying the gastric tube: a new risk factor for anastomotic leakage after esophageal surgery. *Radiology* 2015;274(01):124–132
- 3 Goense L, van Rossum PSN, Weijs TJ, et al. Aortic calcification increases the risk of anastomotic leakage after Ivor-Lewis esophagectomy. *Ann Thorac Surg* 2016;102(01):247–252
- 4 Kornmann VNN, van Werkum MH, Bollen TL, van Ramshorst B, Boerma D. Compromised visceral circulation does not affect the outcome of colorectal surgery. *Surg Today* 2014;44(07):1220–1226
- 5 Mohler ER III, Gornik HL, Gerhard-Herman MD, Misra S, Olin JW, Zierler RE. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing. *J Am Coll Cardiol* 2012;60(03):242–276
- 6 Bulut T, Oosterhof-Berkas R, Geelkerken RH, Brusse-Keizer M, Stassen EJ, Kolkman JJ. Long-term results of endovascular treatment of atherosclerotic stenoses or occlusions of the coeliac and superior mesenteric artery in patients with mesenteric ischaemia. *Eur J Vasc Endovasc Surg* 2017;53(04):583–590