

## Renin-Angiotensin system (RAS) in Hematopoietic Stem/Progenitor Cells (HS/PC) Predicts Vaso-reparative Dysfunction and Progression of Diabetic Retinopathy (DR)

Yaqian Duan<sup>1,2</sup>, Eleni Beli<sup>2</sup>, Sergio Li Calzi<sup>2</sup>, Judith L. Quigley<sup>2</sup>, Rehae C. Miller<sup>2</sup>, Leni Moldovan<sup>2</sup>, Tatiana Salazar<sup>2</sup>, Sugata Hazra<sup>3</sup>, Jude Al-Sabah<sup>2</sup>, Kakarla V. Chalam<sup>4</sup>, Thao Le Phuong Trinh<sup>1,2</sup>, Ruchi J. Vyas<sup>5</sup>, Matthew C. Murray<sup>5</sup>, Patricia Parsons-Wingenter<sup>5</sup>, Gavin Y. Oudit<sup>6</sup>, and Maria B. Grant<sup>1,2</sup>

### Author Affiliations:

1. Department of Integrative and Cellular Physiology, Indiana University School of Medicine, Indianapolis, IN
2. Department of Ophthalmology, The Eugene and Marilyn Glick Eye Institute, Indiana University School of Medicine, Indianapolis, IN
3. Department of Internal Medicine, University of Utah, Salt Lake City, UT
4. Department of Ophthalmology, University of Florida, Jacksonville, FL
5. Space Life Sciences Research Branch, NASA Ames Research Center, Moffett Field, CA
6. Department of Medicine, Mazankowski Alberta Heart Institute, University of Alberta, Edmonton, Canada.

**Purpose:** We tested the hypothesis that loss of angiotensin converting enzyme 2 (ACE2) within diabetic HS/PCs would be detrimental to HS/PC reparative function, and alter their ability to contribute to vascular remodeling in human subjects and rodent models of DR.

**Methods:** Subjects (n=52) were recruited as controls (n=13) or diabetics (n=39) with either no DR, mild non-proliferative DR (NPDR), moderate NPDR, severe NPDR or proliferative DR (PDR). Fluorescein angiograms were analyzed using Vessel Generation Analysis (VESGEN) software in a cohort of subjects. CD34<sup>+</sup> HS/PCs were isolated from peripheral blood. RAS gene expression and migration was measured. Diabetic ACE2 knockout (KO)/C57BL/6-*Ins2* (Akita) mice at 3, 6 and 9 months of diabetes were compared to age-matched controls. Bone marrow HS/PC populations were analyzed by flow cytometry and migration and proliferation studies performed.

**Results:** ACE2 gene expression in human CD34<sup>+</sup> cells from diabetics without DR was increased compared to controls (p=0.0437). Mas receptor mRNA was also increased in diabetics without DR, but reduced with the onset of NPDR (p=0.0002), suggesting a loss of compensation. DR was associated with CD34<sup>+</sup> cell migratory dysfunction. By VESGEN analysis, vessel density measured by several confirming parameters in early NPDR (n=3) was greater than in normal retina (n=6) in both arteries and veins, which suggests active retinal remodeling. ACE2KO-Akita and Akita cohorts showed reduced retinal thickness by OCT at 9 months of diabetes. Absence of ACE2 in 9-month Akita mice led to an accelerated

increase in acellular capillaries compared to diabetic alone. Electroretinogram (ERG) in ACE2KO-Akita mice resulted in persistent deterioration of the neural retina. Reparative function studies showed that ACE2KO exacerbated diabetes-induced impairment of LK cell migration and proliferative functions as early as 3-month of diabetes ( $p=0.0019$ ).

**Conclusions:** Retinopathy and adverse vascular remodeling in subjects with diabetes was associated with a loss of the protective arm of RAS in HS/PCs. Loss of ACE2 exacerbated vascular dysfunction in diabetic mice.