

A Quick Summary of IMERG Versions and Features

**George J. Huffman(1), David T. Bolvin(1,2),
Dan Braithwaite(3), Kuolin Hsu(3), Robert Joyce(4,5),
Christopher Kidd(1,6), Eric Nelkin(1,2), Soroosh Sorooshian(3),
Jackson Tan(1,7), Pingping Xie(5)**

(1) NASA/GSFC Earth Sciences Division – Atmospheres

(2) Science Systems and Applications, Inc.

(3) Univ. of California Irvine

(4) Innovim

(5) NOAA/NWS Climate Prediction Center

(6) Univ. of Maryland / ESSIC

(7) Univ. Space Res. Assoc.

1. INTRODUCTION

Input precip estimates


- GPROF (LEO passive microwave [PMW])
- PERSIANN-CCS (GEO infrared)

Goal: seek the longest, most detailed record of “global” precip

IMERG is a unified U.S. algorithm that takes advantage of

- Kalman Filter CMORPH (lagrangian time interpolation) – NOAA
- PERSIANN-CCS (IR) – U.C. Irvine
- TMPA (inter-satellite calibration, gauge combination) – NASA
- PPS (input data assembly, processing environment) – NASA

GSMaP is the JAXA merged product



Ascending passes (F08 descending); satellites depicted above graph precess throughout the day.
Image by Eric Nelkin (SSAI), 14 December 2017, NASA/Goddard Space Flight Center, Greenbelt, MD.

2. IMERG DESIGN – Data Sets

Multiple runs accommodate different user requirements for latency and accuracy

- “Early” – 4 hr (flash flooding)
- “Late” – 14 hr (crop forecasting)
- “Final” – 3 months (research)

Time intervals are half-hourly and monthly (Final only)

0.1° global CED grid

- merged PMW precip 90° N-S
- morphed precip 60° N-S for now
- probability of liquid precip 90° N-S

User-oriented services by archive sites

- interactive analysis (Giovanni)
- alternate formats (TIFF files, ...)
- value-added products

New in V05

	<i>Half-hourly data file (Early, Late, Final)</i>
1	[multi-sat.] precipitationCal
2	[multi-sat.] precipitationUncal
3	[multi-sat. precip] randomError
4	[PMW] HQprecipitation
5	[PMW] HQprecipSource [identifier]
6	[PMW] HQobservationTime
7	IRprecipitation
8	IRkalmanFilterWeight
9	[phase] probabilityLiquidPrecipitation
10	precipitationQualityIndex
	<i>Monthly data file (Final)</i>
1	[sat.-gauge] precipitation
2	[sat.-gauge precip] randomError
3	GaugeRelativeWeighting
4	probabilityLiquidPrecipitation [phase]
5	precipitationQualityIndex

3. VERSION 04 IMERG – Upgrades

Use Version 04 precip from sensors using GPROF2014v2 algorithm

Reduce Final Run latency from 3.5 to 2.5 months

- change how ancillary data are handled

Shift from static to dynamic calibration of PERSIANN-CCS by PMW precip

Extend PMW gridders to 90° N-S

Reduce blockiness

- turn off volume adjustment in gauge analysis
- screen off-shore gauge influence
- spatially average 2BCMB-GMI calibrations


Correct bug that placed morphed values one gridbox south of actual location

- found thanks to a user's question

Adjust 2BCMB to the zonal-mean GPCP (land and ocean, except low-latitude ocean)

Calibrate all microwave sensors to 2BCMB

3. VERSION 04 IMERG – High-Latitude Seasons for Merged Microwave (HQ)



David Bolvin (SSAI; GSFC)

Warm-season estimates appear useful at high latitudes

Input precip estimates are still deficient in snow/ice-covered surface regions

- still screening out PMW estimates in snow/ice areas and use PMW-calibrated PERSIANN-CCS estimates

3. VERSION 05 IMERG – GPM Core Products Are Low in Extratropical Oceans

Ocean-only zonals for 2015

V05 GPM products are similar, by design


- V05 IMERG calibrated by 2BCMB at low latitudes

GPCP is higher in the extratropics

- Version 2.3 of community standard
- Behrangi Multi-satellite CloudSat, TRMM, Aqua (MCTA) product confirms GPM bias
 - includes CloudSat rain, snow, mixed
 - higher than GPCP in mid-latitudes
 - roughly agrees at high latitudes

Adjust IMERG V04, and now V05 to GPCP at higher latitudes with seasonal “climatology”

- provided reasonable IMERG bias in V04
- low biases in GPM products addressed in V05, but still low, still require GPCP



3. VERSION 05 IMERG – GPM Core Product Biases Vary by Latitude

Land-only zonals for 2015


V05 GPM products tend to show more spread

GPCP is higher in the extratropics

- V05 IMERG similar (both use GPCC gauge analysis)
- MCTA n/a over land

Adjust IMERG to GPCP for V04 and now V05 at all latitudes with a seasonal “climatology”

- first cut at the adjustment to gauges that the final calibration in IMERG enforces
- biases in GPM products addressed in V05, but still low, still require GPCP









4. VALIDATION – Accumulations over South Carolina, 1-5 October 2015

Bias decreases from V03 to V04 to V05

- the gauge-only analysis shows more than MRMS
- V03 and V04 lack the split near the coast, closer hint in V05
- V05 still puts the maximum too far inland
- IMERG higher over the ocean, but need to consider radar range artifacts for MRMS

Late V05 is not yet available

- the uncal field in Final should be approximately the same



4. VALIDATION – Half-Hourly V04 IMERG Sources and MRMS over South Carolina, 2-4 October 2015


“Violin diagram” for individual sources of the half-hourly IMERG estimates

- width shows relative contribution for each difference bin
- V03(V04) on left(right)

All rainfall rates, over land

V04 is an improvement for all sensors

No-PMW (interpolated and with IR) data are competitive with the skill for most of the sensors




4. VALIDATION – Half-Hourly V04 IMERG Sources and MRMS over South Carolina, 2-4 October 2015

This diagram focuses solely on heavy rain

- both ≥ 10 mm/h
- small sample size for AMSR2, GMI, ATMS
- V04 better than V03
- GMI and SSMIS are near zero bias
- new ATMS has issues (but low number of samples)

V05 2BCMB has a better PDF at high rates


- recall: it is the calibrator
- expect improved IMERG performance in flood situations



4. VERSION 05 IMERG – Quality Index (QI)

Half-hourly QI


- approx. Kalman Filter correlation
 - time to nearest PMWs
 - (not-morphed) IR (when used)
 - set to 1 when a PMW is used
- thin strips due to inter-swath gaps
- blocks due to regional variations
- low values at high lat. due to using IR with PMW masked out over snow



Half-Hr Qual. Index 00 UTC 3 Dec 2016

Monthly QI

- Equivalent Gauge (Huffman et al. 1997) in gauges / 2.5° x2.5°
- invert random error equation
- largely tames the non-linearity due to rain amount
- some residual issues at high values



Month Qual. Index Dec 2016

D.Bolvin (SSAI; GSFC)

5. FUTURE – Version Transitions

Early Spring 2017: Version 04, first-generation GPM-based IMERG archive, March 2014–present (Early, Late, Final)

Fall 2017: Version 05 IMERG, March 2014–present (Final; 1 December 2017-present for Early, Late, with retrospective back to March 2014 underway)

- DPR calibration change
- “minor”, but important upgrades to other algorithms
- IMERG Quality Index
- still no morphing outside 60° N-S

Spring 2018: TRMM V8/GPM V05 TRMM/GPM-based IMERG archive, 1998–present

Late Spring 2018: Legacy TMPA products retired

~2 years later: Version 06 IMERG

6. FINAL COMMENTS

Versions 04 and 05 address issues uncovered in each previous version

- swaths gridded over entire globe
- GPCP calibration in many locations
- improved input retrievals

Versions will move quickly over the next 12 months

- GPM era being upgraded to Version 05 (Final done, Early and Late in process)
- TRMM-GPM eras reprocessed in Version 05 in Spring 2018
- TMPA to be run through Spring 2018

The future holds some “interesting” challenges, technical and institutional

george.j.huffman@nasa.gov

pmm.nasa.gov



“Last Week of IMERG” at
<https://svs.gsfc.nasa.gov/cgi-bin/details.cgi?aid=4285>



On the Hyperwall Tues. 10 a.m., Wed. 4 p.m.

extra slides

3. VERSION 04 IMERG – 2BCMB Largely Behaves as Expected for Spring 2015

Low-latitude ocean not adjusted; highest latitudes still show deficits



- regional biases are modest



4. VALIDATION – Hurricane Harvey, 25-31 August 2017, IMERG and MRMS (1/2)

Harvey loitered over southeast Texas for a week

- Multi-Radar Multi-Sensor (MRMS) considered the best estimate
 - some questions about the details of the gauge calibration of the radar estimate
 - over land
- Late Run IMERG V04 under(over)-estimated in Area 1(2)



4. VALIDATION – Hurricane Harvey, 25-31 August 2017, IMERG and MRMS (2/2)

The differences between MRMS (blue) and IMERG (orange) tend to be of the same sign as the event-average difference

- less true in Area 2
- some jumpiness in IMERG is due to overpasses by different sensors
- opposite-sign differences occurred at the same time in the two areas
- PMW-calibrated IR (green) is mostly less than MRMS in both areas

This presumably tells us about the meteorology

- “juicy”, liquid-process tropical convection in Area 1
- drier, more continental convection in Area 2
- deviations from global calibration are regionally correlated

