



© 2011 The Authors
© 2011 Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas 10 (2): 155 - 158
BLACPMA ISSN 0717 7917

Artículo Original | Original Article

Chemical constituents from *Zanthoxylum setulosum* (Rutaceae)

[Constituyentes químicos de *Zanthoxylum setulosum* (Rutaceae)]

Soledad MORA¹, Víctor CASTRO¹, Luis POVEDA², Max CHAVARRÍA¹ & Renato MURILLO¹

¹Escuela de Química and CIPRONA, Universidad de Costa Rica, 2060, San José, Costa Rica. ²Escuela de Ciencias Ambientales, Facultad de Ciencias de la Tierra y el Mar, Universidad Nacional, 3000, Costa Rica.

Contactos | Contacts: Max CHAVARRIA E-mail address max.chavarría@ucr.ac.cr

Abstract

Following our phytochemical studies of Costa Rican plants, in this work we report the isolation and identification of eight compounds from aerial parts of *Zanthoxylum setulosum* (Rutaceae). They were identified as the alkaloid skimmianine, the lignans savinin, kusunokinin, sesamin, syringaresinol and the isopentenyl ether of pluviatol, the amide aurantiamide acetate, and the triterpen lupeol. This is the first report of isolation of skimmianine from the leaves of *Z. setulosum* and its presence confirm that quinoline and benzophenanthridine alkaloids, can be considered as chemotaxonomic markers of this genus. All the isolated compounds were characterized by spectroscopic methods (including ¹H-NMR, ¹³C-NMR, , HMQC, HMBC and NOESY) and comparison with the literature data.

Keywords: Rutaceae, *Zanthoxylum*, alkaloids, skimmianine, lignans.

Resumen

Continuando con el estudio fitoquímico de plantas de Costa Rica, en este trabajo informamos el aislamiento e identificación de ocho compuestos de las partes aéreas de *Zanthoxylum setulosum* (Rutaceae). Los compuestos fueron identificados como el alcaloide skimmianina, los lignanos savinina, kusunokinina, sesamina, siringaresinol y el éter isopentílico del pluviatol, la amida conocida como acetato de aurantiamida, y el triterpeno lupeol. Este es el primer informe del aislamiento de skimmianina en las hojas de *Z. setulosum*, lo cual confirma que alcaloides quinolínicos y benzofenantridínicos pueden ser considerados marcadores quimiotaconómicos en éste género. La estructura de los compuestos aislados fue caracterizada por métodos espectroscópicos (incluyendo ¹H-NMR, ¹³C-NMR, HMQC, HMBC y NOESY) y comparación con datos de la literatura.

Palabras Clave: Rutaceae, *Zanthoxylum*, alcaloides, skimmianina, lignanos.

Recibido | Received: December 18, 2010.

Aceptado en versión corregida | Accepted in revised form: February 23, 2011.

Publicado en línea | Published online: March 30, 2011.

Declaración de intereses | Declaration of interests: The financial support of Deutscher Akademischer Austauschdienst (DAAD).

Este artículo puede ser citado como / This article must be cited as: Soledad MORA, Víctor CASTRO, Luis POVEDA, Max CHAVARRÍA, Renato MURILLO. 2011. Chemical constituents from *Zanthoxylum setulosum* (Rutaceae). Bol Latinoam Caribe Plant Med Aromat 10 (2): 155 – 158.

INTRODUCTION

The *Zanthoxylum* genus that belongs to the family Rutaceae, comprises about 200 species of aromatic trees and shrubs native to the middle latitudes of North and South America, Africa, Asia, and Australia (Talapatra *et al.*, 1973). Several members of this genus are used in traditional medicine around the world. In Costa Rica there are 12 species of this genus (Instituto Nacional de Biodiversidad INBio, <http://www.inbio.ac.cr/es/default.html>) and some of them are used to cure toothache, snake bites, enteritis, diarrhea, urethritis, stomatitis, rheumatism, bronchitis and hemorrhoids (Ngane *et al.*, 2000; Islam *et al.*, 2001; Matu and van Staden, 2003; Ross *et al.*, 2004). In terms of phytochemistry, more than 90 species have been studied and among secondary metabolites that appear most frequently are alkaloids (Ahmad *et al.*, 2003; de A Gonzaga *et al.*, 2003; Jiang *et al.*, 2007; Huang *et al.*, 2008), terpenes (Mathur *et al.*, 1967) and lignans (Marcos *et al.*, 1990; Chen *et al.*, 1999; Fiorentino *et al.*, 2007; Chen *et al.*, 2008).

In this study, we describe the isolation and identification of eight compounds from aerial parts of *Zanthoxylum setulosum* (Rutaceae).

MATERIALS AND METHODS

General

Column chromatography (CC) was performed on silica gel 70-230 mesh (Merck). Thin layer chromatography (TLC) was conducted on silica gel 60 F₂₅₄ (Merck). NMR spectra were obtained on Varian Mercury 400 MHz instrument, using TMS as internal standard.

Plant material

The aerial parts of *Z. setulosum* were collected in Esparza, Alajuela province, Costa Rica, in February 2005. A voucher specimen has been deposited at the herbarium Juvenal Valerio (Heredia, Costa Rica) under reference N° 12091.

Extraction and Isolation

Air-dried and powdered aerial parts (1.35 kg of leaves) were extracted with methyl-*tert*-butyl-ether (MTBE): methanol (MeOH) 9:1 at room temperature for 24 hours. The residue (23.2 g) was obtained after removing the solvent under reduced pressure and resuspended in CHCl₃. The CHCl₃ extract was subjected to silica gel column chromatography (CC) by gradient elution with hexane: MTBE (85:15- 0:100)

and finally with MTBE:MeOH (90:10-80:20) to give seven fractions (F_a-F_g).

Fraction F_b was separated by silica gel flash CC eluting with hexane: MTBE (85:15- 0:100) to obtain twenty-five fractions (F_{b1-25}). Fraction F_{b23} was repurified by preparative TLC, eluted with benzene: MTBE 9:1 to yield sesamin (**1**, 54 mg) (Bedigian *et al.*, 1985). Fractions F_{b24} was repurified in a similar way to yield lupeol (**6**, 22 mg) (Wenkert *et al.*, 1978). Fraction F_c was separated with silica gel flash CC and eluted with hexane: MTBE (85:15- 0:100) to obtain forty fractions (F_{c1-40}). Fraction F_{c35} was repurified by preparative TLC, eluted with benzene: CH₂Cl₂ 1:1 to yield the isopentenyl ether of pluviatol (**3**, 5 mg) (Whiting, 1987). Fractions F_{c36} and F_{c38} were repurified in a similar way to give savinin (**4**, 9 mg) (Schrecker and Hartwell, 1954) and kusunokinin (**5**, 16 mg) (Lopes *et al.*, 1983) respectively.

Fraction F_d was subjected to silica gel flash CC and eluted with hexane: MTBE (85:15- 0:100) to give fifty fractions (F_{d1-50}). Fraction F_{d43} was repurified by preparative TLC, eluted with benzene: MTBE 9:1 to yield aurantiamide acetate (**7**, 34 mg) (Talapatra *et al.*, 1980).

Finally, fraction F_e was separated by silica gel flash CC eluting with hexane: MTBE (85:15- 0:100) to produce sixty fractions (F_{d1-60}). Fraction F_{e55} was repurified by preparative TLC, eluted with benzene: CH₂Cl₂: MTBE 4:4:2 to yield skimmianine (**8**, 5 mg) (Dreyer and Brenner, 1980) and syringaresinol (**2**, 14 mg) (Sharp *et al.*, 2001).

The structures of the eight compounds (figure 1), including one alkaloid, five lignans, one triterpen, and one amide were identified by spectroscopic methods (¹H and ¹³C NMR, including COSY, HMQC and HMBC) and comparison with the literature data.

RESULTS AND DISCUSSION

The plants of genus *Zanthoxylum* are well known to contain several types of compounds as amides, coumarins, flavonoids, and mainly lignans and alkaloids which show a wide range of pharmacological activities. Previous phytochemical investigations on the aerial parts of *Z. setulosum* have yielded the lignans sesamin, syringaresinol and methylarctigenin, the alkaloid kokusaginin, the sterol 22-oxo-24-methylcholest-5-en-3β-ol, the triterpen lupeol, and other components as cerotic acid, pungenin and several flavanones (Angulo Ortiz and Cuca Suarez, 2002).

In this work, we report for the first time that *Z. setulosum* is a source of skimmianine. Previous studies have shown that this alkaloid presents several biological activities e.g *in vitro* antiviral effect against hepatitis B virus (Yang and Chen, 2008), cytotoxic activity (Chen *et al.*, 2005) and antimicrobial activity

(Hanawa *et al.*, 2004). Moreover, the isolation of this compounds confirm that quinoline and benzophenanthridine alkaloids, can be considered as chemotaxonomy markers of this genus (Sun and Duan, 1996).




Figure 1. Structures of compounds 1–8 isolated from *Z. setulosum*: skimmianine (8), the lignans savinin (4), kusunokinin (5), sesamin (1), syringaresinol (2) and the isopentenyl ether of pluviatol (3), the amide aurantiamide acetate (7), and the triterpen lupeol (6).

In addition, the lignans savinin, kusunokinin, the isopentenyl ether of pluviatol, and the amide aurantiamide acetate are also report for the first time in the aerial parts of *Z. setulosum*. Finally, in agreement with the previous report by Angulo Ortiz and Cuca Suarez (2002) the lignans sesamin and syringaresinol and the triterpen lupeol were also isolated from *Z. setulosum*.

CONCLUSIONS

In this new phytochemical study of the aerial parts of *Z. setulosum*, we report the isolation and identification of eight known compounds. They were identified as the alkaloid skimmianine, the lignans savinin, kusunokinin, sesamin, syringaresinol and the isopentenyl ether of pluviatol, the amide aurantiamide acetate, and the triterpen lupeol.

ACKNOWLEDGEMENTS

We thank the financial support of Deutscher Akademischer Austauschdienst (DAAD).

REFERENCES

- Ahmad MU, Rahman MA, Huq E, Chowdhury R. 2003. Alkaloids of *Zanthoxylum budrunga*. *Fitoterapia*. 74: 191 - 193.
- Angulo Ortiz AA, Cuca Suarez LE. 2002. Nuevo esteroil y otros constituyentes de *Zanthoxylum setulosum*, Rutaceae. *Rev. Colomb. Quím.* 31: 87 - 93.
- Bedigian D, Seigler DS, Harlan JR. 1985. Sesamin, sesamol and the origin of sesame. *Biochem. Syst. Ecol.* 13: 133 - 139.
- Chen IS, Chen TL, Chang YL, Teng CM, Lin WY. 1999. Chemical constituents and biological activities of the fruit of *Zanthoxylum integrifoliolum*. *J. Nat. Prod.* 62: 833 - 837.
- Chen JJ, Fang HY, Duh CY, Chen IS. 2005. New indolopyridoquinazoline, benzo[c]phenanthridines

- and cytotoxic constituents from *Zanthoxylum integrifolium*. *Planta Med.* 71: 470 - 475.
- Chen JJ, Wang TY, Hwang TL. 2008. Neolignans, a coumarinolignan, lignan derivatives, and a chromene: anti-inflammatory constituents from *Zanthoxylum avicennae*. *J. Nat. Prod.* 71: 212 - 217.
- de A Gonzaga W, Weber AD, Giacomelli SR, Dalcol, II, Hoelzel SC, Morel AF. 2003. Antibacterial alkaloids from *Zanthoxylum rhoifolium*. *Planta Med.* 69: 371 - 374.
- Dreyer DL, Brenner RC. 1980. Alkaloids of some Mexican *Zanthoxylum* species. *Phytochemistry.* 19: 935 - 939.
- Fiorentino A, DellaGreca M, D'Abrosca B, Oriano P, Golino A, Izzo A, Zarelli A, Monaco P. 2007. Lignans, neolignans and sesquilignans from *Cestrum parqui l'Her.* *Biochem. Syst. Ecol.* 35: 392 - 396.
- Hanawa F, Fokialakis N, Skaltsounis AL. 2004. Photoactivated DNA binding and antimicrobial activities of furoquinoline and pyranoquinolone alkaloids from rutaceae. *Planta Med.* 70: 531 - 535.
- Huang HY, Ishikawa T, Peng CF, Tsai IL, Chen IS. 2008. Constituents of the root wood of *Zanthoxylum wutaense* with antitubercular activity. *J. Nat. Prod.* 71: 1146 - 1151.
- Islam A, Sayeed A, Bhuiyan MS, Mosaddik MA, Islam MA, Astaq Mondal Khan GR. 2001. Antimicrobial activity and cytotoxicity of *Zanthoxylum budrunga*. *Fitoterapia.* 72: 428 - 430.
- Jiang H, Wei-Dong Z, Yun-Heng S, Chuan Z, Lei X, Run-Hui L, Bin W, Xi-Ke X. 2007. Alkaloids from *Zanthoxylum nitidum* (Roxb.) DC. *Biochem. Syst. Ecol.* 35: 114 - 117.
- Lopes LMX, Yoshida M, Gottlieb OR. 1983. Dibenzylbutyrolactone lignans from *Viola sebifera*. *Phytochemistry.* 22: 1516 - 1518.
- Marcos M, Jimenez C, Villaverde MC, Riguera R, Castedo L, Stermitz F. 1990. Lignans and other constituents from South and central american *Zanthoxylum* species. *Planta Med.* 56: 89 - 91.
- Mathur RK, Ramaswamy SK, Rao AS, Bhattacharyya SC. 1967. Terpenoids. 108. Isolation of an oxidodiol from *Zanthoxylum rhetsa*. *Tetrahedron.* 23: 2495 - 2498.
- Matu EN, van Staden J. 2003. Antibacterial and anti-inflammatory activities of some plants used for medicinal purposes in Kenya. *J. Ethnopharmacol.* 87: 35 - 41.
- Ngane AN, Biyiti L, Zollo PH, Bouchet P. 2000. Evaluation of antifungal activity of extracts of two Cameroonian rutaceae: *Zanthoxylum leprieurii* Guill. et Perr. and *Zanthoxylum xanthoxyloides* Waterm. *J. Ethnopharmacol.* 70: 335 - 342.
- Ross SA, Sultana GN, Burandt CL, ElSohly MA, Marais JP, Ferreira D. 2004. Syncarpamide, a new antiplasmodial (+)-norepinephrine derivative from *Zanthoxylum syncarpum*. *J. Nat. Prod.* 67: 88 - 90.
- Schrecker AW, Hartwell JL. 1954. The structure of savinin. *J. Am. Chem. Soc.* 76: 4896 - 4899.
- Sharp H, Thomas D, Currie F, Bright Z, Sarker SD, Nash RJ. 2001. Pinosesinol and syringaresinol: two lignans from *Avicennia germinans* (Avicenniaceae). *Biochem. Syst. Ecol.* 29: 325 - 327.
- Sun XW, Duan ZX. 1996. Progress in the studies on medicinal plants of the genus *Zanthoxylum* Linn. *Acta Pharm. Sin.* 31: 231 - 240.
- Talapatra SK, Dutta SK, Talapatra B. 1973. Alkaloids and terpenoids of *Zanthoxylum ovalifolium*. *Phytochemistry.* 12: 729 - 730.
- Talapatra SK, Mallik AK, Talapatra B. 1980. Pongaglabol, a new hydroxyfuranoflavone, and aurantiamide acetate, a dipeptide from the flowers of *Pongamia glabra*. *Phytochemistry* 19: 1199 - 1202.
- Wenkert E, Baddeley GB, Burfitt IR, Moreno LN. 1978. Carbon-13 nuclear magnetic resonance spectroscopy of naturally-occurring substances-LVII Triterpenes related to lupane and hopane. *Org. Magn. Resonance.* 11: 337 - 343.
- Whiting DA. 1987. Lignans, neolignans, and related compounds. *Nat. Prod. Rep.* 4: 499 - 525.
- Yang G, Chen D. 2008. Alkaloids from the roots of *Zanthoxylum nitidum* and their antiviral and antifungal effects. *Chem. Biodivers.* 5: 1718 - 1722.