


1-(N-Alkoxyamino)pyridine Derivatives and others N-Alkoxyhydrazines

Shtamburg V.G.^a, Tsyhankov A.V.^b, Shtamburg V.V.^a, Klotz E.A.^c, Shishkina S.V.^d,
Zubatyuk R.I.^d, Mazepa A.V.^e

^aUkrainian State University of Chemical Technology, e-mail: stamburg@gmail.com,

^bKirovograd Flight Academy of National Aviation University, ^cKirovograd State Pedagogical University; ^dSSI "Institute for Single Crystals" of NAS of Ukraine; ^eA.V. Bogatsky Physiko-Chemical Institute of NAS of Ukraine

N-Chloro-N-alkoxycarbamates **1a-d** react with 4-dimethylaminopyridine (DMAP) yielding unknown chlorides of 1-(N-alkoxyamino)-4-dimethylaminopyridines **2a-d**. N-Chloro-N-alkoxyureas **3a-c** with DMAP form N-1-(4-dimethylamino)pyridinium-N-alkoxyureas chlorides **4a-c**. N-Chloro-N-alkoxy derivatives of benzamide **5a,b** and N-*tert*-alkylamine **8** converted in N-pyridinium-N-alkoxycompound **6,9**, respectively, by the action of DMAP. N-Chloro-N-alkoxycarbamates **1a** has been transformed in N-alkoxyhydrazines **10, 11**.



The structures of compounds **2a,b**,
discussed.

and **4a, 9-11** had been